

THE
JOURNAL
OF THE
BOMBAY NATURAL HISTORY SOCIETY

EDITED BY
SIR REGINALD A. SPENCE, KT., F.Z.S., and S. H. PRATER, C.M.Z.S.

VOL. XXXI

Parts 1 & 2

Containing 2 Coloured Plates, 52 Black and White Plates,
39 Text-figures and 6 Diagrams.

Dates of Publication.

Part I. (Pages 1 to 232) ... 15th May, 1926.
,, II. (,, 233 to 532) ... 15th August, 1926.

LONDON AGENTS
DULAU & Co., Ltd., 34—36, Margaret Street, Cavendish Square, W.

PRINTED AT THE DIOCESAN PRESS, MADRAS.

CONTENTS OF VOLUME XXXI.

No. 1.

	PAGE
THE MAMMALS AND BIRDS OF KASHMIR AND THE ADJACENT HILL PROVINCES. Part VI. (<i>With a plate.</i>) By Col. A. E. Ward.....	1
FACTS AND HYPOTHESES IN THE PROBLEM OF EVOLUTION. By E. Blatter, S.J., PH.D., F.L.S.	12
SCIENTIFIC RESULTS OF THE MAMMAL SURVEY, NO. XLV. By Oldfield Thomas, F.R.S.....	22
THREE MONTHS UP THE VALLEY OF THE SUTLEJ RIVER. (<i>With four plates.</i>) By Lt.-Col. R. W. Burton, I.A. (Retd.).....	23
BOMBAY NATURAL HISTORY SOCIETY'S MAMMAL SURVEY OF INDIA, BURMA AND CEYLON. Report No. 38. By Mrs. Helen M. Lindsay.....	40
THE IDENTIFICATION OF INDIAN BUTTERFLIES. Part IX. (<i>With a plate.</i>) By Col. W. H. Evans, D.S.O., F.Z.S.....	49
THE RETICULATE PYTHON. By Col. F. Wall, C.M.G., K.H.S.	84
ADDITIONAL NOTES ON THE AVIFAUNA OF IRAQ. (<i>With three plates.</i>) By C. B. Ticehurst, M.A., M.B.O.U., F.R.G.S.....	91
MAHSEER FISHING IN THE DECCAN LAKES. (<i>With two plates.</i>) By Major W. B. Trevenen.....	120
ON A COLLECTION OF <i>GEOMETRIDÆ</i> FROM UPPER BURMA MADE BY CAPT. A. E. SWANN. (<i>With a plate.</i>) By L. B. Prout, F.E.S.....	129
AN ORIENTAL <i>SPHEX</i> OR HUNTING WASP. Part II. By Major R. W. G. Hingston, I.M.S.	147
INDIAN DRAGONFLIES. Part XXIII. (<i>With a plate and three text figures.</i>) By Major F. C. Fraser, I.M.S., F.E.S.....	158
HUNTING STRAIGHT-HORNED MARKHOR. (<i>With three plates.</i>) By Major C. H. Stockley, D.S.O., O.B.E., M.C.....	172
NOTES ON THE SEASONAL OCCURRENCE OF RANGOON EARTH- WORMS. By G. E. Gates.....	180
BIRDS NESTING IN THE DRAS AND SURU VALLEYS. By B. B. Osmaston, C.I.E., I.F.S. (Retd.).....	186
REPORT OF THE WORK OF THE BOMBAY NATURAL HISTORY SOCIETY AND OF THE PROGRESS OF THE NATURAL HISTORY SECTION, PRINCE OF WALES' MUSEUM.....	197

	PAGE
EDITORIAL.....	207
REVIEWS—THE VEGETATION OF BURMA FROM A ECOLOGICAL STANDPOINT. By L. Dudley Stamp.....	210
INDIAN NATURE STUDIES FOR CHILDREN. By C. H. Donald, F.Z.S., M.B.O.U.....	211
A HANDBOOK OF THE BIRDS OF EASTERN CHINA. B. J. D. La Touche, C.M.Z.S., C.F.A.O.U., M.B.O.U. Part I.....	211
THE FAMILIES OF FLOWERING PLANTS. By J. Hutchinson, F.L.S.....	212
MISCELLANEOUS NOTES:—	
I. Wild Animals in Central India. By Lt.-Col. R. W. Burton, Indian Army. (Retd.).....	215
II. About Tigers and Leopards. By Brig.-Genl. R. G. Burton.....	216
III. Cannibalism amongst Panthers. By Capt. H. J. Rice, I.M.S.....	219
IV. Jungle Tragedies. By Lt.-Col. R. W. Burton.....	220
V. On the Colour of the Eye of the Gaur or Indian Bison. By A. A. Dunbar Brander.....	220
VI. Malformed Heads of Markhor and Ibex (<i>With a photo.</i>) By Lt.-Col. R. W. Burton.....	222
VII. Weights and Measurements of the Nilghiri Langur (<i>Pithecus johnii</i>). By C. Leigh, S.J.....	223
VIII. The Nesting of the Open-Bill Stork (<i>Anastomus oscitans</i>) in Purulia, Manbhum District. By Satya Churn Law.....	223
IX. Occurrence of the Sheldrake (<i>Tadorna tadorna</i>) in Bihar. By John A. Bucknell, F.Z.S., M.B.O.U.....	224
X. Birds Nesting with a Camera. A Correction. By B. B. Osmaston, I.F.S.....	224
XI. Notes on Birds uncommon in, or unrecorded from the Simla Hills. By S. Basil-Edwardes, M.B.O.U.....	224
XII. Record of a death from the bite of a Hamadryad or King Cobra (<i>Naia hanna</i>). By Ralph C. Morris (<i>a correction</i>).....	226
XIII. Notes on Snakes. By C. Leigh, S.J.....	227
XIV. A Bull Frog. (<i>Rana tigrina</i>) swallowing a Rat. By B. N. Zutshi.....	228
XV. Fishing in the Persian Gulf. By Lt.-Commander G. E. Boulton, R.N.....	228

	PAGE
XVI. On the Occurrence of the Giant Water-Bug. (<i>Belostoma indicum</i>) in Simla. By S. Basil-Edwardes, M.B.O.U.....	229
PROCEEDINGS.....	230

No. 2.

THE GAME BIRDS OF THE INDIAN EMPIRE. Part I. (Genus <i>Rallus</i>) (with a coloured plate.) By E. C. Stuart Baker, J.P., F.Z.S., F.L.S., M.B.O.U., C.F.A.O.U.....	233
SCIENTIFIC RESULTS FROM THE MAMMAL SURVEY, No. XLVI. By Helen M. Lindsay, M.A., B.Sc.....	239
THE MASON WASP (<i>Eumenes conica</i>). Part I. (With a plate and text figures.) By Major R. W. G. Hingston, I.M.S...	241
NOTES ON SOME NEW AND INTERESTING BUTTERFLIES FROM INDIA AND BURMA. Part I. (With one coloured and one black and white plate.) By Major-General H. C. Tytler, C.B., C.M.G., C.I.E., D.S.O., F.E.S.....	248
A CONTRIBUTION TO THE ORNITHOLOGY OF DELHI. Part I. (With two plates.) By the late Mr. S. Basil-Edwardes, M.B.O.U., R.A.O.U.....	261
TAKIN SHOOTING IN THE SPRING. (With a plate.) By Capt. E. Maxwell West.....	274
BIRDS NESTING WITH A CAMERA IN INDIA. Part VI. (With fourteen plates.) By Capt. R. S. P. Bates.....	277
NOTES ON A COLLECTION OF MOTHS OF THE FAMILY <i>GEOMETRIDÆ</i> MADE BY CAPT. A. E. SWANN. Part II. (With a plate.) By Louis B. Prout, F.E.S.....	308
THE COMMON BUTTERFLIES OF THE PLAINS OF INDIA. Part XXXVI. By T. R. Bell, C.I.E., I.F.S. (Retd.).....	323
THREE MONTHS UP THE VALLEY OF THE SUTLEJ RIVER. Part II. (With two plates.) By Lt.-Col. R. W. Burton, I.A. (Retd.).....	352
ON THE DOWN PLUMAGES OF SOME INDIAN BIRDS. (With a plate). By C. B. Ticehurst, M.A., M.R.C.S., M.B.O.U., F.R.G.S. (Late Capt. R.A.M.C.).....	368
BOMBAY NATURAL HISTORY SOCIETY'S MAMMAL SURVEY OF INDIA, BURMA AND CEYLON. Report Nos. 40, 41 and 42. By Mrs. Helen M. Lindsay, M.A., B.Sc.	379
INDIAN DRAGONFLIES. Part XXIV. (With two plates and five text-figures.) By Major F. C. Fraser, I.M.S.....	408
THE IDENTIFICATION OF INDIAN BUTTERFLIES. Part X. (With a plate.) By Col. W. H. Evans, D.S.O., F.Z.S., F.E.S.	427

	PAGE
ON SOME INTERESTING FEATURES OF THE FAUNA OF THE WESTERN GHATS. By S. L. Hora, D.Sc.....	447
RECENT ADDITION TO INDO-CEYLONESE COCCID FAUNA, WITH NOTES ON KNOWN AND NEW FORMS. By T. V. Ramakrishna Aiyar.....	450
A NOTE ON THE BIRDS OF KULU. By H. Whistler, F.L.S., F.Z.S.....	458
LADY (ELIZABETH) GWILLIM. By C. A. Wood, M.D., M.B.O.U.	486
SOME NOTES ON THE SECOND EDITION OF THE FAUNA OF BRITISH INDIA—BIRDS. Vols. I and II. By C. B. Ticehurst, M.A., M.R.C.S., M.B.O.U., F.R.G.S.....	490
REVIEWS :—INDIAN BIRD LIFE. By Douglas Dewar.....	500
MANUAL OF THE BIRDS OF CEYLON. By W. E. Wait.....	500
A HANDBOOK OF THE BIRDS OF EASTERN CHINA. By J. D. D. La Touche. Part II.....	501
THE AQUARIUM BOOK. By E. G. Boulenger....	502
ACCOUNT OF A PHOTOGRAPHIC EXPEDITION TO THE SOUTHERN GLACIERS OF KANCHENJUNGA IN THE SIKKIM HIMALAYAS. By N. A. Tombazi.....	503
LIFE OF PLANTS. By Sir Frederick Keeble....	504
A CORRECTION.....	505
EDITORIAL.....	506
OBITUARY :—MR. S. BASIL-EDWARDES.....	508
H. H. THE MAHARAJA OF DHAR.....	508
MISCELLANEOUS NOTES :—	
I. Measurements of Game Animals. By Major C. H. Stockley, O.C.E., M.C., D.S.O.....	509
II. Comparative Shape and Measurement of the Fore Feet in Tuskers and Tuskless Elephants. By J. L. Cameron.....	512
III. A Wonderful Sight. By R. C. Morris.....	513
IV. Tigers and Elephants. By R. K. Anderson.....	514
V. The 'Sambhur' call of Tigers. By R. C. Beadon.	515
VI. Wild Dogs in Mysore. By Capt. G. E. Windle.....	516
VII. Solitaires. By R. C. Morris.....	517
VIII. A Solitary Cow Gaur (<i>with a photo</i>). By Salim A. Ali.....	518
IX. A Cross between an Ibex and a Tame Goat (<i>with a photo</i>). By Col. A. E. Ward.....	519
X. The Nilgiri Tahr (<i>Hemitragus hylocrius</i>) (<i>with a plate</i>). By A. P. Kinloch, F.Z.S., M.B.O.U.....	520

	PAGE
XI. The 'Clicking' of the Muntjac (<i>M. vaginalis</i>). By Max Frei.....	521
XII. Measurements and Weight of a Malay Tapir. By S. H. Prater, C.M.Z.S.....	522
XIII. The Occurrence of the Pink-footed Goose (<i>Anser brachyrhynchus</i>) and the Mallard (<i>Anas boscas</i>) in Jaipur, Rajputana. By Major W. M. Logan Home.	522
XIV. On the occurrence of Pallas' Sand Grouse (<i>Syrhaptes paradoxurus</i>) within Indian limits. (With a plate.) By S. H. Prater, C.M.Z.S.	522
XV. The Occurrence of the Giant Heron (<i>Ardea goliath</i>) in the Khulna District, Bengal. By S. H. Prater, C.M.Z.S.....	523
XVI. The Rock-Horned Owl in Kashmir. (With a photo.) By B. B. Osmaston, C.I.E., I.F.S. (Retd.)....	523
XVII. Mating Habits of the Common Kite. (<i>Milvus migrans govinda</i> .) By Salim A. Ali.....	524
XVIII. Brahminy Kite (<i>Haliastur indus indus</i> .) Swimming. By S. H. Prater, C.M.Z.S.....	526
XIX. An Albino Bustard (<i>Eupodotis edwardsi</i> .) By Maharaja Kumar Shree Vijayarajji.....	526
XX. The Transport of Birds' Eggs. (With a text-figure.) By General R. M. Betham, M.B.O.U.....	527
XXI. Earth Snake (<i>Silybura</i> sp.) and Chicken. By A. P. Kinloch, F.Z.S., M.B.O.U.....	528
XXII. An Interesting Encounter with a Cobra. By W. H. Theobald.....	528
XXIII. The Occurrence of the Butterfly <i>Appias indra</i> var. <i>aristoxenus</i> in the Nilghiris. By J. Florence.....	529
XXIV. On <i>Psara phæopteralis</i> as a Pest on Grasses in South India. (With a plate.) By M. C. Cherian, B.Sc. and C. J. George, M.A.....	529
PROCEEDINGS.....	531

ALPHABETICAL LIST OF CONTRIBUTORS

VOLUME XXXI

Nos. 1 and 2.

	PAGE		PAGE
AIYAR, T. V. RAMAKRISHNA ; Recent Additions to the Indo- Ceylonese <i>Coccid</i> Fauna, with notes on known and new forms	450	BELL, T. R., C. I. E., I. F. S. (Retd.) ; The Common Butter- flies of the Plains of India (in- cluding those met with in the Hill Stations of the Bombay Presidency). Part XXXVI ...	323
ALI, SALIM A. ; A Solitary Cow Gaur (<i>with a photo</i>)	518	BETHAM, R. M., M. B. O. U. ; The Transport of Birds' Eggs. (<i>with a text-figure</i>)	527
————— ; Mating Habits of the Common Kite (<i>Milvus</i> <i>migrans govinda</i>)	524	BLATTER, E., S. J., Ph. D., F. L. S. ; Facts and Hypotheses in the Problem of Evolution ...	12
ANDERSON, R. K. ; Tigers and Elephants	514	BOMBAY NATURAL HISTORY SOCIETY'S MAMMAL SURVEY OF INDIA. Report No. XL, Gwalior State. By Helen M. Lindsay.	379
BAKER, E. C. STUART, J. P., F. Z. S., F. L. S., M. B. O. U., C. F. A. O. U. The Game Birds of the Indian Empire, Part I (<i>with a coloured plate</i>) of (<i>Rallus aquaticus indicus</i>) ...	233	BOULTBEE, Lt.-COM., G.E., R.N. ; Fishing in the Persian Gulf ...	228
BASIL-EDWARDES, The Late S., M.B.O.U. ; Notes on Birds Un- common in, or Unrecorded from the Simla Hills	224	BRANDER, A. A. DUNBAR ; On the Colour of the eye of the Gaur or Indian Bison. (<i>Bibos</i> <i>gaurus</i>)	220
————— S., M.B.O.U. ; On the Occurrence of the Giant Water-Bug. " (<i>Belostoma indi-</i> <i>cum</i>) in Simla	229	————— Solitaries ...	517
————— S., M. B. O. U., R. A. O. U. ; A Contribution to the Ornithology of Delhi, Part I (<i>with two plates</i>) ...	261	BUCKNELL, JOHN A., F. Z. S., M. B. O. U. ; Occurrence of the Sheldrake (<i>Tadorna tadorna</i>) in Bihar	224
BATES, CAPT. R. S. P. ; Birds and Nesting with a Camera in India, Part VI. (<i>with fourteen</i> <i>plates</i>)	277	BURTON, BRIGADIER-GENERAL, R. G. ; About Tigers and Leo- pards	216
BEADON, W. R. COLERIDGE ; The ' Sambhur Call of Tigers ' ...	515	BURTON, Lt.-Col., R. W. I. A. (Retd.) ; Jungle Tragedies ...	220
		————— I. A. (Retd.) Three Months up the Valley of the Sutlej River. Part I (<i>with</i> <i>four plates</i>). Part II (<i>with two</i> <i>plates</i>)	23 352

	PAGE
BURTON, Lt.-Cl., R.W. Malformed heads of Markhor and Ibex ...	222
----- Wild animals in Central India ...	215
CAMERON, I. L. ; Comparative shape and measurement of the Fore-feet in Tuskers and Tuskless Elephants ...	512
CHEESMAN, Major R. E., M. B. O. U., F. R. G. S. (See Ticehurst, C. B.).	
CHERIAN, M. C., B. A., B. Sc., D. I. C. and George, C. J., M. A. ; On <i>Psara phaeopteralis</i> as a Pest on Grasses in South India (with a plate) ...	529
CORRECTION ...	505
COX, SIR PERCY, G. C. M. G., G. C. I. E., K. C. S. I., M. B. O. U. ; (See Ticehurst, C. B.).	
SALIM, A. ALI ; Mating Habits of the Common Kite (<i>Milvus migrans govinda</i>) ...	524
EDITORIAL ...	207, 506
EVANS, COL. W. H., D. S. O., F. Z. S., F. E. S. ; The Identification of Indian Butterflies. Part IX (with a plate), Part X (with a plate) ...	49, 427
EVANS, COL. W. H. ; The occurrence of the Butterfly <i>Appias indra</i> var. <i>aristoxenus</i> in the Nilgiris, commenting on the above ...	529
FLORENCE, J. ; The Occurrence of the Butterfly <i>Appias indra</i> var. <i>aristoxenus</i> in the Nilgiris. ...	529
FRASER, MAJOR, F. C., I. M. S., F. E. S. Indian Dragonflies, Part XXIII (with a plate and three text-figures) ...	158
----- Indian Dragonflies, Part XXIV (with two plates and five text-figures) ...	408
FREI, MAX. ; The 'Clicking' of the Muntjac (<i>M. vaginalis</i>) ...	521
GATES, G. E. ; Notes on the Seasonal Occurrence of Rangoon Earthworms ...	180

	PAGE
HINGSTON, MAJOR R. W. G., I. M. S. ; An Oriental <i>Sphex</i> or Hunting Wasp, Part II ...	147
----- The Mason Wasp (<i>Eumenes conica</i>), Part I (with a plate and two text-figures) ...	241
HOME, MAJOR W. M. LOGAN. ; The Occurrence of the Pink-footed Goose and the Mallard in Jaipur, Rajputana ...	522
HOPWOOD, S. F. ; Tigers and Elephants ...	514
HORA, SUNDER LAL, D. Sc. ; On Some Interesting Features of the Fauna of the Western Ghats ...	447
KINLOCH, A. P., F. Z. S., M.B.O.U. ; The Nilgiri Tahr (<i>Hemitragus hylocrius</i>) (with a plate) ...	520
----- ; Earth Snake (<i>Silybura</i> sp.) and Chicken ...	528
LAW, SATYA CHURN ; The Nesting of the Open-billed Stork (<i>Anastomus oscitans</i>) in Purulia, Manbhum District ...	223
LEIGH, C., S. J. ; Weights and Measurements of the Nilgiri Langur (<i>Pithecus johnii</i>) ...	223
----- ; Notes on Snakes ...	227
LINDSAY, MRS. HELEN M. ; Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Report No. 38, Sind ...	40
----- M. A., B. Sc. ; Scientific Results from the Mammal Survey No. XLVI ...	239
----- ; See Bombay Natural History Society's Mammal Survey of India.	
MILROY, A. J. W. ; Comparative Shape and Measurement of the Fore-feet in Tuskers and Tuskless Elephants ...	512
MORRIS, RALPH C. ; Record of a Death from the Bite of a Hamadryad or King Cobra (<i>Naia hanna</i>) (a correction) ...	226

	PAGE		PAGE
MORRIS, RALPH C.; A Wonder- ful Sight	513	REVIEWS :—' Indian Bird Life '...	501
—————; Solitaries ...	517	' A Handbook of the Birds of Eastern China '	501
OBITUARY :—S. Basil-Edwardes.	508	' The Aquarium Book ' ...	502
—————; H.H. Sir Udaji Rao Maharaja of Dhar	508	' Account of a Photographic Expedition to the Southern Glaciers of Kanchenjunga in the Sikkim Himalayas ' ...	503
OSMASTON, B.B., C.I.E., I.F.S., (Retd.); Birds Nesting in the Dras and Suru Valleys... ..	186	' Life of Plants '	504
—————; Birds Nesting with a Camera (<i>a correction</i>)	224	RICE, CAPT. H. G., I. M. S.; Cannibalism amongst Panthers	219
—————; The Rock Horned Owl in Kashmir (<i>with a photo</i>)	523	STOCKLEY, MAJOR C. H., D.S.O., O.B.E., M.C.; Hunting Straight- horned Markhor (<i>with three plates</i>).	172
PROCEEDINGS	230, 531	—————; Mea- surements of Game Animals ...	509
PRATER, S. H., C. M. Z. S.; Measurements and Weights of a Malay Tapir	522	THEOBALD, WILLIAM H.; An Interesting Encounter with a Cobra	528
—————; On the Occur- rence of Pallas' Sandgrouse (<i>Syrnhaptes paradoxurus</i>) with- in Indian limits (<i>with a plate</i>).	522	THOMAS, OLDFIELD, F.R.S.; Scientific Results of the Mam- mal Survey, No. XLV	22
—————; The Occurrence of the Giant Heron (<i>Ardea goliath</i>) in the Khulna District, Bengal	523	TICEHURST, C.B., M.A., M.B. O.U., F.R.G.S. (Late Capt. R.A.M.C.) COX, SIR PERCY, G.C.M.G., G.C.I.E., K.C.S.I., M.B.O.U., AND CHEESMAN, MAJOR R.E., M.B.O.U., F.R. G.S. Additional Notes on the Avifauna of Iraq (<i>with 3 plates</i>),	91
—————; Brahminy Kite (<i>Haliastur indus indus</i>) Swim- ming	526	TICEHURST, C.B., M.A., M.B. O.U., F.R.G.S. (Late Capt. R.A.M.C.) (<i>with one plate</i>) ...	368
PROUT, LOUIS B., F.E.S.; On a Collection of Moths of the Family <i>GEOMETRIDÆ</i> from Upper Burma made by Captain A. E. Swann (<i>with a plate</i>) ...	129	—————; Some Notes on the Second Edition of the Fauna of British India-Birds, Vol. I and II	490
—————; Part II (<i>with a plate</i>)	307	TREVENEN, MAJOR W.B.; Mah- seer Fishing in the Deccan Lakes (<i>with two plates</i>) ...	120
REPORTS :—On the work of the Bombay Natural History Society and of the Progress of the Natural History Section, Prince of Wales' Museum ...	197	TYTLER, MAJOR-GENERAL, H.C., C.B., C.M.G., C.I.E., D.S.O., F.E.S.; Notes on Some New and interesting Butterflies from India and Burma, Part I (<i>with one coloured and one black and white plate</i>)	248
REVIEWS :—' The Vegetation of Burma from an Ecological Stand-point '	210	VIJAYARAJI, MAHARAJA KUMAR; An Albino Bustard (<i>Eupodotis edwardsi</i>)	526
' Indian Nature Studies for Children '	211		
' A Handbook of the Birds of Eastern China '	211		
' The Families of Flowering Plants '	212		

LIST OF CONTRIBUTORS

xi

	PAGE		PAGE
WALL, COL. F., C.M.G., K.H.S.;		WEST, E. MAXWELL; T a k i n	
The Reticulate Python (<i>Python</i>		Shooting in the Spring (<i>with a</i>	
<i>reticulatus</i>)	84	<i>plate</i>)	274
WARD, COL. A. E. (Retd.); The		WHISTLER, HUGH.; A Note on	
Mammals and Birds of Kash-		the Birds of Kulu	458
mir and the Adjacent Hill		WINDLE, CAPT. G.E.; Wild Dogs	
Provinces. Part VI (<i>with a</i>		in Mysore	516
<i>plate</i>)	1	WOOD, CASEY A.; Lady [Eliza-	
_____ ; A		beth] Gwillim, Artist and Orni-	
Cross between an Ibex and a		thologist	486
Tame Goat (<i>with a photo</i>) ...	519	ZUTSHI, B.N.; A Bull-Frog (<i>Rana</i>	
		<i>tigrina</i>) swallowing a Rat ...	228

LIST OF PLATES

VOLUME XXXI

No. 1

	PAGE
The Mammals and Birds of Kashmir and the adjacent Hill Provinces—	
Plate (A) Pallas's Cat (<i>Felis manul</i>) from the Nutra Valley ..	1
(B) Leopard 7' 8" from Plains of India... ..	1
Three Months up the Valley of the Sutlej River—	
Plate I—(A) Imposing Edifice at Fort Samarkot	24
(B) A Temple at Soondri	24
Plate II—(A) Two Tibetan Damsels	25
(B) Two of my Baggage-carrying Damsels and a Lad of Taranda	25
Plate III—(A) Panorama from Chini Forest	34
(B) View from Chini Forest Bungalow	34
Plate IV—(A) View from Snow Storm Camp	35
(B) The Thaman River at Rupa	35
Indian Butterflies—	
Plate <i>Ismeninae</i> , <i>Hesperiidae</i> , <i>Celaenorrhinae</i>	49
Avifauna of Iraq—	
Plate I—(A) Striated Scops Owl (<i>Otus brucei</i>). Baghdad ...	97
(B) Nest and Three Eggs of Bifasciated Lark (<i>Alaemon alaudipes cinerea</i>) on Acacia. Aquar Quf ...	97
Plate II—(A) Pelican's Eggs at Khor Milab	114
(B) Nest and Eggs of Slender-billed Gull (<i>Larus genis</i>)	114
(C) Flamingos Nests on the Khor Milab, Bubyen Islands	114
Plate III—(A) Lesser Crested Terns (<i>Sterna bengalensis bengalensis</i>)	115
(B) Colony and Eggs of Lesser Crested Tern (<i>Sterna bengalensis bengalensis</i>)	115
Mahseer Fishing in the Deccan Lakes—	
Plate I—(A) English Pattern Spoon, mounted with Single Hook and Triangle	123
(B) Hog-backed Spoon, mounted with Single Hook and Triangle	123
Plate II—(A) 3" Copper Flange Spinner, painted with Aluminium Paint	124
(B) 2½" Aluminium Reflex Minnow	124
Moths of the Family <i>Geometridae</i> from Upper Burma—	
Plate I— <i>Geometridae</i>	129, 309
Indian Dragonflies—	
Plate—Dorso-lateral views of <i>Davidius</i> , spp.	166
Hunting straight-horned Markhor—	
Plate I—(A) The Chickalli 'Tangi'	172
(B) Camp with Markhor Ground behind it	172
Plate II—(A) Markhor Ground	173
(B) The Dragon's Nostrils	173

	PAGE
Plate III—The Crest of the Main Ridge with Awal Shah and Jalpara.	177
Report on the Work of the Bombay Natural History Society and of the Progress of the Natural History Section, Prince of Wales' Museum—	
Plate I—Lammergeyer or Bearded Vulture (<i>Gypaëtus barbatus</i>)..	198
Plate II—Peacock Pheasant (<i>Polyplectron bicalcaratum</i>) ...	199
Plate III—(A) Indian Bison, Mounted in Society's Laboratory, being let down from the first floor windows, and ...	202
(B) <i>en route</i> to the Museum	202
Plate IV—Desert Country at Golamala, Karachi District, Sind, a Portion of the Group exhibited in the Prince of Wales' Museum	203
Plate V—Nesting group of Night Herons (<i>Nycticorax nycticorax</i>). Exhibited in the Bird Gallery, Prince of Wales' Museum	204
Plate VI—The Estuary Snake (<i>Hurria rynchops</i>) Photograph of a Cast exhibited in the Reptile Gallery	205
Plate VII—Photo of an Enlarged Model (X 40) of the Bed-Bug (<i>Cimex rotundatus</i>). One of a Series of Insect Models prepared in the Society's Laboratory to illustrate the case on 'Insects in Relation to Disease'	206
Plate (A) Cormorants and Cattle Egrets. Roosting-site in the Islands of Saheb Bandh, Purulia	223
(B) Nesting-site of Open-billed Stork (<i>Anastomus oscitans</i>) in the Island of Saheb Bandh, Purulia	223

No. 2

The Game Birds of the Indian Empire—	
Plate—The Indian Water Rail (<i>Rallus aquaticus indicus</i>) ...	233
The Mason Wasp. (<i>Eumenes conica</i>)—	
I—Completed Nest with Apertures of exit made by Young Wasps..	244
II—Cells	244
III—Engaged at Building her Wall	244
IV—Introducing her Eggs	244
New and Interesting Butterflies from India and Burma, Plates 1 and 4.	253, 257
The Ornithology of Delhi—	
Plate—(A) Babool Trees along the Muttra Road; Haunts of <i>Argya malcolmi</i> , <i>Phoenicurus o. phoenicuroides</i> ; <i>Pericrocotus p. peregrinus</i> ; <i>Lanius vittatus</i> , etc. ...	262
(B) Babool Jungle at Raisina, Delhi; Haunts of a Variety of Small Birds	262
Plate—(A) The Canal at Okla, Delhi; Haunts of <i>Ceryle e. leuc melanura</i> , <i>Ardeola grayii</i> , etc.; Clump of Trees in Background—Haunts of <i>Cuncuma leocorypha</i> ...	263
(B) A Portion of the Ridge at Raisina, Delhi; Haunts of <i>Argya c. caudata</i> , <i>Franklinia buehanani</i> , <i>Lanius e. lahtora</i> , <i>Uroloncha malabarica</i> , <i>Cyrtostomus a. asiaticus</i> .	263
Takin Shooting in the Spring—	
(A) Typical Takin Country	274
(B) The End of a Perfect Morning	274

	PAGE
Birds Nesting with a Camera in India—	
Plate I—(A) The Brown Rock-chat (<i>Cercomela fusca</i>)	278
(B) The Large Grey Babbler (<i>Argya malcolmi</i>)	278
(C) The Ceylon Red-vented Bulbul (<i>Molpastes haemorrhous haemorrhous</i>)	278
(D) The Southern Indian Stone-Chat (<i>Saxicola caprata atrata</i>)	278
Plate II—(A) The White-spotted Fantail Fly-catcher (<i>Rhipidura pectoralis</i>)	279
(B) The Southern Red-whiskered Bulbul (<i>Otocompsa emeria fuscicaudata</i>)	279
(C) The Nilgiri Laughing Thrush (<i>Trochalopteron cachinans cachinans</i>)	279
Plate III—(A) The White-eared Bulbul (<i>Molpastes leucogenys leucotis</i>)	280
(B) The Central Indian Red-vented Bulbul (<i>Molpastes haemorrhous pallidus</i>)	280
Plate IV—(A) The Southern Indian Black Bulbul (<i>Microscelis psaroides ganeesa</i>)	281
(B) The Southern Red-whiskered Bulbul (<i>Otocompsa emeria fuscicaudata</i>)	281
Plate V—(A) The Dusky Horned Owl (<i>Bubo coromandus</i>)	282
(B) The Smaller White Scavenger Vulture (<i>Neophron percnopterus ginginianus</i>)	282
Plate VI—(A) The Indian Purple Sunbird (<i>Leptocoma asiatica asiatica</i>)	283
(B) The Malabar Small Minivet (<i>Pericrocotus peregrinus malabaricus</i>)	283
Plate VII—The Southern Jungle Crow (<i>Corvus coronoides culminatus</i>)	284
Plate VIII—The Nilgiri Blackbird (<i>Turdus merula simillimus</i>)	285
Plate IX—(A) The Black-throated Weaver-Bird (<i>Ploceus bengalensis</i>)	286
(B) The Indian White-backed Vulture (<i>Pseudogyps bengalensis</i>)	286
Plate X—(A) The White-spotted Fantail Fly-catcher (<i>Rhipidura pectoralis</i>)	287
(B) The Spotted Babbler (<i>Pellorneum ruficeps ruficeps</i>)	287
Plate XI—(A) The Ashy Wren-warbler (<i>Prinia socialis socialis</i>)	290
(B) The Ceylon Wren-warbler (<i>Prinia inornata jerdoni</i>)	290
Plate XII—(A) and (B) The Ashy Wren-warbler (<i>Prinia socialis socialis</i>)	291
Plate XIII—(A) The Ceylon Hoopoe (<i>Upupa epops ceylonensis</i>)	306
(B) The Southern Grey Tit (<i>Parus major mahrattarum</i>)	306
Plate XIV—The Black and Orange Fly-catcher (<i>Ochromela nigrorufa</i>)	307
Three Months up the Valley of the Sutlej River—	
Plate (A) A Fine Burhel, 26½"	308
(B) In the Tadong Gud—' River Ravine and Splintered Precipice	309

	PAGE
Plate (C) The Timorous Gargaram crossing the Sutlej ...	361
(D) Three Handsome Men of Charong ...	361
Plate—The Down-plumages of Some Indian Birds ...	369
Indian Dragonflies—	
Plate I—Lateral views of <i>Burmagomphus</i> , spp. ...	410
Plate II—Lateral views of 1. <i>Davidioides martini</i> , ♂; 2. <i>Anisogomphus occipitalis</i> , ♂; 3. <i>Anisogomphus caudalis</i> , ♂; 4. <i>Temnogomphus bivittatus</i> , ♂; 5. <i>Platygomphus dolabratus</i> , ♂; 6. <i>Platygomphus fecæ</i> , ♂; 7. <i>Anisogomphus orites</i> , ♀ ...	422
The Identification of Indian Butterflies—	
Plate XXXI.— <i>Hesperiidæ</i> :— <i>Celænorrhinæ</i> , <i>Hesperiinæ</i> <i>Pamphilinæ</i>	430
The Nilghiri Tahr (<i>Hemitragus hylocrius</i>)—	
Plate (A) A Good Head in the Society's Collection, R. = $16\frac{1}{2}$ "	
L = $16\frac{1}{4}$ " ...	520
(B) Young Female Nilgiri Tahr ...	520
(C) Pallas' Three-toed Sandgrouse (<i>Syrrhaptes paradoxurus</i>).	522
Plate— <i>Psara phaeopteralis</i> as a Pest on Grasses in South India ...	529

INDEX TO ILLUSTRATIONS

VOLUME XXXI

Nos. I and 2

	PAGE		PAGE
<i>Alæmon alaudipes cinerea</i> , Pl.		<i>Burmagomphus laidlawi</i> , Pl. i, fig. 5.	
fig. B	97	Lateral view of ♂	410
<i>Amblypodia maymyoica</i> , Pl. fig. 19	257	Fig. 7. Genitalia... ..	418
———— <i>suffusa atarana</i> , Pl.		———— <i>pyramidalis</i> , fig.	
fig. 13	257	Wings	409
<i>Anastomus oscitans</i> , Pl. fig. B ...	223	Pl. i, fig. 4. Lateral view ...	410
<i>Angus Bull</i> , Transverse section		Fig. 4. Genitalia... ..	418
of the eye, fig.	221	———— <i>siamensis</i> , Pl. i, fig.	
<i>Anisogomphus caudalis</i> , fig. 12.		1. Lateral view of ♀	410
Genitalia	418	———— <i>sivalikensis</i> , Pl. i,	
Pl. ii, fig. 3. Lateral view of		fig. 4. Lateral view of ♂ ...	410
male	422	Fig. 1. Genitalia... ..	419
Fig. 5 <i>b</i> and appendage of male		———— <i>vermiculatus</i> , Pl. i,	
seen from above	424	fig. 2. Lateral view of ♂ ...	410
———— <i>occipitalis</i> , fig. 5.		———— <i>v. flavum</i> , Pl. i, fig.	
Genitalia	418	7. Lateral view of ♀	410
Fig. 4. Wings of male... ..	420	Burmese Moths :—	
Pl. ii, fig. 2. Lateral view of		<i>Geometridæ</i> Pl.	129
male	422	<i>Celænorhincæ</i> , spp. Pl.xxxi. Wings	430
Fig. 5 <i>c, d</i> . Anal appendage of		<i>Cercomela fusca</i> , Pl. i, fig. 1 ...	278
male seen from side or above	424	<i>Cervus, f. megaceros</i> , fig. ...	222
———— <i>orites</i> , Pl. ii, fig. 7.		<i>Cimex rotundatus</i> , Pl.	206
Lateral view of female... ..	422	<i>Cleora coniozona</i> , Pl. ii, fig. 3 ...	309
<i>Aquila vindhiana, preplumula</i> ,		———— <i>cryptogonia</i> , Pl. ii, fig. 7... 309	
Pl. fig. B.	369	<i>Corvus coronoides culminatus</i> , Pl.	
<i>Argya malcolmi</i> , Pl. 1, fig. 2 ...	278	vii	284
<i>Bibos gaurus</i> , Pl.	202	<i>Ctenognophos methoria</i> , Pl. ii,	
<i>Boarmia virguncula</i> , Pl. ii, fig. 4	309	fig. 12	309
<i>Bubo coromandus</i> , Pl. v, fig. 1 ...	282	<i>Cyclogomphus ypsilon</i> , fig. Wings	158
<i>Bubulcus coromandus</i> , Pl. fig. A.	223	<i>Davidioides martini</i> , fig. 6. Geni-	
<i>Bug, Bed</i> . Photo of an enlarged		talia	418
model (to illustrate the case on		Pl. ii, fig. 1. Lateral view ...	422
Insects in Relation to Disease,		<i>Davidius aberrans</i> , Pl. fig. 1.	
Pl.	206	Dorso-lateral view	166
<i>Burmagomphus cauvericus</i> , Pl. i,		———— fig. Dorsal	
fig. 6. Lateral view of ♂	410	view of anal appendages, ♂ ...	168
Fig 3. Genitalia	418	———— <i>dauidi assamensis</i> , Pl.	
———— <i>hasimaricus</i> , fig. 2.		fig. 2. Dorso-lateral view ...	166
Genitalia	418		

	PAGE
<i>Davidius assamensis</i> , fig. Ventral processes seen from the side, ♂	168
----- fig. 3.	
Genitalia, ♂	168
----- <i>davidi</i> , Pl. fig. 6.	
Dorso-lateral view	166
----- <i>kumaonensis</i> , Pl. fig. 7.	
Dorso-lateral view	166
----- <i>malloryi</i> , Pl. fig. 5.	
Dorso-lateral view	166
----- fig. Dorsal view of anal appendages, ♂	168
----- fig. Lateral view of anal appendages, ♂	168
----- fig. Genitalia, ♂	168
----- <i>zollorensis delineatus</i> , fig. 4,	164
----- Pl. fig. 4. Dorso-lateral view	166
----- fig. Dorsal view of anal appendages, ♂	168
----- fig. Lateral view of anal appendages, ♂	168
----- fig. Genitalia, ♂	168
----- <i>zallorensis</i> , Pl. fig. 3. Dorso-lateral view	166
Delhi :—Babool Trees along the Muttra road, Pl. fig. 1.	262
Babool Jungle at Raisina, Pl. fig. 2	262
Canal at Okla, Pl. fig. 1	263
Portion of ridge at Raisina, Pl. fig. 2	263
Down Plumages of some Indian Birds, Pl.	369
Dragon Flies, Indian Pl.	166
----- Figs. 158, 164, 168	
<i>Ectropis æthreogenes</i> , Pl. ii. fig. 18	309
----- <i>chrysotenecta</i> , Pl. ii. fig. 22	309
----- <i>embolochroma</i> , Pl. ii. fig. 8	309
----- <i>hiulca</i> , Pl. ii, fig. 17	309
----- <i>lophomeris</i> , Pl. ii, fig. 21.	309
----- <i>pulvicopia</i> , Pl. ii, fig. 6.	309
----- <i>zotica</i> , Pl. ii, fig. 20	309

	PAGE
<i>Elaphas</i> , figs. Hindfoot, fore-foot	513
<i>Enantiodes consanguinea</i> , Pl. ii, fig. 10.	309
<i>Erebia mani kamriana</i> , Pl. i, fig. 12.	257
----- <i>lorimeri</i> , Pl. i, fig. 10.	257
----- <i>shandura</i> , Pl. i, fig. 9.	257
----- <i>yasina</i> , Pl. i, fig. 11 ..	257
<i>Eumenes conica</i> , figs.	241
----- Pl. 1. Completed nest, 2. Cells, 3. engaged at building wall, 4 introducing eggs	244
<i>Felis manul</i> , Pl. fig. A	1
----- <i>pardus</i> , Pl. fig. B	1
<i>Gaur</i> , photo of skull	518
<i>Gypaëtus barbatus</i> , Pl.	198
<i>Hemitragus hylocrius</i> , Pl.	520
Hérons, Night, Nesting groups of, Exhibited on the Bird Gallery, Prince of Wales' Museum	204
<i>Hesperinæ</i> , spp. Pl. xxxi. Wings.	430
<i>Heterolocha segregis</i> , Pl. ii, fig. 2.	309
<i>Hirasa plagiochorda</i> , Pl. ii, fig. 13	309
<i>Hurria rynchops</i> , Pl.	205
Ibex, Malformed head of, fig.	222
----- X tame goat, Photo	519
Indian Butterflies :—	
Plate. Wings of <i>Ismeninæ</i> , <i>Hesperiidæ</i> , <i>Celænorhina</i>	49
<i>Indogomphus longistigma</i> , fig. 10.	
Genitalia	418
----- <i>martini</i> , fig. 9.	
Genitalia	418
Karachi District. Sind desert country at Golamala. One portion of group exhibited in the Prince of Wales' Museum, Pl....	203
<i>Krananda orthotmeta</i> , Pl. iii, fig. 1	309
<i>Larus genei</i> . Nest and eggs, Pl. fig. B	114
<i>Leptocoma asiatica asiatica</i> , Pl. vi, fig. 1	283
<i>Loxotephria perileuca</i> , Pl. ii, fig. 5	309
<i>Lycæna astorica</i> , Pl. fig. 3	257
----- <i>baroghila</i> , Pl. fig. 1	257
----- <i>christophi lesliei</i> , Pl. fig. 2	257
----- <i>florenciæ</i> , Pl. fig. 4	257

	PAGE		PAGE
<i>Lycaena galathea depreei</i> , Pl. i, fig. 5	257	<i>Otocompsa emeria fuscicaudata</i>	
— <i>metallica chitralensis</i> ,		Pl. iv, fig. 2	281
Pl. i, fig. 6	257	<i>Otus brucei</i> , Pl. fig. A	97
— <i>gilgitica</i> , Pl. i,		Owl, Rock Horned, Photo of	
fig. 7	257	3 young in nest	523
— <i>orbitulus astorica</i> , Pl. i,		<i>Pamphilinæ</i> spp., Pl. xxxi. Wings	430
fig. 3	257	<i>Parnassius delphius florenciæ</i> ,	
Mahseer Fishing in the Deccan		Pl. iv, fig. 9	253
Lakes—		— <i>rileyi</i> Pl. iv,	
Plate 1, fig. A. English pattern		fig. 6, 7	253
Spoon	123	— <i>zogilaica</i> , Pl.	
Fig. B. Hog-backed		iv, fig. 8	253
spoon	123	— <i>discobolus baroghila</i> ,	
Plate 2, fig. A. 3" copper		Pl. iv, fig. 1	253
hange spinner	124	— <i>hunzaica</i> , Pl.	
Fig. B. 2½" alumi-		iv, fig. 2	253
um reflex minnow	124	— <i>dongalaica</i> Pl. iv, fig. 5.	253
Markhor, Hunting Straight-horned—		— <i>jacquemontii shandura</i> ,	
Plate 1, fig. A. Chickalli		Pl. iv, figs. 3, 4	253
'Tangi'	172	— <i>simo lovimeri</i> , Pl. iv, fig 10	253
Fig. B. Camp with		<i>Parus major mahrattarum</i> , Pl. xiii,	
markhor ground		fig. 2	306
behind it	172	<i>Pelecanus onocrotalus</i> , eggs, Pl. fig.	
Plate 2, fig. A. Markhor		A.	114
ground	173	<i>Pellorneum ruficeps ruficeps</i> , Pl.	
Fig. B. The Dragon's		x, fig. 2	287
nostrils	173	<i>Pericrocotus pericrocotus malaba-</i>	
Plate 3, Crest of the main		<i>ricus</i> , Pl. vi, fig. 2	283
ridge	177	<i>Phalacrocorax javanicus</i> , Pl. fig.	
Markhor, Malformed head of, fig.	222	A.	223
<i>Medasina lasiochora</i> , Pl. ii, fig. 11	309	<i>Phœnicopterus ruber antiquorum</i> ,	
<i>Micrabraxas anisonoma</i> , Pl. ii,		Nests, Pl. fig. C.	114
fig. 9	309	<i>Platygomphus dolabratus</i> , fig. 2.	
<i>Microscelis psaroides ganeesa</i> , Pl.		wings of male	416
iv, fig. 1	281	— fig. 11.	
<i>Molpastes hæmorrhous hæmorrhous</i> ,		Genitalia	418
Pl. 1, fig. 3	278	— P. ii.	
— <i>pallidus</i> ,		fig. 5. Lateral view of	
Pl. iii, fig. 2	280	male	422
— <i>leucogenys leucotis</i> , Pl		— <i>feae</i> , P. ii, fig. 6.	
iii, fig. 1	280	Lateral view of male	422
<i>Neophron percnopterus ginginea-</i>		<i>Ploceus bengalensis</i> , Pl. ix, fig. 1	286
<i>nus</i> , Pl. v, fig. 2...	282	<i>Polyommatus iolas gilgitica</i> , Pl. i,	
<i>Nycticorax nycticorax</i> , Pl.	204	fig. 16	257
<i>Ochroinela nigrorufa</i> , Pl. xiv	307	— <i>sarta rupala</i> , Pl. figs.	
<i>Ocelophora agana</i> , Pl. ii, fig. 16	309	14, 15	257
<i>Orsonoba æthocrypta</i> , Pl. ii, fig.		<i>Polyplectron bicalcaratum</i> , Pl.	199
14	309	<i>Prinia inornata jerdoni</i> , Pl. xi.	
<i>Orthomella putaoica</i> , Pl. fig. 20	257	fig. 2	290
<i>Otocompsa emeria fuscicaudata</i> ,		— <i>socialis socialis</i> , Pl. xi, fig. 1	290
Pl. ii, fig. 2	279	and Pl. xii.	291

	PAGE
<i>Psara phaeopteralis</i> , P. Life-history	529
<i>Pseudogyps bengalensis</i> . Pl. ix, fig. 2 ...	286
<i>Pterocles senegalensis</i> , dorsal view of downy chick, Pl. fig. iii. ...	369
<i>Pyrrhulanda grisea</i> , down-plumage and juvenile feather Pl. fig. A	369
<i>Rallus aquaticus indicus</i> Pl. ...	233
<i>Rapala rubida</i> , Pl. fig. 18 ...	257
<i>Rhipidura pectoralis</i> , Pl. ii, fig. 1	279
----- Pl. x, fig. 1	287
<i>Saxicola caprata atrata</i> , Pl. i, fig. 4	278
<i>Semiothisa ageta</i> , Pl. ii, fig. 15 ...	309
Snake, Estuary photograph of a cast, exhibited in the Reptile Gallery, Pl.	205
<i>Sterna bengalensis bengalensis</i> , Pl. figs. A, B	115
<i>Streptopelia meena</i> , down plum- age and featherbud, Pl. fig. D	369
Sutlej Valley—	
Pl. 1, fig. A. Edieice at fort Samarkot ...	24
Fig. B. Temple at Soondri	24
Pl. 2, fig. A. Two Tibetan damsels ...	25
Fig. B. Two baggage car- rying damsels and a lad of Taranda ...	25
Pl. 3, fig. A. Panarama from Chini Forest ...	34
Fig. B. View from Chini Forest Bungalow	34

	PAGE
Pl. 4. fig. A. View from snow storm camp ...	35
Fig. B. Thaman River at Rupa ...	35
Sutlej, Three months up the Valley of the—	
A fine burhel 26½", Pl. i, fig. 1	360
In the Tadong Gud, Pl. i, fig. 2	360
The Timorous Gangaram cross- ing the Sutlej, Pl. ii, fig. 1 ...	361
Three handsome men of Cha- rong, Pl. ii, fig. 2	361
<i>Syrrhaptis paradoxurus</i> , Pl. ♂ and ♀	522
<i>Sysstema aulotis</i> , Pl. ii, fig. 19 ...	309
Takin shooting :—next line Typi- cal Takin country, Pl. fig. 1... ..	274
End of a perfect morning, Pl. fig. 2	274
<i>Temnogomphus bivittatus</i> , fig. 8. Genitalia	418
Plate ii, fig. 4 Lateral view of male	422
Fig. 5 (a) Anal appendage of male seen from above ...	424
<i>Trochalopteron cacinans cachi- nans</i> , Pl. ii, fig. 3	279
<i>Turdus merula similimus</i> , Pl. viii	285
<i>Upupa epops ceylonensis</i> , Pl. xiii, fig. 1	306
<i>Virachola dhoertyi</i> , Pl. i, fig. 17...	257

ERRATA

Volume XXXI, No. 1

Page 112, line 19 for *Burhinius* read *Burhinus*.
 „ 188 „ 51 „ *leucocephala* read *leucocephala*.
 „ 190 „ 48 „ *Cyanosyloia* read *Cyanosylvia*.
 „ 192 „ 43 „ *albitrons* read *atrifrons*.
 „ 194 „ 10 „ *latteral* read *lateral*.
 „ „ „ 22 „ *bonapartci* read *bonapartei*.
 „ 195 „ 24 „ *Bulfinch* read *Bullfinch*.
 „ 203 „ 35 „ J. P. Mills read T. P. M. O'Callaghan.

No. 2

Page 235 line 37 for mounted read moulted.
 „ 250 „ 4 „ *Shandun* read *Shandur*.
 „ „ „ 21 „ „ „
 „ „ „ 32 „ „ „
 „ 251 „ 19 „ *nicerillei* read *nicevillei*.
 „ 264 „ 23 „ *Dicurus* read *Dicrurus*.
 „ 269 „ 22 „ *phillipinus philippinus* read *phillippinus philip-*
pinus.
 „ 270 „ 34 „ *per* read *par*.
 „ „ „ 48 „ *Ptyonoprogne* read *Ptyonoprogne*.
 „ 371 „ 20 „ *Rhiphidura* read *Rhipidura*.
 „ 434 „ 7 „ the Tring types read these particular Tring
 types.
 „ 479 „ 7 „ *Hicætus* read *Hierætus*.
 „ 484 „ 47 „ *indicius* read *indicus*.
 „ 491 „ 3 „ *Phyrrhrocorax* read *Pyrrhocorax*.
 „ 494 „ 4 „ *Acreuthornis* read *Arceuthornis*.
 „ 522 „ 14 „ *Anas* read *Anser*.
 „ „ „ 15 „ *brachyrynchus* read *brachyrhynchus*.
 „ „ „ 21 „ „ „
 „ „ „ 28 „ *onocrocotalus* read *onocrotalus*.
 „ „ „ 33 „ *paradoxurus* read *paradoxus*.
 „ „ „ 35 „ „ „

Legend on plate—

Page 524 line 15 „ *Osmatson* read *Osmaston*.
 „ 526 „ 44 „ *Eupoditis* read *Eupodotes*.

BOMBAY NATURAL HISTORY SOCIETY

OFFICE BEARERS, 1927

Patron.

H.E. The Viceroy of India, G.M.S.I., G.M.I.E., H.R.H. The Prince of Wales, K.G.

Vice-Patrons.

H.H. The Maharao of Cutch, G.C.S.I., G.C.I.E.
H.H. The Maharaja of Jodhpur, K.C.S.I., K.C.V.O.
H.H. The Maharaja of Rewa, K.C.S.I.
H.H. The Maharani of Dhar.
Mr. F. V. Evans, Haymans Green, Liverpool.

President.

H.E. The Right Hon'ble Lt.-Col. Sir Leslie Wilson, P.C., G.C.I.E., C.M.G., D.S.O.

Vice-Presidents.

H.H. The Maharao of Cutch, G.C.S.I., G.C.I.E.
Rev. E. Blatter, S.J., Ph.D., F.L.S.
The Hon'ble Mr. J. E. B. Hotson, C.S.I., I.C.S.

Managing Committee.

Mr. R. D. Bell, C.I.E., I.C.S.	Bombay.
Mr. H. A. W. Brent	"
Prof. V. N. Hate, M.A.	"
Mr. J. G. Kidland	"
Mr. P. M. D. Sanderson, F.Z.S.	"
Lt.-Col. F. P. Mackie, I.M.S.	"
Mr. J. B. Greaves (<i>Honorary Treasurer</i>)	"
Sir Reginald Spence, <i>Kt.</i> (<i>Honorary Secretary</i>)	"
Mr. T. Bainbrigge Fletcher, F.E.S.	Pusa.
Mr. T. R. Bell, C.I.E., I.F.S. (Retd.)	Karwar.
Col. W. H. Evans, R.E., C.I.E.	Quetta.
Lt.-Col. F. C. Fraser, I.M.S.	Waltair.
Dr. F. H. Gravely, D.Sc.	Madras.
Mr. C. M. Inglis, M.B.O.U., C.M.Z.S.	Darjeeling.
Lt.-Col. C. H. Stockley, D.S.O., M.C., O.B.E.	Shanghai.
Major R. B. Seymour-Sewell, I.M.S.	Calcutta.

LIST OF MEMBERS

LIFE MEMBERS

Aga Khan, H. H. Aga Sir Sultan Mahomed Shaha (G.C.S.I., G.C.I.E.)	Bombay.
Aitken, C. E.	Bombay.
Alwar, Lt.-Col. H. H. Sawai Maharaja Sir Jaisingh Bahadur (K.C.S.I., G.C.I.E.)	Alwar.
Armstrong, Dr. J. S. (R.A.M.C.)	New Zealand.
Bahawalpur, H.H. The Nawab Capt. Sir Sadiq Moham- mad Khan Abassi (K.C.V.O.)	Bahawalpur.
Baker, E. C. Stuart (F.Z.S.)	England.
Balkrishna Vinayek Wassoodeo	Bombay.
Bansda, Maharaul Shri Indra Sinhji Pratap Sinhji, Raja Saheb of	Bansda.
Baroda, H.H. The Maharaja Sir Sayaji Rao Gaekwar (G.C.S.I., K.C.S.I.)	Baroda.
Barton, E. L.	England.
Barwani, Capt. H. H. Rana Ranjitsinghji Bahadur (K.C.S.I.)	Barwani.
Bikaner, Col. H.H. The Maharaja Sir Gunga Singhji Bahadur (G.C.S.I., G.C.I.E., G.C.V.O.)	Bikaner.
Bikaner, Capt. H.H. Maharaj Kumar Shri Sadul Singhji Bahadur (C.V.O.)	Bikaner.
Bikaner, Maharaj Shri Bijey Singhji Bahadur	Bikaner.
Bodenham, Capt. K. L.	England.
Bradshaw, J. P.	Bombay.
Bridgeman, Lt.-Col. The Hon'ble H. G. O. (R.F.A.)	Birmingham.
Brook, Carlton P.	England.
Burton, Brig.-Gen. R. G.	England.
Burton, Lt.-Col. R. W.	England.
Bute, The Most Hon'ble The Marquis of	Scotland.
Butler, His Excellency Sir Harcourt (G.C.I.E., K.C.S.I., I.C.S.)	Burma.
Caccia, Major A. M. (I.F.S.)	England.
Cambay, H.H. Nawab Hussain Yavarkhan Jafar Ali- khan Saheb Bahadur	Cambay.
Carter, Major H. St. M. (D.S.O., R.A.M.C.)	England.
Cassamali Jairajbhoy Peerbhoy	Bombay.
Christopher, Sydney A. (Bar-at-law)	Rangoon.
Clarke, L. O. (I.C.S.)	England.
Clutterbuck, J. E. (R.E.)	Bhurkunda.
Clutterbuck, Sir Peter H.	England.
Cooch Behar, Maharaj Kumar Victor N. Narayan	Cooch Behar.
Coode, John M.	Ootacamund.
Cowie, Rev. A. G. G.	Rawalpindi.
Cox, Lt.-Genl. Sir Percy (G.C.I.E., K.C.S.I., K.C.M.G., K.C.I.E., F.Z.S., F.R.G.S.)	England.

Craw, H. H. (I.C.S.)	Rangoon.
Crawford, W. M. (F.E.S., I.C.S.)	England.
Culbertson, J. M. S. (A. M. I.C.E.)	Karachi.
Cursetji, Khan Bahadur C. M.	Bombay.
Cutch, H. H. The Maharao Sir Shri Kengurji Sawai Bahadur (G.C.I.E., G.C.S.I.) (<i>Vice-Patron and Vice-President</i>)	Bhuj-Cutch.
Cutch, Kumar Shri Godji	Cutch.
D'Almeida, J. F. R. (B.A., B.Sc. (Hon.)	Bandra.
Davidson, Lt.-Col. J. (D.S.O., I.M.S.)	Edinburgh.
Dhar, Her Highness Laxmibai Saheb Puar (<i>Vice-Patroness</i>)	Dhar.
Dharampur, H. H. The Maharana Shree Vijayadevji Rana, Maharaja Saheb of	Dharampur.
Dhrangadhra, H.H. Maharana Shri Sir Ghanshyam Sinhji (K.C.S.I.)	Dhrangadhra.
Dhunjibhoy Bomanji, Sir, <i>Kt.</i>	Bombay.
Drake-Brockman, Lt.-Col. H. E. (F.Z.S., I.M.S.)	England.
Duxbury, Lt.-Col. C. D.	Poona.
Elgee, Colonel J. W. L.	England.
Ellison, Bernard C. (C.M.Z.S., F.R.G.S.)	Bhopal.
English, E. E.	Cairo.
Erb, E.	Bombay.
Evans, F. V. (<i>Vice-Patron</i>)	England.
Fairbank, Rev. H.	Ahmednagar.
Foulkes, Lt.-Col. T. H. (I.M.S.)	Europe
Frei, Max	Switzerland.
Frere, Major A. G. (I.A.)	Cannanore.
Gammie, Prof. G. A.	England.
Gauripur, Raja Prabhatchandra Barna	Gauripur.
Gharpurey, Lt.-Col. K. G. (I.M.S.)	Sholapur.
Gonsalves, A. F.	Bandra.
Gregory-Smith, Capt. H. G.	England.
Hide, P.	Indore.
Hill, Major R. D. O.	England
Holkar, H. H. Tukuji Rao	Indore.
Hopwood, S. F. (I.F.S.)	England.
Hotson, Hon'ble Mr. J. E. B. (C.S.I., I.C.S.)	Bombay.
Hoyos, Count E.	Austria.
Husbands, Major H. W. S. (M.C., A.M.I.C.E., S.M.R.E.)	England.
Hyam, Khan Bahadur Judah (G.B.V.C., F.Z.S.)	Poona.
Ichalkaranji, The Hon'ble Meherban Narayan Govind <i>alias</i> Babasaheb Ghorpade	Ichalkaranji.
Idar, H. H. Maharaja Dhiraj Lt.-Col. Shree Sir Dowlat Singhji (K.C.S.I.)	Himatnagar.
Inglis, Chas M. (F.Z.S., M.B.O.U.)	Darjeeling.
Ingoldby, Capt. C. M. (R.A.M.C.)	England.
Ivens, J. H. (F.W.D.)	Europe.
Jamkhandi, Shrimant Shankarrao Parsharamrao <i>alias</i> Appasaheb Patwardhan	Jamkhandi.
Jind, H. H. The Maharaja Sir Ranbir Singh (K.C.S.I. G.C.I.E.)	Jind.

Jodhpur, Lt.-Col. H. H. The Maharaja Sir Umaid Sing Bahadur (K.C.S.I., K.C.V.O.) (<i>Vice-Patron</i>)	...	Jodhpur.
Kaiser Shumsher Jung Bahadur Rana, S. M., Lt.-Gen.	...	Nepal.
Khairpur, H. H. Mir Ali Nawazkhan	...	Khairpur.
Khilchipur, H. H. Durjansal Singh	...	Khilchipur.
Kirk-Greene, Mrs. H.	...	England.
Kolhapur, H. H. The Maharaja Sir Rajaram Chhatra- pati (G.C.I.E.)	...	Kolhapur.
Kotah, H. H. The Maharajadhiraj, Lt.-Col. Sir Umed Singhji Saheb Bahadur (G.C.I.E., C.B.E.)	...	Kotah.
Kurundwad Senior State, Bhalchandra Rao Chintaman <i>alias</i> Anna Saheb Patwardhan	...	Kurundwad.
Lee, D. H.	...	Europe.
Long, G. R. (I.F.S.)	...	England.
Lorimer, Major D. L. R. (C.I.E., I.A.)	...	England.
Lowndes, Capt. D. G.	...	Europe.
Ludlow, F. (M.A., M.B.O.U., I.E.S.)	...	England.
Mackenzie, J. M. D. (I.F.S.)	...	Rangoon.
Manavadar, Khan Shree Fatehdinkhan	...	Manavadar.
Maudlik, Narayen V.	...	Bombay.
Marshall, Arch McL.	...	England.
Marshall, J. McL.	...	Scotland.
Martin, Col. Gerald	...	England.
Maxwell, M.	...	Europe.
McConaghy, Lt.-Col. C. B. (I.M.S.)	...	Nepal.
Millard, A. W. P.	...	Bombay.
Milliard, W. S. (F.Z.S.)	...	England.
Mills, Major J. D.	...	England.
Miraj, Shrimat Gungadhar Rao Ganesh <i>alias</i> Baba- saheb Patwardhan	...	Miraj.
Mitchell, H. H. G.	...	England.
Montagnon, D. J.	...	Bindukuri.
Monteath, G. (I.C.S.)	...	Belgaum.
Morris, Lt.-Col. D. O.	...	England.
Morvi, H. H. The Maharaja Lakhdhiri Waghji	...	Morvi.
Mosse, Lt.-Col. A. H. E. (I.A.)	...	Bombay.
Murland, Major H. F.	...	Mandalay.
Mysore, H. H. The Maharaja Krishnaraj Woodayar Bahadur (G.C.S.I.)	...	Mysore.
Namjoshi, V. K.	...	Cambay.
Naraenji Dwarkadas	...	Bombay.
Narayanlal Bansilal, Raja	...	Bombay.
Narrotum Morarji Goculdas	...	Bombay.
Nawanagar, H. H. The Jamsaheb (G.C.S.I., C.B.E., K.C.S.I.)	...	Jamnagar.
Nepal, Supradipta Manyaber General Mohun Shum- shere Jung Bahadur Rana (K.C.I.E.)	...	Nepal.
Noyce, F. (I.C.S.)	...	Bombay.
Nurse, Lt.-Col. C. G. (F.R.S.)	...	England.
Oberholser, Harry C.	...	Washington D. C.
Ogilvie, G. H. (I.F.S.)	...	Myitkyina.
Olivier, Lt.-Col. H. D. (R.E., F.Z.S.)	...	England.

LIST OF MEMBERS

xxv

Page, A. J. (I.C.S.)	Rangoon.
Palitana, H. H. The Thakar Saheb Bahadur Singhji	Palitana.
Parlett, L. M. (I.C.S.)	England.
Patiala, H. H. The Maharaja Sir Bhupendar Singh	Patiala.
(G.C.S.I., G.C.I.E., G.C.B.E.)	Patiala.
Patiala, Capt. Rao Raja Birindra Singhji	Patiala.
Patuck, P. S. (I.C.S.)	England.
Peacock, E. H. (B.F.S.)	Mawlaik.
Pestonji Jivanji (N.C.S.)	Hyderabad, Dn.
Petit, Dhunjibhoy Bomanji	Bombay.
Petit, Jehangir Bomanji	Bombay.
Phipson, H. M. (F.Z.S.)	England.
Poncins, Viscount Edmund de	France.
Pratap Singh Lal Ram (B.A., L.L.B.)	Dehra Dun.
Radhanpur, H. H. Shri Jalaluddin Khanji	Radhanur.
Rae, Major M. E. (I.A.)	Peshawar.
Rajgarh, H. H. Raja Rawat Sir Birendra Singh	Rajgarh.
(K.C.I.E.)	Rajgarh.
Ravenshaw, Col. C. W.	England.
Reid, Mrs. D. W.	Bombay.
Rewa, H. H. The Maharaja Gulabsingh Bahadur	Rewa.
(Vice-Patron)	Rewa.
Ribeiro, J. (L.C.E.)	Santa Cruz.
Ross, Major Tyrell	England.
Roumania, H. R. H. The Crown Prince Carol	Roumania.
Sanders, Major C. W.	Jullundur.
Sangli, Shrimant Chintamanrao Appasaheb Patwar-	Sangli.
dhan	Sangli.
Savile, L. H. (C.B.)	England.
Schmid, W.	Bombay.
Seton-Karr, Capt. H. W.	England.
Smith, A. Kirke	Bombay.
Smith, H. C.	Rangoon.
Spence, Dr. Arthur D. (M.B.Ch. B. (Edin.))	Rangoon.
Spence, Sir Reginald (Kt.)	Bombay.
Spooner, T. J. (C.E.)	Europe.
Standen, Sir B. (K.C.I.E., C.I.E., I.C.S.)	Europe.
Stanton, W. C.	England.
Stephens, T. R.	Bombay.
Stewart, C. G. (I.P.S., M.A.)	England.
Suket State H. H. Raja Lakshman Sen	Suket.
Swaine, J. K.	Matang.
Swithinbank, Mrs. B. W.	Rangoon.
Tata, Sir Dorabji J. (Kt.)	Bombay.
Tehri Garhwal, H. H. Raja Narendra Shah Saheb	Tehri.
Bahadur	Tehri.
Thomson-Glover, Capt. J. W.	Hangu.
Tilly, T. H.	Europe.
Vaughan, Wm. (F.E.S.)	Europe.
Venning, Col. F. E. W. (C.B.E., D.S.O.)	Meerut.
Venour, Lt.-Col. W. E. (I.A.)	Europe.
Vernay, A. S.	England.

Vesugar, Lt. Jamshed...	Europe.
Waite, H. W.	Phillaur, Punjab
Wales, H. R. H. Edward Prince of (K.G., M.C., etc.)	England.
(Patron)	Bombay.
Walker, Roland	Secunderabad.
Wauchope, Major R. S. (I.A.)	England.
Whistler, Hugh (M.B.O.U., F.Z.S., F.L.S.)	England.
Williamson, W. J. F. (C.M.G., F.Z.S., M.B.O.U.)	England.
Wilson, Sir Arnold (C.S.I., C.M.G., D.S.O., K.C.I.E.)	Kenya Colony.
Wyndham, P. (C.B.E., I.C.S.)	England.
Yerbury, Col. J. W. (R.A.)	England.

MEMBERS

Adam, J. B. Mercer (I.F.S.)	Tavoy.
Addyman, J. (M.L.C.)	Bombay.
Aga Khan, H. H. Aga Sir Sultan Mahomed Shah	Bombay
(G.C.S.I., G.C.I.E.) (<i>Life Member</i>)	Aurangabad.
Ahmadi, Daud I.	Nilgiris.
Aird, J.	Lahore.
Aitchison College, The Principal	Mandalay.
Aitken, Major A. B., (O.B.E., M.C., M.E.I.E.)	Bombay.
Aitken, C. E. (<i>Life Member</i>)	Coimbatore.
Aiyar, T. V. Ramkrishna (B.A., F.Z.S.)	Bombay.
Aldous, A. J.	Bombay.
Alfrey, J. I.	Bombay.
Ali, Salim A.	Bombay.
Alipur, Agri. and Horticultural Society of India, The	Alipur.
Secretary	Ali Rajpur.
Ali Rajpur, Maharaj Kumar Capt. Fattehsing	England.
Alington, Capt. N. S.	Allahabad.
Allahabad University, The Head of the Department of	Shillong.
Botany	Kirkee, Poona.
Allen, Major C. H. P. (I.M.S.)	Maymyo.
Allen, B. C. A.	Alwar.
Alleyne, Lt. H. Massy	Ahmedabad.
Alwar, Lt.-Col. H. H. Sawai Maharaja Sir Jaisingh	New York, U.S.A.
Bahadur (K.C.S.I., G.C.I.E.) (<i>Life Member</i>)	Thayetmyo.
Ambalal Sarabhai	England.
American Museum of Natural History, The Librarian	England.
Anderson, R. K.	Cinnamara.
Anderson-Morshead, Capt. R. Y.	Rangoon.
Andrew, G. P. (I.C.S.)	Bombay.
Andrews, E. A.	Madras.
Angelo, Capt. N. L. (I.A.)	Siliguri.
Annesley, F. C.	Bombay.
Anstead, R. D. (M.A.)	New Zealand.
Antram, Chas B. (F.E.S.)	Bombay.
Aquino, V.	England.
Armstrong, Dr. J. S. (R.A.M.C.) (<i>Life Member</i>)	England.
Ash, H. D.	England.
Ashfield, R. C. (I.C.S.)	England.

Atkinson, D. J. (I.F.S.)	Dehra Dun.
Atkinson, G.	Dehra Dun.
Atkinson, G. R.	Bezwada.
Australian Museum, The Secretary	Sydney.
Awati, Prof. P. R. (I.E.S.)	Bombay.
Bacon, A. L.	Magok.
Bahawalpur, H.H. The Nawab Capt. Sir Sadiq Moham- mad Khan Abassi (<i>Life Member</i>)	Bahawalpur.
Bailey, Lt.-Col. F. M. (C.I.E.)	Gangtok.
Baini Parshad, Dr. (M.Sc.)	England.
Baird, Col.-Comdt. H. B. D. (C.M.G., C.I.E., D.S.O.)	England.
Baker, E. C. Stuart (F.Z.S.) (<i>Life Member</i>)	England.
Balasinor, Babi Shri Jamiatkhanji Manverkhanji	Balasinor.
Bale Sons and Danielson, Ltd., Messrs. John	England.
Balkrishna Vinayek Wassoodew (<i>Life Member</i>)	Bombay.
Ball, G. F. (I.F.S.)	England.
Ball, H. P.	Bombay.
Bally, P. O. R.	Bombay.
Banatvala, Col. Sir H. E. (I.M.S., C.S.I.)	Bandra.
Bang-Haas, Herr Otto	Germany.
Bansda, Maharaul Shri Indra Sinhji Pratap Sinhji Raja Saheb of (<i>Life Member</i>)	Bansda.
Banwell, G. E.	Thaton.
Baria, H.H. The Maharaja Ranjit Singhjee	Baria.
Barker, A. N. (I.F.S.)	Rangoon.
Barlee, K. W. (I.C.S.)	Bombay.
Barnard, J. T. O. (C.I.E.)	Rangoon.
Baroda Museum, The Curator	Baroda.
Baroda, H.H. The Maharaja Sir Sayaji Rao Gaekwar, (G.C.S.I., K.C.S.I.) (<i>Life Member</i>)	Baroda.
Barraud, P. J. (F.Z.S., F.E.S.)	Kasauli.
Barrett, Capt. M. L.	England.
Barrington, A. H. M.	Rangoon.
Barron, W. J.	Rangoon.
Barton, C. S.	Scotland.
Barton, E. L. (<i>Life Member</i>)	England.
Barwani, Capt. H.H. Rana Ranjit Singhji Bahadur (K.C.S.I.) (<i>Life Member</i>)	Barwani.
Bassett, A. E.	Rampur Hat.
Bates, Capt. R. S. P.	England.
Battye, Lt. B. W. (R.A.)	Jubbulpore.
Baxter, N. B.	Nasik.
Bayley d'Castro, Dr. A.	Port Blair.
Bazett, H. M.	England.
Beadnell, C. B.	Vellore.
Beadon, W. R. C. (F.G.S.)	Rangoon.
Beckett, James	Cuddapah.
Beeson, C. F. C. (M.A.)	France.
Beg, Nawab Nazir Jung Bahadur Mirza Nazir	Hyderabad, Dn.
Begbie, Lt.-Col. A. S.	England.
Bell, R. D. (C.I.E., I.C.S.)	Bombay.
Bell, T. R. (C.I.E.)	Kerwar.

Bell, V. G.	Sungei Patani, F.M.S.
Benson, Capt. C. E.	England.
Benson, Major R. L. (D.S.O.)	England.
Bernard, J. L.	Delhi
Berriff, A. H.	Simla.
Betham, Brig.-Genl. R. M. (C.I.E.)	England.
Bewsher Lt.-Col., F. C. (D.S.O., O.B.E., M.C., B.A., F.Z.S., F.R.G.S.)	Suraband.
Beynon, F. C.	Kasauli.
Bharatpur, H.H. Maharaja Lt.-Col. Sawai Kishen Singh, K.C.S.I.	Bharatpur.
Bharucha, F. E. (L.M.E., M.I.M.E., M.I.E.)	Bombay.
Bhatia, Capt. S. L. (I.M.S.)	England.
Bignell, Major G. N.	Landi Kotal.
Bignell, Major R. G. (R.G.A.)	Gibraltar.
Bikaner, Col. H.H. The Maharaja Sir Gunga Singhji Bahadur (G.C.S.I., G.C.I.E., G.C.V.O.) (<i>Life Mem-</i> <i>ber</i>)	Bikaner.
Bikaner, Capt. H.H. Maharaj Kumar Shri Sadul Singhji Bahadur (C.V.O.) (<i>Life Member</i>)	Bikaner.
Bikaner, Maharaj Shri Bijey Singhji Bahadur (<i>Life</i> <i>Member</i>)	Bikaner.
Billimoria, N. M.	Karachi.
Binny, Major A. C. M.	Rawalpindi.
Birch, C. V.	New Zealand.
Bird, B. H. (I.C.S.)	England.
Bird, Capt. William (R.A.M.C.)	Simla Hills.
Bisgaard, L.	Bangkok.
Bissett, Frank	Peermade.
Bissett, J. E.	Meppadi.
Blackie, Capt. F. B.	Calcutta.
Blackwood, Robert C.	Shillong.
Blair, Capt. D. P. (R.A.M.C.)	Liverpool.
Blandy, Major R. (M.C.)	Bombay.
Blanford, H. R. (O.B.E., I.F.S.)	Rangoon.
Blatter, Rev. E. (S.J.)	Panchgani.
Blunt, H. R.	Srinagar.
Boag, G. T. (I.C.S.)	England.
Boalth, V. H.	Lahore.
Boas, H. A.	Kollengode.
Bobriskoy, Count (F.R.G.S.)	England.
Bodding, Rev. P. O.	Mohulpahari.
Bodenham, Capt. K. L. (<i>Life Member</i>)	England.
Bolitho, Capt. E. H. W. (R.F.A.)	England.
Boller, A.	Bombay.
Bor, N. L. (I.F.S.)	Kohima.
Botham, A. W. (C.I.E., I.C.S.)	Shillong.
Bott, J. R. E.	Sunthaik.
Boyd, Major J. E. M. (R.A.M.C.)	Bombay.
Bradley, J. W.	Insein.
Bradshaw, J. P. (<i>Life Member</i>)	Bombay.

LIST OF MEMBERS

313

Braham, N. C.	Lakon Lampang.
Brasney, Lt.-Col. L. P. (I.M.S.)	Jhelum.
Brayne, F. L. (I.C.S.)	Gurgaon.
Brent, H. A. W.	Bombay.
Bridge, M. F. (I.F.S.)	England.
Bridgeman, Lt.-Col. The Hon'ble H. G. O. (R.F.A.)
(Life Member)	Birmingham.
Bridges, Major Hugh C.	Calcutta.
Briggs, Revd. Frank S.	Simla.
Bristow, C. H. (I.C.S.)	Ahmedabad.
Britton, Major J. A. H.	England.
Brodie, N. S. (I.C.S.)	England.
Brodrick, Capt. W. L. C.	Bombay.
Brook, Carlton P. (Life Member)	England.
Brooke, Capt. A. S.	Bombay.
Brooke-Purdon, Major W. (D.S.O., M.C., O.B.E., R.A.M.C.)	Bangalore.
Broughton, Lt.-Col. T. D. (R.F., C.R.F.)	Naini Tal.
Brown, A. F. R. (I.F.S.)	Pakokku.
Brown, George	Ratnapura.
Brown, J. S.	Bombay.
Brown, L. N. (I.C.S.)	Sukkur.
Browne, Lt.-Col. C. H.	Madras.
Browne, E. G.	Gorakhpur
Bryden, Lt.-Col. R. A. (R.A.M.C., D.S.O.)	England.
Burder, John H.	Calcutta.
Burgess, F. G. (I.F.S.)	Maymyo.
Burnett, Prof. K.	Hyderabad, Dn.
Burrows, G. T.	England.
Burton, Brig.-General R. G. (Life Member)	England.
Burton, R. N.	Gauhati.
Burton, Lt.-Col. R. W. (Life Member)	England.
Busher, Rev. R. C.	Naini Tal.
Bute, The Most Hon'ble The Marquis of (Life Member).	Scotland.
Butler, S. G.	Silchar.
Butler, His Excellency Sir Harcourt (G.C.I.E., K.C.S.I., I.C.S.) (Life Member)	Burma.
Buxton, Dr. P. A.	England.
Caccia, Major A. M. (I.F.S.) (Life Member)	England.
Calcutta Zoological Gardens, The Honorary Secretary.	Calcutta.
Caldecott, C. D'a	England.
Calvocoressi, Stephen	Bombay.
Cambay, H. H. Nawab Hussain Yavarkhan Jafar Alikhan Sahib Bahadur (Life Member)	Cambay.
Cambridge, University Museum of Zoology, The Superintendent	Cambridge.
Cameron, Thos. H. (F.Z.S.)	Devicoolam.
Cameron, I. L.	Eheliyagoda
Campbell, A. S.	Karachi.
Campbell, J. A. G.	Bokakhat.
Campbell, R. G.	England.

LIST OF MEMBERS

Campbell, S.	Saktipur.
Canning, Fred. (I.F.S.)	Barielly.
Capito, C. Erik (O.B.E., M. INST. C.E.)	Bushire.
Cardew, Capt. C. G. (R.E.)	Bombay.
Cardwell, R.	Hongkong.
Carter, Major H. St. M. (D.S.O., M.D.)	(Life Member).			England
Cartland, Murray A.	Ghazipur.
Cassamali Jairajbhoy Peerbhoy	(Life Member)			Bombay.
Cassell, R. H.	Moniarah.
Casserly, Lt.-Col. Gordon	France.
Castens, H. E. (I.F.S.)	Rangoon.
Cater, A. N. L. (C.I.E., I.C.S.)	Jaipur.
Cave, Capt. F. O.	Srinagar.
Cave, W. A.	Australia.
Chaldecott, H. D.	Kollengode.
Chamba State, H. H. Raja Ram Singh	Chamba.
Chambers, Capt. G. H.	Jubbulpore.
Chamier, Major R. O.	Shillong.
Champion, F. W. (I.F.S.)	Lansdowne.
Champion, H. G. (I.F.S.)	England.
Chapple, E.	Bangkok.
Chartres, C. B. (M.L.A.)	Calcutta.
Chasen, F. N.	Singapore.
Cheesman, Major R. E. (I.A.)	England.
Chevenix-Trench, C. G. (I.C.S.)	Udaipur.
Chislett, E. V. D.	Bombay.
Chitham, C. C.	Nagpur.
Christopher, Sydney A. (BAR-AT-LAW)	(Life Member).			Rangoon.
Chrystall, A.	Salona.
Clarke, A. M.	Bombay
Clarke, L. O. (I.C.S.)	(Life Member)			England.
Clarke, M. H.	Jakhlbandha.
Clifford, Lt.-Col. R.	England.
Clifford, W. Walters	Jalgaon.
Clive, Capt. J. M'c C. (M.C.)	Nowshera.
Clode, C. W.	Suntikoppa.
Clough, R. J.	Calcutta.

CLUBS—

Ahmednagar Club, The Honorary Secretary	...	Ahmednagar.
Bombay Club, The Honorary Secretary	...	Bombay.
Bombay Gymkhana Club, Ltd., The Secretary	...	Bombay.
Byculla Club, The Honorary Secretary	...	Bombay.
Club of Central India, The Secretary	...	Mhow.
Club of Western India, The Honorary Secretary	...	Poona.
Madras Club, The Secretary	...	Madras.
Mawlaik Gymkhana Club, The Honorary Secretary	...	Mawlaik.
Mercara Book Club, The Honorary Secretary	...	Mercara.
Muktesar Shikar Club, The Honorary Secretary	...	Retani.
Naini Tal Club, Ltd., The Secretary	...	Naini Tal.
Nerbudda Club, The Secretary	...	Jubbulpore.

LIST OF MEMBERS

xxxi

Pegu Club, The Secretary	Pegu.
Port Blair Settlement Reading Club, The Principal	Port Blair
Punjab Club, The Secretary	Lahore.
Quetta Club, Ltd., The Secretary	Quetta.
Royal Bombay Yacht Club, The Secretary	Bombay.
Sardar Club, The Honorary Secretary	Jodhpur.
Secunderabad Club Library, The Honorary Secretary	Secunderabad.
Sholapur Club, The Honorary Secretary	Sholapur.
Sind Club, The Honorary Secretary	Karachi.
Srinagar Club, The Secretary	Srinagar.
United Service Club, The Secretary	Bangalore.
United Service Club, Ltd.,	Simla.
Wellington Gymkhana Club, The Honorary Secretary	Wellington.
Clutterbuck, J. E. (R.E.) (<i>Life Member</i>)	Bhurkunda.
Clutterbuck, Sir Peter H. (<i>Life Member</i>)	England.
Cochrane, R. A.	Katha.
Coldstream, J. (I.C.S.)	Lahore.
Cole, Lt.-Col. B. L.	England.
COLLEGES AND SCIENTIFIC INSTITUTIONS—				
Agricultural College, Burma, The Principal	Mandalay.
American College, The Treasurer	Madura.
Government Collegiate High School, The Head
Master	Bangalore.
Bombay Veterinary College, The Principal	Bombay.
Burma Forest School, The Director	Pyiumana.
Carmichael Medical College, The Principal	Calcutta.
Central Research Institute, The Director	Kasauli.
Agricultural College and Central Research Institute,
The Librarian	Coimbatore.
Kings Institute, The Director	Guindy.
Dow Hill Training College, The Principal	Kurseong.
Ewing Christian College, The Professor	Allahabad.
Forest Research Institute and College, The Forest
Botanist	Dehra Dun.
Forest Research Institute and College, The Principal	Dehra Dun.
Gujrat College, The Principal	Ahmedabad.
Institute of Plant Industry, The Director	Indore.
Punjab Veterinary College, The Principal	Lahore.
Madras Forest College, The Principal	Coimbatore.
Presidency College, The Principal	Madras.
Mayo College, The Principal	Ajmer.
Pasteur Institute, The Director	Rangoon.
Agricultural College, Punjab, The Principal	Lyallpur.
Rajkumar College, The Principal	Rajkot.
Ravenshaw College, The Principal	Cuttack.
St. Mary's High School, The Rector	Bombay.
Royal Institute of Science, The Principal	Bombay.
St. Joseph's College, The Principal	Teppakulam.
Victoria College, Palghat, The Principal	Palghat.
Victoria Technical Institute, The Curator	Nagpur.
Wilson College, The Principal	Bombay.
Central College, The Principal	Bangalore.

Collings, T. H. O.	England.
Colombo Museum, The Librarian	Colombo.
Colthurst, Dr. Ida	Calcutta.
Colvin, Major E. J. D. (I.A.)	Rewa.
Commissioner, The Hon'ble The Chief, North-West Frontier Province	Peshawar.
Condon, E. C. H. (M.A.I., A.M.I.C.E., M.A.)	Bombay.
Congreve, C. R. T.	Valparai.
Conley, Andrew	Liguanla.
Connor, Lt.-Col. Sir Frank (D.S.O., I.M.S.)	Calcutta.
Cooch Behar, Maharaj Kumar Victor N. Narayan	Cooch Behar
(Life Member)	Cooch Behar
Coode, John (I.F.S.)	Ootacamund
Coode, J. M. (Life Member)	England.
Cooke, E. Bernard	Poona.
Cooper, G. E. R.	Maymyo.
Cooper, H. L.	Naini Tal.
Cooper, H. R. (B.Sc., F.C.S.)	Cinnamara.
Corbett, R. H.	Papun.
Corbett, Major T. W. (M.C.)	England.
Cosens, Major F. R.	Fatehgarh.
Coulton, G. F.	Bombay.
Covernton, S. H. (I.C.S.)	Karachi.
Cowie, Rev. A. G. G. (Life Member)	Rawalpindi
Cox, C. E. C. (I.F.S., F.Z.S.)	England.
Cox, Lt.-Genl. Sir Percy (G.C.I.E., K.C.S.I., K.C.M.G., K.C.I.E., F.Z.S., F.R.G.S.) (Life Member)	England.
Craw, H. H. (I.C.S.) (Life Member)	Rangoon.
Crawford, Malcolm M.	England.
Crawford, William G. (I.F.S.)	Rangoon.
Crawford, W. M. (F.E.S., I.C.S.) (Life Member)	England.
Crerar, Hon'ble Mr. J. (C.S.I., C.I.E., I.C.S.)	Delhi.
Crofton, R. M. (I.C.S.)	Chikalda.
Crondace, Capt. C. G.	England.
Cruickshank, F. C. B. (I.C.S.)	Sitapur.
Cuffe, Lady C. T. Wheeler	Ireland.
Culberton, J. M. S. (A.M., I.C.E.) (Life Member)	Karachi.
Cule, T. G.	Lumding
Cumberlege, G. F. J. (D.S.O., M.C.)	Bombay.
Cumming, Capt. O. S. (R.A.)	Jhansi.
Cunningham, Col. A. H. (R.E.)	England.
Cunningham, G. (C.I.E., O.B.E., I.C.S.)	Peshawar.
Currie, Elliott S.	England.
Currie, A. J.	Bombay.
Currie, M. M. L. (I.C.S.)	England.
Cursetji, Khan Bahadur C. M. (Life Member)	Bombay.
Cutch, H. H. The Maharao Sir Shri Kengurji Sawa Bahadur (G.C.I.E., G.C.S.I.) (Life Member and Vice-Patron)	Bhuj-Cutch.
Cutch, Kumar Shri Godji (Life Member)	Cutch.
D'Almeida, J. F. R. (B.A., B.Sc. (HON.)) (Life Member)	Bandra.

Dale, Capt. E. C. B. (R.A.)	England.
Daly, M. O.	Jakhlabandha.
Danson, J. W. W.	Rangoon.
Darby, A. W. (O.B.E.)	Katni.
Dare, I. C. (M.I.S.E.)	Bombay.
Darjeeling, The Curator, Nat. Hist. Mus.	Darjeeling.
Dart, G. Wesche	Punjab.
Das, A. N. (M.R.A.S. (LOND.) F.Z.S., (LOND.))	Calcutta.
Datia, H.H. The Maharaja Lokendra Sir Govind Singh Bahadur (K.C.S.I.)	Datia.
Davenport, Alex L.	England.
Daver, Framroze A.	Bombay.
Davey, G. H.	Kottayam.
Davidson, Lt.-Col. J. (D.S.O., I.M.S.) (<i>Life Member</i>)	Edinburgh.
Davies, R. W. H. (I.C.S.)	Larkana.
Davies, Capt. V. K. N. (I.A.)	Bombay.
Davis, G. (I.C.S.)	Poona.
Davis, P. W. (I.F.S.)	England.
Davis, Dr. W. St. T.	Dwarbund.
Dawes, Lt.-Col. C. D. (I.M.S.)	London.
Dawkins, C. G. E. (I.F.S.)	Rangoon.
Dawson, Capt. W. G. (F.E.S.)	England.
Dean, Edwin	Peshawar.
De Carteret, St. G.	England.
Delacour, Mons. Jean	France.
Delaney, M. F.	Bombay.
Demetriadi, C. C.	Karachi.
deRhe-Philipe, G. W. V. (F.E.S.)	London.
Dhar, Her Highness Laxmibai Saheb Puar (<i>Life Member and Vice-Patroness</i>)	Dhar.
Dharampur, H.H. The Maharana Shree Vijayadevji Rana, Maharaja Saheb of (<i>Life Member</i>)	Dharampur.
Dhrangadhra, H. H. Maharana Shri Sir Ghanshyam Sinhji (K.C.S.I.) (<i>Life Member</i>)	Dhrangadhra.
Dhunjibhoy Bomanji, Sir, <i>Kt.</i> (<i>Life Member</i>)	Bombay.
Dickins, K. F. R. (I.F.S.)	Chanda.
Dickson, Major H. R. P. (C.I.E.)	Bikaner.
Director of the Deutsches Entomologisches Institut der Kaiser-Wilhelm-Gesellschaft	Germany.
Dix, J. W.	Mohammerah.
Dixon, A. F. W. (I.C.S.)	Koraput.
Dobson, F.	Panposh.
Dods, W. K. (C.I.E.)	England.
Dogra, H. R. (B.Sc.)	Madanapalle.
Donald, C. H.	Dharamsala.
Donaldson, A. E.	Rangoon.
Donovan, Lt.-Col. C. (I.M.S.)	England.
d'Orle', Mademoiselle Francoise M.	Bombay.
Dow, H. (I.C.S.)	Hyderabad.
Drake Brockman, D. L. (I.C.S.)	Jodhpur.
Drake-Brockman, Lt.-Col. H. E. (I.M.S., F.Z.S.) (<i>Life Member</i>)	England.

Drummond, Capt. J. G. P.	Delhi.
Dubash, J. K.	Bombay.
Duke, A. H.	Bangkok.
Duke, J. A.	Bombay.
Dunbar-Brander, A. A. (O.B.E., I.F.S.)	Scotland.
Duncan, John E. (P.W.D.)	Rangoon.
Dunlop, Dr. W.	Baghdad.
Dunn, C. W. (C.I.E., I.C.S.)	Maymyo.
Dunsterville, Col. K. S. (R.A.)	London.
Duxbury, Lt.-Col. C. D. (<i>Life Member</i>)	Poona.
Dyer, J. C.	England.
Eates, K. R.	Karachi.
Ebden, J. W.	Delhi.
Edge, Mrs. R. C.	England.
Edie, A. G. (I.F.S.)	Poona.
Elgee, Col. J. W. L. (<i>Life Member</i>)	England.
Elliot-Lockhart, Lt. Wm.	Peshawar
Elliott, A. (C.I.E.)	London.
Elliott, J.	England.
Ellis, C. P. (I.C.S.)	Bassein, Burma
Ellis, R. H. (I.C.S.)	Madras.
Ellis, S. F.	Poona.
Ellison, Bernard C. (C.M.Z.S., F.R.G.S.) (<i>Life Member.</i>)	Bhopal.
English, E. E. (<i>Life Member</i>)	Cairo.
Erb, E. (<i>Life Member</i>)	Bombay.
Evans, F. V. (<i>Vice Patron and Life Member</i>)	England.
Evans, Col. G. H. (C.I.E., F.L.S.)	London.
Evans, T. M. (J.P.)	Bombay.
Evans, Col. W. H. (R.E., C.I.E.)	Peshawar.
Evershed, John	England.
Ezra, Alwyn	Bombay.
Ezra, David	Calcutta.
Fairbank, Rev. H. (<i>Life Member</i>)	Ahmednagar.
Faith, N. W.	Bombay.
Farley, T.	Balipara.
Farr, E. T. C.	Koni.
Farrell, R. C. (I.F.S.)	Hyderabad, Dn.
Farquharson, G. N.	Calcutta.
Farquharson, G. H.	Calcutta.
Faunthorpe, Lt.-Col. J. C.	Lucknow.
Fawcus, L. R. (I.C.S.)	Europe.
Fenton, Major L. S. (I.A.)	London.
Ferguson, Ed. T. D.	Tonk Raj.
Fernandes, B. A.	Bandra.
Ferrar, M. E.	Metali.
Ferrar, Lt.-Col. M. L. (I.A., O.B.E.)	Port Blair.
Field, Capt. F. D. S. (I.M.S.)	Bombay.
Field Frank (M.B.O.U.)	London.
Field, G. G. (I.P.S.)	Moradabad.
Fields-Clarke, V. H. T. (I.F.S.)	London.
Finlay, J. D.	Clarkabad.
Fitz Gerald Moore, A. N.	Bombay.

Fleming, Col. A. M. (I.M.S.)	England.
Fleming, E. W.	London.
Fletcher, Thos. Bainbrigge (F.E.S.)	Pusa.
Florence, James	Kievcoote.
Flynn, A. A. (C.M.Z.S.)	Karachi.
Fooks, H. A.	Calcutta.
Forrington, A.	Bombay.
Forsyth, Dr. C. E.	Europe.
Forsyth, Dr. Wm.	Scotland.
Foster, Major R. T.	England.
Foster, Major Rodney (I.A.)	Mussoorie.
Foulkes, Lt.-Col. T. H. (I.M.S.) (<i>Life Member</i>)	Europe.
Foulkes, R.	Madura.
Fountaine, Miss Margaret (F.E.S.)	London.
Fowler, C. C.	Bombay.
Fraser, Capt. C. J. S.	Putao.
Fraser, Duncan	London.
Fraser, Lt.-Col. F. C. (I.M.S., M.D., F.L.E.S.)	Waltair.
Fraser, Sir Stuart (I.C.S., K.C.S.I., C.I.E.)	Bombay.
Fraser, J. S.	Madras.
Fraser, Major S. G. G.	Bombay.
Frederickson, H.	Calcutta.
Frei, Max (<i>Life Member</i>)	Switzerland.
French, H. A. L.	Karachi.
Frenchman, D. P. (B.Sc.)	Gandamanayakanur.
Freud, G. V. R.	Balchonnur.
Frere, Major A. G. (I.A., F.Z.S.) (<i>Life Member</i>)	Cannanore.
Frohlick, H.	Bombay.
Fry, John T.	Bombay.
Gairdner, K. G.	Lakon-Lampang.
Gamlen, R. L.	Hyderabad, Dn.
Gammie, Prof. G. A. (<i>Life Member</i>)	London.
Garbett, C. C. (I.C.S.)	Bombay.
Garnett, Lt. A. M. (R.E.)	London.
Gatherer, Major R. G. T.	London.
Gauripur, Raja Prabhatchandra Barna (<i>Life Member</i>)	Gauripur.
Gaye, W. C.	England.
Gentle, J. A. R. (I.F.S.)	England.
Geoghegan, Mrs. O. A.	Lahore.
Gervers, Col. F. R. S. (R.E., C.I.E.)	Murree.
Gharpurey, Lt.-Col. K. G. (I.M.S.) (<i>Life Member</i>)	Sholapur.
Ghosh, Prof. Ekendranath (M.Sc., M.D., F.Z.S., F.R.M.S.)	Calcutta.
Gibbs, A. F.	Umaria.
Gibson, E. C. (I.C.S.)	London.
Gibson, R. E. (C.I.E., I.C.S.)	Hyderabad-Sind
Gilbert, C. E. L. (I.F.S.)	Karachi.
Gilbert, T.	Europe.
Gill, E. H. N.	Ghazipur.
Gill, H.	Kalain.
Gillett, B. G. (O.B.E.)	Bombay.
Gillum, S. J.	Bombay.

Gimlette, Col. G. H. D. (C.I.E., I.M.S.)	England.
Gimson, C. (I.C.S.)	Assam.
Girsperger, F.	Bombay.
Gladstone, Capt. H. S. (F.Z.S., F.R.S.E.)	Scotland.
Gladstone-Solomon Capt. W. E. (I.E.S.)	Bombay.
Glascock, L. C. (M.V.O., C.I.E.)	Lyallpur.
Glass, E. L.	Nagpur.
Glazebrook, E. K.	Rangoon.
Gleadow, F. (F.R.M.S.)	England.
Glendenning, P. G.	Moulmein.
Glennie, Capt. E. A. (D.S.O., R.E.)	Europe.
Glover, H. M.	Lahore.
Godding, Capt. H. C. (R.A.M.C.)	London.
Godfrey, E. J. (B.Sc.)	Bangkok.
Goldie, Dr. E. A. (M.C., I.M.S.)	Lahoal, Assam.
Gondal, Kumar Shree Nutversinhji	Jetalsar.
Gonsalves, A. F. (<i>Life Member</i>)	Bandra.
Goodall, Major M.	Sadiya.
Gordon, F. W. (M.C., M.A. (OXON) I.F.S.)	Rangoon.
Gordon, Major J. W.	Jodhpur.
Gordon, R. G. (I.C.S.)	England.
Gore, F. W.	England.
Goschen, The Viscountess	Madras.
Gothorp, P....	Assam.

GOVERNMENT DEPARTMENTS—

The Agricultural Engineer, Burma	Mandalay.
Conservator of Forests, N.C. Division, C.P.	Jubbulpore.
Deputy Director of Agriculture, Burma	Mandalay.
Deputy Director of Agriculture, Myingyan Circle	Meiktila.
Deputy Director of Agriculture, East Central Circle	Pyinmana.
Director of Agriculture, Bombay	Poona.
Director, Bureau of Science	Manila.
Director of Agriculture, B. & O.	Sabour.
Entomologist, The Burma	Mandalay.
Gould, B. J. (C.I.E., I.C.S.)	Peshawar.
Gow, Cedric J.	Madras.
Graham, A. A.	Mussoorie.
Graham, L. (I.C.S.)	London.
Grant, F. A.	Rangoon.
Grant, Lt. R. A.	Bombay.
Gravely, Dr. F. H.	Madras.
Greaves, J. B.	Bombay.
Green, E. Ernest (F.E.S.)	England.
Gregory-Smith, Capt. H. G. (<i>Life Member</i>)	London.
Gregson, T. S.	Bombay.
Griffin, A. M.	Cocanada.
Griffiths, D. T. (B.A., I.F.S.)	Shwebo.
Griffiths, H. M. (P.W.D.)	Lahore.
Grose, F. S.	Taunggyi.
Grossenbacher, G.	Bombay.
Hagenbuch, Fred.	Bombay.

Haigh, P. B. (I.C.S.)	Bombay.
Hamber, Capt. L. G. W.	Poona.
Hamid, A. Ali	Baroda.
Hamid Khan, M. (M.Sc., F.R.M.S.)	Lahore.
Hamilton, K. L. B. (I.C.S.)	Raipur.
Hamilton, A. P. F. (D.S.O., I.F.S.)	Kulu.
Hampson, A.	Calcutta.
Hancock, Capt. C. P.	Rajkot.
Hanhart, S.	Bombay.
Hare, Lt.-Col. G. A. (R.A.)	Kirkee.
Harper, Lt.-Col. A. Forrest	Bombay.
Harris, D. G. (C.I.E.)	Ghazipur.
Harris, W. (M.R.C.V.S.)	Gauhati.
Harrison, Mrs. H. Z.	Bombay.
Harrison, R. T. (A.M.I.C.E.)	Bombay.
Hartnoll, E. S. (I.F.S.)	Moulmein.
Harty, Major A. H. (I.M.S.)	Poona.
Harvey-Kelly, Major C. H. G.	Peshawar.
Harwood, Capt. G. E.	Bombay.
Hasted, Capt. J. S.	Bombay.
Hastings, J. R. G.	Bassein.
Haswell, Capt. F. W.	Bombay.
Hâte, Prof. Vinayakrao N. (B.Sc.)	Bombay.
Haughton, Lt.-Col. H. L.	Bombay.
Hauser, E.	Bombay.
Hawes, C. G.	Bombay.
Hawley, Lt.-Col. W. G. B. J.	London.
Hay, Dr. Alfred	England.
Hay, Capt. R. (I.M.S.)	Seistan.
Headlam, Capt. E. James (R.I.M., C.S.I., C.M.G., D.S.O.)	Bombay.
Heard, Mrs. Freda	Lahore.
Hearsey, Capt. L. D. W. (M.C., V.D.)	Kheri.
Hector, G. P. (M.A., B.Sc.)	Dacca.
Hemmons, Capt. A. J.	Bassein.
Henderson, Major D.	London.
Henderson, J. L.	Quilon.
Herbert, Capt. V. A. (I.A.R.)	Naini Tal.
Heron, A. M. (D.Sc.)	Calcutta.
Heron, F. A. T.	Europe.
Hickie, Major F. C.	England.
Hickie, W. A.	Europe.
Hickin, S. B.	Sukkur.
Hickman, Col. R. St. J. (C.I.E.)	London.
Hide, P. (<i>Life Member</i>)	Indore.
Higginbottom, Dr. Sam	Allahabad.
Higgins, J. C. (I.C.S.)	Imphal.
Hiley, A. C. (I.F.S.)	Dharwar.
Hill, K. A. L. (I.C.S.)	Europe.
Hill, Major R. D. O. (<i>Life Member</i>)	London.
Hingston, Major R. W. G. (I.M.S.)	London.
Hislop, Major J. H. (M.C., I.M.S.)	Europe.
Hodgson, H. P.	London.

Hogg, Capt. R. G.	England.
Holkar, H. H. Tukuji Rao (<i>Life Member</i>)	Indore.
Holland, L. B. (I.F.S.)	Rawalpindi.
Holmes, H. R.	England.
Homfray, Jeston (I.F.S.)	Calcutta.
Hopwood, S. F. (I.F.S.) (<i>Life Member</i>)	London.
Horner, Capt. B. Stuart	Bushire.
Hotson, Hon'ble Mr. J. E. B. (C.S.I., I.C.S.) <i>Life Member</i>	Bombay.
Hotz, E.	Agra.
Howard-White, Lt.-Comdt. C. T. (R.N.)	London, W. 14.
Howell-Jones, Lt.-Col. T. H. (C.I.E., D.S.O., R.A.O.C.)	London.
Howell, Lt.-Col. E. Berkley (C.I.E., I.C.S.)	Dera Ismailkhan.
Hoyos, Count E. (<i>Life Member</i>)	Austria.
Hudson, Col. C. (C.I.E., D.S.O., I.M.S.)	Bombay.
Hudson, Sir Leslie (<i>Kt.</i>)	Bombay.
Huggins, J. R.	Waltair.
Hughes, Major J. E.	Bombay.
Humm, Lt. P. S.	Kasauli.
Humphrys, Lt.-Col. Sir Francis (C.I.E.)	Peshawar.
Hundley, G.	Moulmein.
Hunt, Lt.-Col. S. (I.M.S.)	London.
Hunt, Vice-Admiral Sir Thomas (R.N., K.C.B., C.S.I.)	London.
Husbands, Major H. W. S. (M.C., A.M.I.C.E., S.M.R.E.) (<i>Life Member</i>)	London.
Hutchinson, Lt.-Col. L. T. Rose (I.M.S.)	Bombay.
Hutchinson, Lt.-Col. W. Gordon (I.A., O.B.E.)	London.
Hutchison, L. P.	Azamgarh.
Hutton, C. H.	England.
Hutton, C. I.	Insein.
Hyam, Khan Bahadur Judah (G.B.V.C., F.Z.S.) (<i>Life Member</i>)	Poona.
Hyde, H. A. (M.C., P.W.D.)	Indore.
Ibbotson, Capt. A. W. (M.C., M.B.E., I.C.S.)	Puri.
Ichalkaranji, The Hon'ble Meherban Narayan Govind <i>alias</i> , Babasaheb Ghorpade (<i>Life Member</i>)	Ichalkaranji.
Idar, H. H. Maharaja Dhiraj Lt.-Col. Shree Sir Dowlat Singhji, K.C.S.I. (<i>Life Member</i>)	Himatnagar.
Imperial Library, The Librarian	Calcutta.
Inglis, Chas M. (F.Z.S., M.B.O.U.) (<i>Life Member</i>)	Darjeeling.
Ingoldby, Capt. C. M. (R.A.M.C.) (<i>Life Member</i>)	London.
Ingram, J. A.	N. Wales.
Irwin, H. E. Lord (P.C., G.M.S.I., G.M.I.E.) (<i>Patron</i>)	India.
Irwin, Capt. H. R. (I.A.S.C.)	Europe.
Isaacs, Miss Mozelle (M.A., M.Sc.)	Dombivli.
Ivens, J. H. (P.W.D.) (<i>Life Member</i>)	Europe.
Jabouille, P.	Annam.
Jackson, Capt. E. H. P. (R.H.A.)	Meerut.
Jackson, Dr. T. S.	Europe.
Jacob, J. R.	Bombay.
Jacob, W. R. Le Grand (I.F.S.)	Shillong.
James, Major F. H.	Bombay.

Jamkhandi, Shrimant Shankarrao Parashuramrao, <i>alias</i> Appasaheb Patwardhan (K.C.I.E.) (<i>Life Member</i>)	Jamkhandi.
Jamsetji M. Doctor (F.Z.S., C.M.Z.S.)	Bombay.
Janson, V. T.	Calcutta.
Jaora, H. H. Col. Sir Iftikhor Alikhan (K.C.I.E.)	Jaora.
Jardine, W. E. (C.I.E., I.C.S.)	London.
Jarman, Capt. G. S.	Bombay.
Jenkin, R. Trevor (I.F.S.)	Chikalda.
Jenkins, J. B.	Ahmedabad.
Jermyn, Lt. R. O.	Jhelum.
Jind, H. H. The Maharaja Sir Ranbir Singh (K.C.S.I., G.C.I.E.) (<i>Life Member</i>)	Jind.
Jodhpur, Lt.-Col. H. H. The Maharaja Sir Umaidsing Bahadur (K.C.S.I., K.C.V.O.) (<i>Life Member and Vice Patron</i>)	Jodhpur.
Johnson, Lt. A.	Europe.
Johnson, Kay	Nazira.
Johory, Prof. I. W. (M.A., B.D.)	Europe.
Jones, A. E.	Simla.
Jones, A. J.	Moulmein.
Jones, Capt. W. H. C.	Europe.
Jones, Major W. E.	Europe.
Jourdain, Rev. F. C. R.	England.
Jukes, J. E. C. (C.I.E., I.C.S.)	Simla.
Julius, Sydney	Colombo.
Junagadh, His Highness the Nawab of Kaiser Shumsher Jung Bahadur Rana, S. M. Lt.-Genl. (<i>Life Member</i>)	Nepal.
Karachi, Victoria Museum, The Curator	Karachi.
Keays, Lt.-Col. R. W. C. (I.A.)	London.
Keene, Hugh	Koraput.
Kemp, W. N. R.	Damodarpor.
Kennion, Lt.-Col. R. L. (F.Z.S., C.I.E.)	England.
Kermode, C. W. D. (I.F.S.)	Rangoon.
Kerr, Dr. A. F. G.	Bangkok.
Kerr, H. E. Sir John (K.C.S.I., K.C.I.E.)	Assam.
Kershaw, Miss C. G.	Ceylon.
Khairpur, H. H. Mir Ali Nawazkhan (<i>Life Member</i>)	Khairpur.
Khareghat, M. P. (I.C.S.)	Bombay.
Kharwa, Rao Saheb Gopal Sinha Rathor	Kharwa.
Khan, Sahebzada Sardar Mahomed	Karachi.
Khattan, Dharamsey Mulraj	Bombay.
Khilchipur, H. H. Durjansal Singh (<i>Life Member</i>)	Khilchipur.
Kiddle, J. A. C.	Papun-Salween.
Kiernander, Major O. G.	Iraq.
Kindersley, A. F. (I.C.S.)	Karwar.
King, Capt. E. O. (I.A.R.O.)	London.
King, Major J. St. Aubyn	Jhelum.
Kinloch, Mrs. A. M.	Kotagiri.
Kirby, Col. A. D. (R.A.)	London.
Kirby, R. R.	Kurseong.

LIST OF MEMBERS

Kirk-Greene, Mrs. H. (<i>Life Member</i>)	England.
Kirwan, Noel G. B.	Mangalore.
Knight, H. F. (I.C.S.)	Bombay.
Knight, L. G.	Kalthuritty.
Knight, Robert	Kuala Lumpur.
Koechlin, M. C.	Munnar.
Kolhapur, H. H. The Maharaja Sir Rajaram Chhatrapati (G.C.I.E.) (<i>Life Member</i>)	Kolhapur.
Kotah, H. H. The Maharajadhiraj Lt.-Col. Sir Umed Singhji Saheb Bahadur (G.C.I.E., C.B.E.) (<i>Life Member</i>)	Kotah.
Kunhardt, Lt.-Col. J. C. G. (I.M.S.)	London.
Kuppuswamy, V. S.	Bellary.
Kuroda, Dr. Nagamichi	Tokyo.
Kurundwad Senior State, Meherban Bhalchandrarao Chintaman <i>alias</i> Anna Saheb Patwardhan	Kurundwad.
Kydd, D.	Bombay.
Lahore, Central Museum, The Curator	Lahore.
Laidlay, J. C.	Scotland.
Laird-MacGregor, E. G. L. (I.C.S.)	Poona.
Lakshminarayanan, C.	Madras.
Lambert, G. B. (I.C.S.)	Naini Tal.
Lampard, L. A.	Quilon.
Lane, H. S.	Guntakal.
Latham, H. D.	Vellore.
Latif, Sarhan C.	Calcutta.
La Touche, J. F. D.	Rangoon.
Laud, D. S.	Bombay.
Laurie, M. V. (I.F.S.)	Madura.
Lawson, D. W. (R.N.)	Mandalay.
Lawther, B. C. A.	Peshawar.
Leach, A. H. L.	Rangoon.
Lee, D. H. (<i>Life Member</i>)	Europe.
Lee, Major Robert (I.M.S.)	Bombay.
Leitch, Dr. J. Neil (M.B.B.S. (Lond.), M.R.C.I., L.R.C.P.)	Moran.
LeMarchand, A. E. M.	Europe.
LeMarchand, W. M.	Calcutta.
Leonard, Lt.-Col. W. H. (I.M.S.)	Bombay.
Leonard, P. M. R.	Kutkai.
Leslie, M.	Bombay.
Lewis, E. S. (P.C.S.)	Delhi.
Lewis, J. D.	Osaka, Japan.
Ley, Hon'ble Mr. A. H. (C.S.I., C.I.E., I.C.S.)	Delhi.
Ley, W. E. (I.C.S.)	London.

LIBRARIES—

Agricultural Library, The Librarian, U.P.	...	Nawabgunj.
Allahabad Public Library, The Secretary	...	Allahabad.
Arthur Library, The Honorary Secretary	...	Castle Rock.
Benares, Hindu University Library, The Librarian	...	Benares.

LIBRARIES—*continued.*

Burma, Bernard Free Library, The Honorary Librarian	Rangoon.
Hyderabad, The Librarian, State Library	Hyderabad Dn.
Lahore, University of Punjab, The Librarian	Lahore.
Librarian, Students Library, St. Xavier's College	Bombay.
Lucknow, Public Library, The Honorary Librarian	Lucknow.
Madras, Connemara Public Library, The Principal Librarian	Madras.
Public Library, Museums and National Gallery of Victoria, The Chief Librarian	Melbourne.
Station Library, The Honorary Secretary	Simla.
United Service Library, The Honorary Secretary	Poona.
Lightfoot, S. St. C.	Taunggyi.
Lightfoot, Capt. G. S. (I.P.)	Assam.
Limbdi, H. H. Maharana Shri Sir Daulatsinhjee (K.C.I.E.)	Limbdi.
Lindley-Hinde, G.	Calcutta.
Lindsay, J. H. (I.C.S.)	Calcutta.
Lindsay-Smith, Col. J. (I.A.)	Murree.
Lister, R. S.	Ghooni, Bengal.
Little, E.	Kirkee, Poona.
Lloyd, Col. C. R. (A.S.C., D.S.O.)	Lahore.
Lloyd, J. B.	London.
Loch, Miss N. F.	Nakuria, E. Africa.
Locket, A.	Numaligarh.
Lodge, G. E.	England.
Logan Home, Major W. M.	Ajmer.
Lomas, H. A. (I.C.S.)	London.
Long, G. R. (I.F.S.) (<i>Life Member</i>)	England.
Longbottom, Capt. J. L.	England.
Lorimer, Major D. L. R. (C.I.E., I.A.) (<i>Life Member</i>)	London.
Lory, F. B. P. (M.A.)	Poona.
Lowndes, Capt. D. G. (<i>Life Member</i>)	England.
Lowndes, R. C.	Bombay.
Lowsley, C. O.	Ahmednagar.
Lowther, E. H. N.	England.
Lucas, Major J. de B. T. (R.F.A.)	Europe.
Lucknow, Provincial Museum, The Curator	Lucknow.
Ludlow, F. (M.A., M.B.O.U., I.E.S.) (<i>Life Member</i>)	England.
Luke, J. H.	Narayanganj.
Luke, K. J.	Barrackpore.
Lunawada, Maharaj Kumar Rangitsingh	Lunawada.
Lunham, Lt.-Col. J. L. (I.M.S.)	Bombay.
Lyall, Lt.-Col. R. A.	S. Africa.
Lynch Blossie, H. R. (I.C.S.)	Lahore.
Lytton, H. E. The Right Hon'ble Victor Alexander George Earl (P.C., G.C.S.I., G.C.I.E.)	Bengal.
MacBey, W. D.	London.
MacColl, H. H. (I.F.S.)	Rangoon.
Macdonald, A. St. J.	Dooriah.

MacDonald, A. B. (R.A.)	Glasgow.
MacDonald, R.	Misa.
Macdonell, J. F.	Bombay.
MacDougall, A.	Bombay.
MacGregor, Capt. R. F. D. (I.M.S.)	Bombay.
MacKenna, Sir James (C.I.E., I.C.S.)	London.
Mackenzie, J. M. D. (I.F.S.) (<i>Life Member</i>)	Rangoon.
Mackenzie, Major L. H. L. (I.M.S.)	Gilgit.
Mackenzie, J. M.	Calcutta
Mackenzie, T. J.	Indore.
Mackenzie, Wm.	Madras.
Mackereth, J. (I.F.S.)	Rangoon.
Mackie, A. W. W. (I.C.S.)	Poona.
Mackinlay, Lt.-Col. Chas	Scotland.
Mackwood, F. E.	Colombo.
Mackwood, F. M.	London.
Mac Lachlan, R. B.	Europe.
Mac Lachlan, Capt. D.	Bombay.
Macleod, A. (I.C.S.)	Peshawar.
Macleod, Sir Norman (<i>Kt.</i>)	England.
Macleod, R. D. (I.C.S.)	Etawah.
MacMichael, N. (I.C.S.)	Madras.
Macnaghten, Sir Henry (<i>Kt.</i>)	London.
Macpherson, J. C.	Amritsar.
Macqueen, H. C.	Dehra Dun.
Madan, F. R.	South Mangalore.
Mahomedbhoy Currimbhoy Ebrahim, Sir, Bart	Bombay.
Main, T. F. (B.sc.)	Poona.
Malcolm, C. A. (I.F.S.)	Nagpur.
Maltby, Major C. M. (I.A., M.C.)	India.
Manavadar, Khan Shree Fatehdinkhan (<i>Life Member</i>)	Manavadar.
Mandlik, Narayan V. (<i>Life Member</i>)	Bombay.
Mann, Dr. H. H. (D.sc.)	Poona.
Marjoribanks, N. E. (C.I.E., C.S.I., I.C.S.)	Madras.
Marris, H. E. Sir William (K.C.S.I., K.C.I.E., I.C.S.)	United Province.
Marryatt, N.	Ajmer.
Marshall, Arch. McL. (<i>Life Member</i>)	England.
Marshall, A. W.	Scotland.
Marshall, Col. F. J. (C.B., C.M.G., D.S.O.)	England.
Marshall, Mrs. H. A.	Travancore.
Marshall, J. McL. (<i>Life Member</i>)	Scotland.
Marshall, Brig.-Genl. T. E. (R.A.)	N. Wales.
Martin, Major C. de. C. (I.M.S.)	Rangoon.
Martin, Col. Gerald (<i>Life Member</i>)	London.
Martin, S. J.	Almora.
Mason, Miss E. D.	Madras.
Matthews, W. H.	Pashok.
Mawson, Mrs. G. T.	Malad-Thana.
Maxwell, R. M. (I.C.S., C.I.E.)	Kaira.
Maxwell, M. (<i>Life Member</i>)	Europe.
Mayes, W.	Europe.

Maynard, H. A. V.	Europe.
McCoard, L. A.	Pyinmana.
McConaghy, Lt.-Col. C. B. (I.M.S.) (<i>Life Member</i>)	Nepal.
McGlashan, J. (C.E.)	Calcutta.
McIntosh, Alex. (C.E.)	England.
McIntosh, K. H.	Mussoorie.
McMahon Museum, The Honorary Secretary	Quetta.
Mears, C. E. D.	Indore.
Mehta, Dr. M. V. (F.R.C.P. (Ireland) L.M. (Rotunda) and R.C.P.I., L.M.S.)	Bombay.
Meinertzhagen, Lt.-Col. R.	London.
Menezes, J. Hector	Goa.
Menon, K. G.	Trichoor.
Mercer, David	Scotland.
Merrikin, M. S.	Rangoon.
Merson, Duncan T.	Mango Range.
Metcalf, Capt. M. R. (I.A.)	Europe.
Millard, A. W. P. (<i>Life Member</i>)	Bombay.
Millard, W. S. (F.Z.S.) (<i>Life Member</i>)	England.
Miller, A. C. (O.B.E., M.A.)	Poona.
Miller, John I. (F.R.G.S., F.Z.S.)	Calcutta.
Mills, Major J. D. (<i>Life Member</i>)	England.
Mills, J. P. (I.C.S.)	Calcutta.
Milner, C. E.	Rangoon.
Milroy, A. J. W.	Gauhati.
Miraj, Shrimant Sir Gungadhar Rao Ganesh <i>alias</i> Babasaheb Patwardhan (K.C.I.E.) (<i>Life Member</i>)	Miraj.
Miranda, F. X. (F.L.S.)	Savantwadi.
Mirchandani, Tolo K. (B.E., A.M.I.E., P.F.S.)	Dharwar.
Mitchell, Lt. E. D. Treneer	Peshawar.
Mitchell, F. J.	Lahore.
Mitchell, H. H. G. (<i>Life Member</i>)	London.
Mitchell, Capt. R. St. J. (R.E.)	Europe.
Modi, Dr. Jivanji Jamshetji (B.A., Ph.D., C.I.E., J.P.)	Bombay.
Montagnon, D. J. (<i>Life Member</i>)	Bindukuri.
Monte, Dr. D. A. de (L.M. & S.)	Bandra.
Monteath, G. (I.C.S.) (<i>Life Member</i>)	Belgaum.
Monteath, J. (I.C.S.)	Bombay.
Montgomerie, A. (I.C.S.)	Satara.
Montmorency, Sir Geoffrey (K.C.V.O., C.I.E., C.B.E., I.C.S.)	Simla.
Mooney, H. F. (I.F.S.)	Sambalpur.
Moore, J.	London.
Moore, Capt. A. C. (I.A.)	Mawlaik.
Morgan, F. I.	Javali.
Morgan, R. W. D.	Scotland.
Morgan, Major J. S. H.	Madras.
Morison, Major J. (I.M.S.)	Bombay.
Morris, Chas. F.	Bombay.
Morris, Lt.-Col. D. O. (<i>Life Member</i>)	London.
Morris, R. C.	Attikan.

Morris, Chas. W. G.	Attikan.
Morton, Geo B.	Calcutta.
Morvi, H.H. The Maharaja Lakhdhiri Waghi (<i>Life Member</i>)	Morvi.
Mosse, Lt.-Col. A. H. E. (I.A.) (<i>Life Member</i>)	Bombay.
Mudhol, Shrimant Sir Malojirao Raje Ghorpade (K.C.I.E.)	Mudhol.
Muir, G. B. F. (I.C.S.)	Europe.
Muir, Wm. Angus (M.A., M.C., I.F.S.)	Bombay.
Mullan, Prof. J. P. (M.A., F.Z.S., F.L.S.)	Bombay.
Muller, Dr. H. C. (D.Sc.)	Bombay.
Mulroney, J. T.	Kurseong.
Mundy, N. S.	Silchar.
Munns, F. A. C.	Barhampur.
Munro, F. H.	Europe.
Murland, Major H. F. (<i>Life Member</i>)	Mandalay.
Murphy, P. J.	Paksey.
Murray, Lt.-Col. J. H. (C.I.E., I.M.S.)	Bombay.
Murray, A. E.	Moran.
Musgrave-Hanna, Capt. J. R.	Mandalay.
Mysore, The Director of Agriculture	Bangalore.
Mysore, Government Museum, The Superintendent	Bangalore.
Mysore, H.H. The Maharaja Krishnaraj Woodayar Bahadur (G.C.S.I., G.B.E.) (<i>Life Member</i>)	Mysore.
Nagpur, Central Museum, The Curator	Nagpur.
Namjoshi, V. K. (<i>Life Member</i>)	Cambay.
Napier-Ford, G. S.	Vandiperiyar.
Naraenji Dwarkadas (<i>Life Member</i>)	Bombay.
Narayanlal Bansilal, Raja (<i>Life Member</i>)	Bombay.
Narrotum Morarji Goculdas (<i>Life Member</i>)	Bombay.
Nash, H. J.	Bombay.
Nawanagar, H.H. The Jam Saheb (G.C.S.I., C.B.E., K.C.S.I.) (<i>Life Member</i>)	Jamnagar.
Needham, J. E. (C.B.E.)	England.
Needham, F. M.	Murkong-Sellek.
Nepal, Supradipta Manyaber General Mohun Sumshere Jung Bahadur Rana (K.C.I.E.) (<i>Life Member</i>)	Nepal.
Nevill, Capt. G. A. (I.A.)	Balipara.
Nevill, Capt. T. N. C.	London.
Newcome, Lionel	Somwarpett.
Newland, Major B. E. M. (I.M.S.)	Bombay.
Newman, Lt.-Col. H. E. M.	Waziristan.
Nicholetts, W. A. B.	Rangajan.
Nicholson, Lt.-Col. F. L. (D.S.O., M.C.)	England.
Niederer, George	Bombay.
Norman, Col. H. H. (R.A.M.C.)	England.
Norman, John B.	Champaran.
Norris, J. P.	U. S. A.
Nougerede, L. J. de la	Sadiya.
Noyce, F. (I.C.S.) (<i>Life Member</i>)	Bombay.
Nurse, Lt.-Col. C. G. (F.E.S.) (<i>Life Member</i>)	England.

LIST OF MEMBERS

xlv

Oberholser, Harry C. (<i>Life Member</i>)	Washington, D. C.
O'Brien, Lt.-Col. Edward	Kolhapur.
O'Brien, H. C.	Johannesburg.
O'Callaghan, T. P. M.	Sadiya.
O'Donel, H. V.	Hasimara.
Ogilvie, G. H. (I.F.S.) (<i>Life Member</i>)	Myitkyina.
Oliver, A. W. L.	Hankow, China.
Oliver, Major D. G.	Srinagar.
Oliver, L. C.	Attikan.
Olivier, Col. H. D. (R.E., F.Z.S.) (<i>Life Member</i>)	England.
Ollenbach, O. C. (F.E.S.)	Dehra Dun.
O'Neill, Major H. J. D.	London.
Ormiston, W.	Ceylon.
O'Rorke, Major J. M. W.	London.
Osborne, A. F.	Tuticorin.
Osborne, W. H. (I.F.S.)	Rangoon.
Osmaston, A. E. (I.F.S.)	Naini Tal.
Osmaston, B. B. (C.I.E., I.F.S.)	England.
Osmaston, B. H.	Haldwani.
O'Sullivan, A. G. J.	Godhra.
Oswald Little, Major J.	Darjeeling.
Paddison, R. H. (P.W.D.)	England.
Page, A. J. (I.C.S.) (<i>Life Member</i>)	Rangoon.
Palanpur, H.H. Sir Nawab Saheb Taley Mahomed, Khan Bahadur (K.C.I.E., K.C.V.O.)	Palanpur.
Palitana, H.H. The Thakor Saheb Bahadur Singhji (<i>Life Member</i>)	Palitana.
Palmes, W. T. (I.C.S.)	London.
Panday, Mrs. J. L.	Bombay.
Parish, N. B.	Europe.
Parker, H. (I.C.S.)	Shwebo.
Parlett, L. M. (I.C.S.) (<i>Life Member</i>)	England.
Parry, N. E. (I.C.S.)	Aijal, Assam.
Paterson, E. A.	Calcutta.
Patiala, H.H. The Maharaja Sir Bhupendar Singh (G.C.S.I., G.C.I.E., G.C.B.E.) (<i>Life Member</i>)	Patiala.
Patiala, Capt., Rao Raja Birindra Singhji (<i>Life Member</i>)	Patiala.
Patterson, A.	London.
Patuck, P. S. (I.C.S.) (<i>Life Member</i>)	London.
Pawsey, C. R. (I.C.S.)	Mangaldai.
Peacock, E. H. (B.F.S.) (<i>Life Member</i>)	Mawlaik.
Pears, Hon'ble Mr. S. E. (C.S.I., C.I.E., I.C.S.)	Coorg.
Pearson, Capt. R. Forbes (M.C.)	Bolarum.
Peck, N. F. (I.C.S.)	Calcutta.
Peddie, G.	Secunderabad.
Peebles, W. J. M.	London.
Peppe, Capt. F. H. (R.F.A.)	Nowshera.
Percival, A. P. (I.F.S.)	England.
Percival, Hon'ble Mr. P. E. (C.I.E., I.C.S.)	Karachi.
Perfect, E.	London.

Perry, Major S. (R.A.)	London.
Pershhouse, Major Stanley	London.
Pestonji Jivanji (N.C.S.) (<i>Life Member</i>)	Hyderabad, Dn.
Petit, Dhunjibhoy Bomanji (<i>Life Member</i>)	Bombay.
Petit, Jehangir Bomanji (<i>Life Member</i>)	Bombay.
Petley, C. A.	Toungoo.
Petrie, Walter M.	Karachi.
Phatak, Major V. M.	Gwalior.
Phayre, Major R. B. (M.C.)	Pachmarhi, C. P.
Philip, C. L. (I.C.S.)	Sambalpur.
Phillips, A. A. (I.S.R.)	Europe.
Phillips, G. C.	Bombay.
Phillips, W. W. A.	Matugama.
Phipson, H. M. (F.Z.S.) (<i>Life Member</i>)	London.
Phythian-Adams, Major E. G. (I.A.)	Nilgiris.
Pilditch, D.	Benares.
Pinhey, Capt. L. A. G.	Hyderabad, Dn.
Pipe, T. S.	Dharwar.
Pitman, Major C.R.S.	Africa.
Pogson, Major F. V.	Bombay.
Poncins, Viscount Edmund de (<i>Life Member</i>)	France.
Pope J. A. (I.C.S.)	Indore.
Pratap Singh Lal Ram (B.A., LL.B.) (<i>Life Member</i>)	Dehra Dun.
Prendargast, H. V.	Cawnpore.
Priestley, E. (D.S.P.)	Dharwar.
Pringle, Sir Norman, Bt.	Prome.
Procter, Sir Henry (<i>Kt.</i> , C.B.E.)	London.
Procter, Miss Joan B. (F.Z.S.)	London.
Pudukkottai, State Museum, The Curator	Pudukkottai.
Purkis, F. C. (B.F.S.)	Rangoon.
Purkis, H. Vernon	Bhadarwa.
Quarry, F. G.	Dehra Dun.
Quarry, P. S.	Chinsirah.
Radhanpur, H. H. Shri Jalaluddin Khanji (<i>Life Member</i>)	Radhanpur.
Rae, Major M. E. (I.A.) (<i>Life Member</i>)	Peshawar.
Ragg, H. A.	Munnar.
Raitt, W. Chas.	Kutta.
Rajadhyaksha, G. S. (I.C.S.)	Satara.
Rajgarh, H.H. Raja Rawat Sir Birendra Singh... (K.C.I.E.) (<i>Life Member</i>)	Rajgarh.
Rajkot, H.H. Sir Lakhajiraj (K.C.I.E.)	Rajkot.
Ralston, Lt.-Col. W. H. (D.S.O., I.A.)	Razani.
Rane, K. R. (M.M. and S.P.)	Andheri.
Rane, Yeshwantrao K.	Andheri.
Ravenshaw, Col. C. W. (<i>Life Member</i>)	London.
Rayner, A. B. (I.S.E.)	Balloki, Punjab.
Readymoney, N. J.	Bombay.
Rees, R.	Calcutta.
REGIMENTAL OFFICERS' MESSES—				
The Royal Engineers	Roorkee.

REGIMENTAL OFFICERS' MESSES—*continued.*

1st Bn. The Worcestershire Regiment	Meerut
1st Kings Shropshire	Poona.
6th D.C.O. Lancers	Meerut.
Royal Deccan Horse	Dera Ismail Khan.
Queen Victoria's Own Madras Sappers and Miners	Bangalore.
1st Bn. 2nd Bombay Pioneers	Bombay.
1st Bn./4th Bombay Grenadiers	Bombay.
4/4th Bombay Grenadiers	Razani.
1/6th Rajputana Rifles	Razmak.
3/6th Rajputana Rifles	Landi Kotal.
1st Royal Bn. 9th Jat Regt. L.I.	Fyzabad.
3/10th Baluch Regiment	Lahore.
4/10th Baluch Regiment	Quetta.
1/12th F.F. Regt. (P.W.O. Sikhs)	Jhansi.
4/12th F.F. Regt. (Sikhs)	Ambala.
5/12th Q.V.O. Guides, F.F.	Mardan.
1/13th F.F. Rifles (Cokes)	Razmak.
6/13th Royal F.F. Rifles	Delhi.
3/15th Punjab Regiment	Allahabad.
1/16th Punjab Regiment	Jhelum.
3/17th Dogra Regiment	Alipore.
2/1st K.G.O. Gurkha Rifles	Razmak.
2nd K.E.O. Gurkhas	Dehra Dun.
1/10th Gurkha Rifles	Fort Sandeman.
Reid, E. C.	Europe.
Reid, Mrs. D. W. (<i>Life Member</i>)	Bombay.
Reuben, D. E. (I.C.S.)	Cuttack.
Rewa, H.H. The Maharaja Gulabsingh Bahadur			
(<i>Life Member and Vice Patron</i>)	Rewa.
Reynolds, L. W. (C.I.E., I.C.S.)	Jodhpur.
Ribeiro, J. (L.C.E.) (<i>Life Member</i>)	Santa Cruz.
Richards, B. D. (B.Sc.)	England.
Richardson, F. D. S.	England.
Richmond, G. N.	Bhamo, Burma.
Richmond, R. D. (I.F.S.)	Ootacamund.
Ridland, J. G.	Bombay.
Rishworth, Dr. H. R. (I.M.S.)	Bhusawal.
Ritchie, Sir Adam (<i>Kt.</i>)	Rangoon.
Ritchie, Lt.-Col. W. D. (I.M.S.)	Europe.
Roberts, A. A. Lane (I.C.S.)	Gurdaspur.
Roberts, Dr. C. L. Digby	Mariani, Assam.
Roberts, H. V. H.	Karachi.
Robertson, Lawrence (C.S.I., I.C.S.)	England.
Robinson, A. C.	England.
Robinson, H. C.	Kuala Lumpur.
Robinson, Capt. J. A.	Miranshah.
Robinson, J. B.	Sagrampur.
Robinson, G. C.	Mercara.
Rodger, A. (I.F.S., O.B.E.)	Dehra Dun.
Rogers, C. G. (I.F.S.)	London,

Rogers, Rev. K. St. A. (C.M.S.)	Nairobi.
Rogerson, A. M.	Rangoon.
Rondano, Rev. A. (S.J.)	Mangalore.
Roosevelt, Kermit	New York.
Roosevelt, Col. Theodore	New York.
Ross, Major Tyrell (<i>Life Member</i>)	England.
Ross, Major E. J.	Dehra Dun.
Roumania, H. R. H. The Crown Prince Carol (<i>Life Member</i>)	Roumania.
Row, Lt.-Col. G. R. (I.A.)	Shillong.
Row, Dr. R. (M.D.)	Bombay.
Rowland, J. W. (I.P.)	Satara.
Rowlands, R. D.	Bombay.
Rowlandson, B. C.	Bombay.
Rowson, C.	Europe.
Rowson, W. S.	Vandiperiyar.
Roy, N.	Dacca.
Roy, R. Trevor	Pegu.
Rozalla, A. A.	Simla.
Ruddle, W. H.	Madras.
Russell, Lt.-Col. G. H. (C.I.E., D.S.O.)	Ambala.
Russell, Major J. Bertrand	Saklaspur.
Ruttledge, Capt. R. F. (M.C.)	Peshawar.
Ryan, J. G.	Cawnpore.
Salkeld, Lt.-Col. R. E.	England.
Sanders, D. F.	Calcutta.
Sanders, Major C. W. (<i>Life Member</i>)	Jullunder.
Sanderson, P. M. D. (F.Z.S.)	Calcutta.
Sandilands, Dr. J. E. (M.C., M.A., M.D.)	Bombay.
Sangli, Srimant Chintamanrao Appasaheb Patwardhan (<i>Life Member</i>)	Sangli.
Sant, Maharana Shri Jarwar Sinhji Raja Saheb of	Sant-Rampur.
Sarangarh, Raja Bahadur Jawahir Singh	Sarangarh.
Sarawak Museum, The Curator	Kuching.
Satya Churn Law, Dr. (M.A., B.L., Ph.D., F.Z.S., M.B.O.U.)	Calcutta.
Saunders, H. F.	Madura.
Saunders, L.	Koraput.
Saunders, Lt.-Col. F. W. (R.E.)	London.
Sausman, V. A. N.	Vaghai.
Savantwadi, H. H., The Sardesai	Savantwadi.
Savile, L. H. (C.B.) (<i>Life Member</i>)	London.
Sawyer, H. H.	Bombay.
Scale, Major J. D. (D.S.O., O.B.E.)	England.
Schmid, W. (<i>Life Member</i>)	Bombay.
Schomberg, Lt.-Col. R. C. F. (D.S.O.)	Nowshera.
Scott, A. G.	Jubbulpore.
Scott, C. W. (I.F.S.)	Insein.
Scott, Major F. B. (I.A.)	Bombay.
Scroope, Capt. C. F.	Bombay.
Searight, Capt. E. E. G. L. (M.C., R.E.)	Roorkee.

Secretary for Agriculture, S. S. and F.M.S.	...	Kuala Lumpur.
Seervai, Dr. Rustom F.	...	Bombay.
Seton-Karr, Capt. H. W. (<i>Life Member</i>)	...	London.
Sevastopulo, D. G. (F.E.S.)	...	Lyallpore.
Seymour-Sewell, Major R. B. (I.M.S.)	...	Calcutta.
Shah, Capt. B. (I.M.S.)	...	London.
Shannon, G. C. (I.C.S.)	...	Thana.
Shaw, G. E.	...	Riyang.
Shaw, G. L.	...	Jalpaiguri.
Shaw, John	...	Hsipaw.
Shaw, Col. H. D.	...	Kohat.
Shebbeare, E. O.	...	Europe.
Shephard, T. F. G.	...	Ajmer.
Sheppard, S. T.	...	Bombay.
Sherred, P. R.	...	Ramgarh.
Sherriff, Albert	...	London.
Shevade, Prof. S. V. (B.Sc., Dip-in-Agr. (Camb.) F.Z.S., F.R.H.S.)	...	Baroda.
Shipp, W. E.	...	Ajmer.
Shirley, G. S.	...	Rangoon.
Short, Capt. J. Mc. L.	...	Nowshera.
Shortt, Major H. E. (I.M.S.)	...	Bombay.
Shortt, W. H. O.	...	Mailani.
Shoveller, H. L. (A.C.A.)	...	Bombay.
Simmonds, A.	...	Dimakusi.
Simmons, R. M.	...	Bombay.
Simonds, Major M. H.	...	England.
Simons, H. B.	...	Ferozepore.
Simpson, Col. J. C.	...	Maymyo.
Sinclair, R. L.	...	Bombay.
Singh, Capt. Malik Anup (M.B., I.M.S.)	...	Taiping.
Sitwell, Major N. S. H. (R.A.)	...	Paris.
Sitzler, E. A. (I.F.S.)	...	Mawlaik.
Skinner, Major J. M. (I.M.S.)	...	Bombay.
Skrine, Capt. P. R. H.	...	Europe.
Sladen, J. M. (I.C.S.)	...	Poona.
Sladen, Dr. R. J. L. (F.R.C.S.)	...	Calcutta.
Slater, A. F. (A.M.I.M.E., M.B.E.)	...	Aligarh.
Smith, A. Kirke (<i>Life Member</i>)	...	Bombay.
Smith, Charles J.	...	Jharia.
Smith, H. C. (<i>Life Member</i>)	...	Rangoon.
Smith, J. I. (I.C.S.)	...	Waltair.
Smith, Dr. Malcolm	...	London.
Smyth, C. P.	...	Alipore.
Snelling, Capt. C. G.	...	Bombay.

SOCIETIES—

Royal Asiatic Society, Bombay Branch, Honorary Secretary	...	Bombay.
New York Zoological Society	...	New York, U.S.A.
Souter, C. A. (I.C.S.)	...	Waltair.
Spalding, Lt.-Col. W. B. (R.A.)	...	Cawnpore.

LIST OF MEMBERS

Sparrow, Col. R.	England.
Spence, Dr. Arthur D. (M.B., CH.B. (EDIN.) (<i>Life Member</i>)	Rangoon.
Spence, Sir Reginald, (<i>Kt.</i>) (<i>Life Member</i>)	Bombay.
Spong, Major William A. (R.A.M.C.)	England.
Spooner, T. J. (C.E.) (<i>Life Member</i>)	Europe.
Spurgin, Capt. A. R.	Bombay.
Stable, Capt. R. H. (D.S.O.)	Ali Masjid.
Stables, Major Alex. (R.A.M.C.)	Scotland.
Standen, Sir B. (K.C.I.E., C.I.E., I.C.S.) (<i>Life Member</i>)	Europe.
Stanford, J. K. (M.C., M.A., M.B.O.U., I.C.S.)	Mawlaik.
Stanton, W. C. (<i>Life Member</i>)	London.
Stent, P. J. H. (I.C.S.)	Nagpur.
Stephens, T. R. (<i>Life Member</i>)	Quetta.
Stevens, Herbert (F.Z.S., M.B.O.U.)	England.
Stevenson, Capt. H. A. (I.A.)	Chaman.
Stewart, C. G. (I.P.S., M.A.) (<i>Life Member</i>)	London.
Stewart, Prof. R. R. (Ph.D.)	Rawalpindi.
Stileman, D. F. (I.F.S.)	Nilambur.
Stirling, J. H.	Jodhpur.
Stirling, Col. Wm. (C.M.G., D.S.O., R.A.)	London.
Stockley, Lt.-Col. C. H. (D.S.O., O.B.E., M.C.)	China.
Stokes, H. G. (C.I.E., I.C.S.)	Chetput.
Stokes, I. W.	England.
Stoney, R. F.	Madras.
Storey, J. O.	Alleppey.
Stuart, J. D.	Rangoon.
Studd, E.	Calcutta.
Sudan Government Museum, The Curator	Khartoum.
Sujan, N. B. (B.Sc. (Edin.) F.Z.S.)	Karachi.
Suket, H. H. Raja Lakshman Sen (<i>Life Member</i>)	Suket.
Suter, Dr. M. F. (D.Sc.)	Bombay.
Swaine, J. K. (<i>Life Member</i>)	Matang.
Swan, E. J. A.	Papun.
Swithinbank, Mrs. B. W. (<i>Life Member</i>)	Rangoon.
Sylvester, Capt. E. C. (R.F.A.)	Vandiperiyar.
Symes, A. W. J.	Sabarmati.
Symons, C. T.	Colombo.
Symons, Major-General T. H. (I.M.S., O.B.E., C.S.I.)	Simla.
Tabuteau, Major G. G. (D.S.O., R.A.M.C.)	Rawalpindi.
Taleyarkhan, K. M. (Bar-at-Law)	Bombay.
Tasker, T. J. (I.C.S., O.B.E.)	Mercara.
Tata, Sir Dorabji J. (<i>Kt.</i>) (<i>Life Member</i>)	Bombay.
Tate, Major J. C. (I.A.)	Chhota Udepur.
Taylor, G. J. S.	S. Rhodesia.
Taylor, Dr. W. R. (M.R.C.S., L.R.C.S.)	Insein.
Taylor, Lt.-Col. J. (D.S.O., I.M.S.)	Rangoon.
Tehri Garhwal State, H.H. Raja Narendra Shah Saheb Bahadur (C.S.I.) (<i>Life Member</i>)	Tehri.
Thatcher, E. C.	Sahuspur.

Theobald, Lt. J. A.	Chakrata.
Thom, W. S.	Paletwa.
Thomas, F.	Bombay.
Thompson, C. H. (I.F.S.)	Rangoon.
Thomson, Major D. B.	England.
Thomson—Glover, Capt. J. W. (<i>Life Member</i>)	Hanga.
Thorns—Roberts, J. W. B.	Rangoon.
Thornton, H. A. (I.C.S.)	Maymyo.
Thursfield, R. M. C.	Rangoon.
Tibbs, Rev. Canon P. G.	Europe.
Ticehurst, Dr. C. B. (M.A., M.B.O.U., R.A.M.C.)	England.
Tilden, H. B. (F.C.S.)	London.
Tilly, T. H. (<i>Life Member</i>)	Europe.
Tombazi, N. A. (F.R.G.S.)	France.
Tomlinson, A. G.	England.
Travers, W. L. (C.B.E.)	Jalpaiguri.
Traylen, G. D.	London.
Treasuryvala, N. J.	Bombay.
Treuthardt, M.	Bombay.
Trevenen, Major W. B.	Poona.
Triggs, Bernard	Rutlan.
Trotter, Genl. E. W.	Bangkok.
Tuggerse, M. S.	Honawar.
Tunnard, T. E.	Ramboda, Ceylon.
Tunstall, A. C.	Cinnamara.
Turkhud, Dr. D. A. (M.B.C.M., etc.)	Kodaikanal.
Turnbull, D.	Tuticorin.
Turnbull, Mrs. S. R.	Madras.
Turner, Sir Montague	London.
Tyers, G. A. V.	London.
Tyler, H. H. F. M. (I.C.S.)	Anantapur.
Tyrrell, Lt.-Col. J. R. (I.M.S.)	London.
Tyson, John D. (I.C.S.)	Calcutta.
Tytler, Major-Genl. Sir Henry (K.B.B., C.I.E., C.M.G., D.S.O.)	Maymyo.
Ubsdell, Lt. A. R.	Calcutta.
University of Amoy, The Head of the Department of Zoology	Amoy, China.
Upton, T. E. T.	Europe.
U. S. Department of Agriculture, The Librarian	Washington, D.C.
Van Ingen, E. M.	Mysore.
Vaughan, Wm. (F.E.S.) (<i>Life Member</i>)	Europe.
Venning, Col. F. E. W. (C.B.E., D.S.O.) (<i>Life Member</i>)	Meerut.
Venour, Lt.-Col. W. E. (I.A.) (<i>Life Member</i>)	Europe.
Vernay, A. S. (<i>Life Member</i>)	London.
Vernon, H. A. B. (I.C.S.)	London.
Vesugar, Jamshed (<i>Life Member</i>)	Europe.
Victoria Memorial Park, The Honorary Secretary	Rangoon.
Vijayarajji, Maharaj Kumar, Shri	Bhaj-Cutch.
Villar, A. R. (I.F.S.)	Europe.
Vitty, C. P.	England.
Volkart, L.	Switzerland.

Wait, W. E.	Colombo.
Wales, H. R. H. The Prince of (K.G., M.C., etc.)				
(Patron and Life Member)	England.
Waite, H. W. (Life Member)	Phillaur, Punjab.
Walker, Major E. Clare (R.E.)	Maymyo.
Walker, Roland (Life Member)	Bombay.
Wall, Col. F. (C.M.G., C.M.Z.S., K.H.S., I.M.S.)			...	London.
Walton, S. W.	Bombay.
Wapshare, J. H.	Guynd, Nilgiris.
Ward, Lt.-Col. A. E.	Pandraithan,
				Kashmir
Waterfield, H. G.	Indore.
Watson, H.W.A. (I.F.S.)	Rangoon.
Watson, Lt.-Col. J. W. (I.M.S.)			...	Bombay.
Wauchope, Major R. S. (I.A.) (Life Member)			...	Europe.
Webb, J. E. N.	Adra.
Webb, Major Geo. A. (V.D.)	Mirik.
Webb Ware, George	Kodaikanal.
Weldon, Dr. R. P. (M.D.)	Chittagong.
West, L. C.	Hubli.
Weston, A. T.	Europe.
Whistler, Hugh (M.B.O.U., F.Z.S., F.L.S.) (Life Member)			...	London.
White, L. S. (I.C.S.)	Cawnpore.
Whitehead, Capt. W. A.	Nowshera.
Whitehead, John (I.F.S.)	Europe.
Whitman, Dr. V. E.	Mohammerah.
Whymper, S. L.	London.
Wickham, P. F. (C.E.)	Rangoon.
Wilkes, J. S.	London.
Wilkie, J. S.	Peermade.
Wilkins, P. M.	Somwarpet.
William, Capt. Chas. H. (I.A.S.C.)			...	Quetta.
Williams, Lt.-Col. C. E. (I.M.S.)			...	London.
Williams, Capt. G. B.	Bombay.
Williams, Lt. P. (R.A.)	Razmak.
Williams, J. L. H.	Valparai.
Williamson, A. (I.C.S.)	Rangoon.
Williamson, F. W. (I.C.S.)	Gangtok.
Williamson, P. R. C.	Trichinopoly.
Williamson, W. J. F. (C.M.G., F.Z.S., M.B.O.U.) (Life Member)	London.
Wilson, Sir Arnold (C.S.I., C.M.G., C.I.E., D.S.O., K.C.I.E.) (Life Member)	London.
Wilson, Charles	Dehra-Dun.
Wilson, Commdr. George (R.N.)			...	G.P.O., Bombay.
Wilson, Major C. H. E. (R.F.A.)			...	Kharaghoda.
Wilson, Mrs. D. W.	Bombay.
Wilson, H.E. The Rt. Hon'ble Lt.-Col. Sir Leslie (P.C., G.C.I.E., C.M.G., D.S.O.)			...	Bombay.
Wilson, R. A. (I.C.S.)	Nagpur.
Winch, H. J.	London.

Winckworth, Major H. C. (R.A.M.C.)	Bangalore.
Withers, D. S.	Mariani.
Withers, F. W. (I.F.S.)	London.
Witt, D. O. (I.F.S.)	Pachmarhi.
Wittwer, Dr. R. (D.Sc.)	Bombay.
Wood, Lt. H. B.	Fort Sandeman.
Wood, Noel H.	Bombay.
Wood, W. S.	Tavoy.
Woods, D. F.	Bombay.
Wooldridge, C. J.	Madras.
Wordsworth, Major R. G.	Bombay.
Worgan, Col. R. B. (C.S.I., C.V.O., D.S.O.)	Delhi.
Workman, W. H. (F.Z.S., M.B.O.U.)	Ireland.
Wrangham—Hardy, G.	Darjeeling.
Wright, H. Courtnay	London.
Wright, J. M. (I.C.S.)	Myaungmya.
Wright, Capt. D. Moncriff	Scotland.
Wright, Rev. G. A. A.	England.
Wright Neville, Capt. V. R.	Hongkong.
Wyndham, P. (C.B.E., I.C.S.) (<i>Life Member</i>)	Kericho.
Yeolekar, T. G. (M.A., B.Sc.)	Poona.
Yerbury, Col. J. W. (R.A.) (<i>Life Member</i>)	London.
Young, Major H. G. (D.S.O., R.F.A.)	England.
Young, R. H.	Karachi.
Zollinger, A. E.	Tuticorin.
Zuberbihler, H. W.	Bombay.
Zurmuhle, E.	Bombay.

INDEX OF SPECIES

	PAGE		PAGE
<i>Abraximorpha chamunda</i>	73	<i>Ægithaliscus concinnus iredalei</i> ...	370, 461
————— <i>dauidii</i>	73	————— <i>leucogenys</i>	491
<i>Acacia arabica</i>	262	————— <i>niveogularis</i>	461
<i>Acanthion hodgsoni</i>	401	<i>Ægithina nigrolutea</i>	263
————— <i>leucurus</i>	382	<i>Ægyptius monachus</i>	108, 477
<i>Acanthis cannabina fringillirostris</i> ...	94	<i>Æmona lena haynei</i>	260
————— <i>flavirostris brevirostris</i> ...	472	<i>Ænanthe ænanthe</i>	103
<i>Acanthopneuste o. occipitalis</i> ...	188, 194	————— <i>argentea</i>	103
<i>Accipiter nisus</i>	480	————— <i>rostrata</i>	103
————— <i>nisosimilis</i>	108	————— <i>alboniger</i>	103
————— <i>virgatus</i>	226	————— <i>finschii barnesi</i>	103
————— <i>affinis</i>	480	————— <i>hispanica</i>	103
<i>Achalarus bifasciatus aborica</i> ...	69	————— <i>melanoleuca</i>	103
————— <i>bifasciatus</i>	69	————— <i>moesta</i>	103
————— <i>casyapa</i>	69	————— <i>brooksbanksi</i>	103
————— <i>liliana</i>	69	————— <i>xanthoprymna cummingi</i> ...	103
————— <i>germanus</i>	69	————— <i>xanthoprym-</i>	
————— <i>nepos</i>	69	————— <i>na</i>	103
————— <i>proximus</i>	69	<i>Æromachus catocyanea catocyanea</i> ...	431
————— <i>simplex</i>	69	————— <i>kali</i>	431
<i>Acidalia actiosaria</i>	140	————— <i>discreta discreta</i>	432
————— <i>geminifera</i>	143	————— <i>javanica</i>	432
————— <i>marcidaria</i>	137	————— <i>piceus</i>	432
————— <i>patularia</i>	139	————— <i>inachus chinensis</i>	431
<i>Aclerda ischœni</i>	454	————— <i>inachus</i>	431
————— <i>japonica</i>	454	————— <i>stigmata</i>	431
<i>Acomys flavidus</i>	407	————— <i>jhora jhora</i>	432
<i>Acridotheres ginginianus</i>	267	————— <i>pygmæus musca</i>	432
————— <i>tristis tristis</i> 267, 286, 371, 471		————— <i>pygmæus</i>	432
<i>Acrocephalus agricola</i>	187, 224	<i>Ætheolepis papillosa</i>	319
————— <i>arundinaceus</i>	100	<i>Æthiopsar fuscus fuscus</i>	471
————— <i>babylonicus</i>	100	<i>Agave americana</i>	262
————— <i>concinens concinens</i> 186, 224		<i>Agnibesa sanguiniplaga</i>	143
————— <i>dumetorum</i> ... 264, 470, 496		<i>Agrobates galactotes familiaris</i> ...	102, 496
————— <i>griseldis</i>	100	<i>Alæmon alaudipes</i>	373
————— <i>palustris</i>	101	————— <i>cinerea</i>	97
————— <i>schoenobænus</i>	101	————— <i>pallida</i>	97
————— <i>scirpaceus scirpaceus</i> ...	101	<i>Alauda arvensis cantarella</i>	96
————— <i>stentoreus</i>	496	————— <i>guttata</i>	475
————— <i>zarudnyi</i>	100	————— <i>gulgula guttata</i>	188, 373
<i>Adhotada vasica</i>	262	————— <i>raytal raytal</i>	273
<i>Adelura cæruleocephala</i>	467, 493	<i>Alcedo atthis</i>	476

INDEX OF SPECIES

lv

	PAGE		PAGE
<i>Alcippe phæocephala phæocephala</i> ...	304	<i>Anthus campestris</i> ...	97
<i>Alectoris græca chukar</i> ...	376, 483	----- <i>griseus</i> ...	97, 272
----- <i>kurdestanicus</i> ...	118	----- <i>cervinus</i> ...	97
----- <i>græca weræ</i> ...	118	----- <i>hodgsoni</i> ...	474
<i>Aloe vera</i> ...	262	----- <i>leneophrys jerdoni</i> ...	474
<i>Alseonax latirostris poonensis</i> ...	371, 469	----- <i>pratensis</i> ...	97
----- <i>rubricaudus</i> ...	463	----- <i>richardi rufulus</i> ...	272, 284, 474
<i>Amandava amandava amandava</i> ...	269	----- <i>roseatus</i> ...	372, 474
<i>Amaurornis fuscus</i> ...	489	----- <i>rufulus</i> ...	372
<i>Amblonax cinerea</i> ...	394	----- <i>sordidus</i> ...	272
<i>Ammomanes deserti</i> ...	97	----- <i>spinoletta blakestoni</i> ...	474
----- <i>cheesmani</i> ...	97	----- <i>jerdoni</i> ...	372
----- <i>coxi</i> ...	97	----- <i>trivialis trivialis</i> ...	97, 272
----- <i>fraterculus</i> ...	97	<i>Antoniria bambusæ</i> ...	453
----- <i>phoenicuroides</i> ...	288	----- <i>maritenia</i> ...	453
<i>Ammoperdix griseogularis</i> ...	118	<i>Aonidia bullata</i> ...	456
----- <i>ter-meuleni</i> ...	118	----- <i>columnifera</i> ...	456
<i>Ampittia dalailama dalailama</i> ...	431	----- <i>mesochitinsa</i> ...	456
----- <i>nanus</i> ...	431	----- <i>mimusopis</i> ...	456
----- <i>dioscorides</i> ...	430	<i>Apithecina viridata</i> ...	144
----- <i>maga</i> ...	431	<i>Apodemus speciosus orestes</i> ...	400
----- <i>trimacula maraoides</i> ...	431	----- <i>griseus</i> ...	407
----- <i>trimacula</i> ...	431	<i>Aporia nabellica ferrari</i> ...	252
----- <i>virgata</i> ...	430	<i>Apostictopterus fuliginosus</i> ...	429
<i>Anaïtis affinis</i> ...	313	<i>Appias indra shiva</i> ...	429
<i>Anas acuta</i> ...	109	----- <i>var. aristoxenus</i> ...	529
----- <i>angustirostris</i> ...	110	<i>Aquila chrysætus daphanæ</i> ...	478
----- <i>boscas</i> ...	522	----- <i>clanga</i> ...	479
----- <i>platyrhyncha</i> ...	110, 485	----- <i>heliaca</i> ...	478
----- <i>pœcilorhyncha pœcilorhyncha</i> ...	287, 484	----- <i>heliaca</i> ...	108
----- <i>strepera</i> ...	109	----- <i>nipalensis orientalis</i> ...	108, 478
<i>Anastomus oscitans</i> ...	223	----- <i>rapax</i> ...	108
<i>Anathana pallida</i> ...	402	----- <i>vindhiana</i> ...	375, 479
<i>Anhinga rufus</i> ...	109	<i>Arboricola torqueola millardi</i> ...	483
<i>Anisogomphus caudalis</i> ...	423	<i>Arceuthornis</i> ...	493
----- <i>occipitalis</i> ...	421	----- <i>pilaris</i> ...	494
----- <i>orites</i> ...	422	----- <i>viscivorus bonapartei</i> ...	467
<i>Anisodes absconditaria</i> ...	136	<i>Archæobalbis</i> sp. ...	131
----- <i>hyriaria</i> ...	136	<i>Arctogalidia leucotis</i> ...	46
----- <i>sp. n.</i> ...	136	----- <i>merriam</i> ...	46
<i>Anomalococcus cremastogastri</i> ...	452	<i>Arctomys caudatus</i> ...	189
----- <i>hirsutus</i> ...	452	<i>Arctonyx collaris</i> ...	393
<i>Anourosorex squamipes</i> ...	391	<i>Ardea cinerea</i> ...	488
<i>Anser anser</i> ...	109, 484	----- <i>cinerea</i> ...	110
----- <i>brachyrhynchus</i> ...	522	<i>Ardea goliath</i> ...	110, 523
----- <i>indicus</i> ...	484	----- <i>manillensis</i> ...	488
<i>Anthoscopus pendulinus persimites</i> ...	99	<i>Ardeola grayi</i> ...	378
<i>Anthropoides virgo</i> ...	112	----- <i>ralloides ralloides</i> ...	111
<i>Anticlea cuprearia</i> ...	310	<i>Argemone mexicana</i> ...	262
----- <i>schistacea</i> ...	308	<i>Argya caudata caudata</i> ...	263, 280, 370
<i>Antilope cervicapra</i> ...	382, 508	----- <i>eclipses</i> ...	491

	PAGE		PAGE
<i>Argya caudata huttoni</i> ...	491	<i>Botaurus stellaris stellaris</i> ...	111
— <i>malcolmi</i> ...	263, 280	<i>Brabira artemidora</i> ...	321
<i>Arnetta atkinsoni</i> ...	432	— <i>atkinsoni</i> ...	321
— <i>kala</i> ...	433	<i>Brachypternus aurantius aurantius</i> ...	284
— <i>verones</i> ...	433	— <i>delulus</i> ...	373
— <i>vindhiana</i> ...	432	<i>Brachypteryx major major</i> ...	300
<i>Artamus fuscus</i> ...	225	<i>Branta ruficollis</i> ..	109
<i>Arvicola melanogaster</i> ...	400	<i>Bubo bubo bengalensis</i> ...	297, 477
<i>Asellia tridens murraiana</i> ...	40	— <i>nikolskii</i> ...	106
<i>Asio otus</i> ...	107	— <i>ruthenus</i> ...	106
<i>Aspidiotus ambalangoda</i> ...	456	— <i>turcomanus</i> ...	477
— <i>calophylli</i> ...	456	— <i>coromandus</i> ...	297, 375
— <i>var. symp-</i>		<i>Bubulcus coromandus</i> ...	378
— <i>locos</i> ...	456	<i>Budorcas taxicolor</i> ...	203, 274, 401
— <i>longispinus</i> ...	456	<i>Bungarus cæruleus</i> ...	228
— <i>perniciosus</i> ...	456	<i>Burhinus oedie nemus</i> ...	112
— <i>pinnulifera var. diversicoor</i>	456	— <i>astutus</i> ...	112
— <i>tripinnatus</i> ...	456	— <i>saharæ</i> ...	112
<i>Aspidoproctus cinerea</i> ...	451	<i>Burmagomphus cauvericus</i> ...	413
<i>Asteriolecanium gutta</i> ...	452	— <i>hasimaricus</i> ...	411
— <i>lineare</i> ...	452	— <i>insularis</i> ...	408
— <i>loranthi</i> ...	452	— <i>jacobseni</i> ...	408
— <i>minutum</i> ...	452	— <i>laidlawi</i> ...	412
— <i>pseudomiliaris</i> ...	452	— <i>pyramidalis</i> ...	409
<i>Astictopterus jama henrici</i> ...	429	— <i>siamensis</i> ...	411
— <i>jama</i> ...	429	— <i>sivalikensis</i> ...	414
— <i>olivascens</i> ...	429	— <i>vermiculatus</i> ...	408
<i>Astur badius dussumieri</i> ...	480	— <i>v-flavum</i> ...	414
— <i>gentilis</i> ...	480	— <i>williamsoni</i> ...	410
<i>Athene noctua bactriana</i> ...	107	<i>Butastur teesa</i> ...	375, 479
— <i>lilith</i> ...	107	<i>Butea frondosa</i> ...	262
<i>Atherurus assamensis</i> ...	401	<i>Buteo buteo japonicus</i> ...	480
<i>Atopophysa indistincta</i> ...	322	— <i>ferox</i> ...	196, 480
<i>Aulocera padma grandis</i> ...	253	<i>Cacomantis merulinus passerinus</i> ...	477
— <i>swaha garuna</i> ...	253	<i>Calandrella amtirostris</i> ...	186
— <i>gilgitica</i> ...	253	— <i>brachydactyla</i> ...	191
— <i>kurrama</i> ...	253	— <i>heinei</i> ...	97
— <i>swaha</i> ...	253	— <i>minor</i> ...	96
<i>Badamia exclamationis</i> ...	64	— <i>rufescens</i> ...	96
<i>Bandicota nemorivaga</i> ...	397	— <i>minor</i> ...	96
— <i>sp.</i> ...	403	— <i>persica</i> ...	96
<i>Baracus hampsoni</i> ...	430	— <i>adamsi</i> ...	373
— <i>vittatus septentrionum</i> ...	430	<i>Callabraxas trigoniplaga</i> ..	312
— <i>subditus</i> ...	430	<i>Callacanthus burtoni</i> ...	194, 472
— <i>vittatus</i> ...	430	<i>Calliana pieridoides</i> ..	65
<i>Barbastella darjelingensis</i> ...	389	<i>Calliope pectoralis pectoralis</i> ...	192, 466
<i>Belostoma indicum</i> ...	228	<i>Callosciurus epomophorus davisoni</i> ...	47
<i>Berberis aristata</i> ...	262	— <i>domelicus</i> ...	47
<i>Bibasis sena sena</i> ...	63	— <i>hastilis</i> ...	46
<i>Bibos gaurus</i> ...	220	— <i>sullivanus</i> ...	47
<i>Botaurus stellaris</i> ...	488	— <i>tabaudius</i> ...	47

INDEX OF SPECIES

xviii

	PAGE		PAGE
<i>Callosciurus erythræus aquilo</i> ...	396	<i>Carterocephalus abax</i>	428
----- <i>erythræus</i> ...	395	----- <i>argyrostigma argyro-</i>	
----- <i>erythrogastr.</i>	395	<i>stigma</i> ...	428
----- <i>nagarum</i> ...	195	----- <i>avanti</i>	428
----- <i>wellsi</i> ...	396	----- <i>montana</i>	428
<i>Calluga costalis</i>	320	----- <i>christophi christophi</i>	429
<i>Calostigia albigirata</i>	144	----- <i>micio</i> ...	429
<i>Calothyranis sp.</i>	136	----- <i>dieckmanni dieck-</i>	
----- <i>correspondens</i>	136	<i>manni</i> ...	428
----- <i>oligoscia</i>	136	----- <i>dulcis</i> ...	428
<i>Canis indicus</i>	393, 402	----- <i>flavomaculatus</i> ...	428
----- <i>kola</i>	41, 381, 405	----- <i>niveomaculatus niveomacu-</i>	
----- <i>lupus laniger</i>	403	<i>latus</i> ...	428
----- <i>pallipes</i>	403	----- <i>thibetanus,</i>	
<i>Cannomys badius</i>	401		428
<i>Capila jayadeva</i>	64	----- <i>palæmon</i>	427
----- <i>mackwoodi</i>	64	----- <i>pulchra</i>	428
----- <i>omeia</i>	64	----- <i>silvius houangty</i> ...	427
----- <i>translucida</i>	64	----- <i>shoaka</i>	427
----- <i>zennara</i>	64	----- <i>silvius</i>	427
<i>Capella gallinago</i>	484	<i>Catreus wallichi</i>	482
----- <i>nemoricola</i>	484	<i>Celænorrhinus ambareesa</i>	65
----- <i>solitaria</i>	484	----- <i>asmara asmara</i> ...	68
----- <i>stenura</i>	484	----- <i>cacus</i>	67
<i>Capparis aphylla</i>	262	----- <i>consertus</i>	67
----- <i>horrida</i>	262	----- <i>aspersa</i>	66
<i>Capricornis sumatrensis rubidus</i> ...	401	----- <i>aurivittata aurivittata</i>	69
<i>Caprimulgus ægyptius ægyptius</i> ...	106	----- <i>cameroni</i>	68
----- <i>asiaticus</i>	297	----- <i>vimana</i>	69
----- <i>europæus meridionalis</i> ...	106	----- <i>badia</i>	69
----- <i>unwini</i>	106, 374	----- <i>consanguinea</i>	69
----- <i>zarudnyi</i>	106	----- <i>dhanada affinis</i> ...	65
----- <i>mahrattensis</i> ..	375	----- <i>andamanica</i>	68
----- <i>sp.</i>	476	----- <i>dentatus</i>	68
<i>Caprona agama</i>	78	----- <i>dhanada</i>	68
----- <i>var. elwesi</i>	79	----- <i>flavocincta</i>	66
----- <i>ransonnettii alida</i>	78	----- <i>inaequalis</i>	69
----- <i>potiphera</i>	78	----- <i>ladana</i>	69
----- <i>ransonnettii</i> ..	78	----- <i>leucocera brahmaputra</i>	66
<i>Caranx jarra</i>	228, 229	----- <i>chinensis</i>	66
<i>Carcharodus alceæ swinhoei</i>	82	----- <i>leucocera</i>	66
----- <i>altheæ dravira</i>	82	----- <i>maculicornis</i>	67
----- <i>boeticus</i>	82	----- <i>maculosa</i>	66
----- <i>lavatharæ</i>	82	----- <i>munda</i>	67
<i>Cardiococcus bivalata</i>	455	----- <i>nigriscans balukinus</i> ...	67
<i>Carduelis caniceps caniceps</i>	189, 194, 472	----- <i>nigriscans</i>	68
----- <i>carduelis niediecki</i>	94	----- <i>patula</i>	68
----- <i>spinus</i>	94	----- <i>pero</i>	66
<i>Carpodacus erythrinus roseatus</i>	189, 190,	----- <i>plagifera</i>	66
	191, 270, 472	----- <i>pulumaya</i>	65
----- <i>rhodochoea</i>	472	----- <i>pyrrha</i>	65

	PAGE		PAGE
<i>Celænorhynchus ruficornis</i> area ...	67	<i>Chloris chloris turkestanica</i> ...	95
----- <i>ruficornis</i> ...	67	<i>Chloroclystis chlorophilata</i> ...	319
----- <i>saturatus</i> ...	67	----- <i>consueta</i> ...	319
----- <i>simula</i> ...	66	----- <i>flicata</i> ...	319
----- <i>snelleni</i> sema n. sp. ...	67	----- <i>inæquata</i> ...	320
----- <i>snelleni</i> ...	67	----- <i>nigroviridata</i> ...	145
----- <i>spilothyrus</i> ...	67	----- <i>papillosa</i> ...	319
----- <i>sumitra</i> ...	66	----- <i>rubroviridis</i> ...	320
----- <i>tibetana</i> ...	68	<i>Chrysocolaptes guttacristatus delesserti</i> ...	292
----- <i>tytleri</i> ...	66	<i>Ciconia ciconia ciconia</i> ...	111
<i>Ceratodus</i> ...	503	<i>Cidaria albigrata</i> ...	144
<i>Cercococcus paradoxa</i> ...	452	----- <i>albofasciata</i> ...	309
<i>Cercomela fusca</i> ...	268, 278	----- <i>aliena</i> ...	311
----- <i>fryeri</i> ...	454	----- <i>cerva</i> ...	308
----- <i>iceryoides</i> ...	454	----- <i>cinereata</i> ...	312
<i>Certhia h. himalayana</i> ...	188, 463, 508	----- <i>inextricata</i> ...	312
<i>Cervus f. megaceros</i> ...	172, 179, 222	----- <i>lipartaria</i> ...	310
<i>Ceryle lugubris guttulata</i> ...	476	----- <i>latifasciaria</i> ...	310
----- <i>rudis rudis</i> ...	106	----- <i>maculata</i> ...	304
<i>Cettia celti orientalis</i> ...	100	----- <i>relata</i> ...	311
<i>Chætolopha incurvata</i> ...	313	----- <i>seriata</i> ...	310
<i>Chaimarrornis leucocephala</i> ...	188, 371, 465	----- <i>subapicaria</i> ...	312
<i>Chalyboclydon flexilinea</i> ...	143	----- <i>viridata</i> ...	144
<i>Charadrius alexandrinus alexandri-</i>		<i>Cidrus gerardiana</i> ...	34
nus ...	113, 377	<i>Cimex rotundatus</i> ...	206
----- <i>dubius jerdoni</i> ...	377, 483	<i>Cinclus cinclus cashmeriensis</i> ...	464
----- <i>mongolus albifrons</i> ...	192	----- <i>pallasii tenuirostris</i> ...	188, 464
----- <i>atrifrons</i> ...	377	<i>Cannyris a. asiatica</i> ...	370
<i>Charmion ficulnea</i> ...	65	----- <i>a. brevirostris</i> ...	373
<i>Charronia flavigula</i> ...	393	<i>Circaetus gallicus</i> ...	479
<i>Chartographa fabislaria trigomiphaga</i> ...	312	<i>Circus æruginosus æruginosus</i> ...	108, 480
<i>Chaulelasmus streperus</i> ...	485	----- <i>cyaneus</i> ...	480
<i>Chelidon urbica</i> ...	473	<i>Cisticola juncidis cursitans</i> ...	264, 370, 470
<i>Cheliones hurrianæ</i> ...	41, 406	----- <i>neurotica</i> ...	102
<i>Chettusia gregaria</i> ...	113	<i>Clamator glandarius</i> ...	106
----- <i>leucura</i> ...	113	----- <i>jacobinus</i> ...	477
<i>Chionaspis acuminata var. atricolor</i> ..	455	<i>Coelops frithi</i> ...	388
----- <i>centripetalis</i> ...	455	<i>Coenotephria mononyssa</i> ...	144
----- <i>elongata</i> ...	455	<i>Coladenia agni</i> ...	76
----- <i>gynandropsidis</i> ...	455	----- <i>agnioides</i> ...	76
----- <i>herbæ</i> ...	455	----- <i>dan dan</i> ...	75
----- <i>linearis</i> ...	455	----- <i>dhyana</i> ...	75
----- <i>longissima</i> ...	455	----- <i>fatih</i> ...	75
----- <i>megaloba</i> ...	455	----- <i>indrani atarana</i> n. sp. ...	75
----- <i>pusa</i> ...	455	----- <i>indra</i> n. sp. ...	75
----- <i>tenera</i> ...	455	----- <i>indrani</i> ...	75
<i>Chiropodomys gliroides</i> ...	400	----- <i>tissa</i> ...	75
<i>Chlamydotis undulata macqueeni</i> ...	112	----- <i>uposathra</i> ...	75
<i>Chlidonias leucoparcia leucoparcia</i> ...	114	----- <i>laxmi</i> ...	76
----- <i>leucoptera</i> ...	114	----- <i>vitrea</i> ...	76
<i>Chloris chloris</i> ...	95	<i>Golias cocandica hindnencica</i> ...	253

	PAGE		PAGE
<i>Colias marcopolo</i>	252	<i>Cryptoloba etaina</i>	322
— <i>weskotti chrysoptera</i>	252	— <i>togulata n. sp.</i>	321
<i>Collocalia fuciphaga brevirostris</i>	476	<i>Cryptolopha x. xanthoschista</i>	187, 371
<i>Collix flavofasciata</i>	314	<i>Cryptostegia grandiflora</i>	262
— <i>ghosha</i>	313	<i>Ctenochiton cinnamomi</i>	455
<i>Columba leuconota</i>	481	— <i>fryeri</i>	455
— <i>leuconota</i>	192	— <i>olivaceum</i>	455
— <i>livia</i>	481	<i>Ctenoptilum multiguttata</i>	77
— <i>gaddi</i>	117	— <i>vasava</i>	77
— <i>palumbus palumbus</i>	117	<i>Cuculus canorus</i>	187, 373, 476
— <i>rupestris</i>	481	— <i>subtelephonus</i>	105
— <i>turkestanica</i>	192	— <i>telephonus</i>	189, 194
<i>Comatibis emerita</i>	111	— <i>poliocephalus poliocephalus</i> ..	187, 477
<i>Comibæna cenocraspis</i>	133	— <i>saturatus</i>	187
— <i>swanni</i>	132	<i>Culicicapa ceylonensis ceylonensis</i> . 268, 469,	494
<i>Comostola virago</i>	135	— <i>meridionalis</i>	494
<i>Conurus cyanocephalus</i>	373	<i>Cursorius coromandelicus</i>	376
— <i>torquatus</i>	373	— <i>cursor cursor</i>	112
<i>Copsyche saularis saularis</i> 269, 371, 467, 493		— <i>gallicus</i>	112
<i>Coracias benghalensis benghalensis</i> 106, 273		<i>Cyanops asiatica</i>	373
— <i>garrula semenowi</i>	187, 475	— <i>flavifrons</i>	373
<i>Corvus collaris</i>	93	<i>Cyanosylvia cyanecula abbotti</i>	469
— <i>corax laurencei</i>	93, 263, 490	— <i>suecica</i>	190
— <i>tibetanus</i>	188, 459	— <i>abbotti</i>	371
— <i>cornix</i>	93	— <i>pallidogularis</i> . 268, 493	
— <i>coronoides</i>	263	— <i>suecica</i>	493
— <i>intermedius</i> . 370, 459, 490		<i>Cyclogomphus heterostylus</i>	160
— <i>frugilegus</i>	93	— <i>gynostylus</i>	162
— <i>tschusii</i>	490	— <i>minusculus</i>	163
— <i>monedula</i>	459	— <i>vesiculosus</i>	161
— <i>sœmmerengii</i>	93	— <i>wilkinsi</i>	161
— <i>paradisi</i>	495	— <i>ypsilon</i>	159
— <i>splendens</i>	459, 508	<i>Cynopterus brachyotis angulatus</i>	411
— <i>splendens</i>	263, 291	— <i>sphinx</i>	386
— <i>zugmayeri</i>	490	<i>Cyornis superciliaris</i>	371, 468
<i>Cosmostolia subtiliaria</i>	135	<i>Cyrtostomus asiaticus asiaticus</i> . 273, 475	
<i>Coturnix coturnix coturnix</i>	118, 483	<i>Dacnomys wroughtoni n. sp.</i>	398
<i>Craspedia bispurcata</i>	137	<i>Daimio bhagava andamanica</i>	74
— <i>undulataria pulverosa</i>	139	— <i>bhagava</i>	74
— <i>straminea</i>	139	— <i>celebica</i>	75
<i>Crateropus altirostri</i>	102	— <i>limax diræ</i>	75
— <i>caudatus huttoni</i>	102	— <i>phisara</i>	74
<i>Cremnoconchus syhadrensis</i>	447	— <i>sinica indica</i>	74
<i>Crirolecanium formicarium</i>	454	— <i>narada</i>	74
— <i>radicicola</i>	454	— <i>sinica</i>	74
<i>Crocidura ilensis</i>	405	— <i>tabrica pinwilli</i>	75
— <i>rubricosa</i>	391, 404	— <i>tabrica</i>	75
<i>Crocopus</i>	375	— <i>tethys birmana</i>	74
<i>Crossiura pennicillatum</i>	64	<i>Darpa hanria</i>	77
<i>Cryptoloba aerata</i>	321		
— <i>cinerea</i>	322		

	PAGE		PAGE
<i>Darpa pteria</i>	77	<i>Dryobates auriceps</i>	373, 475
— <i>striata</i>	77	— <i>himalayensis</i>	475
<i>Davidioides martini</i>	419	— <i>hyperythrus</i>	475
<i>Davidius aberrans</i> ^{aberrans}	169	— <i>medius sancti-johannis</i>	105
— <i>senchalensis</i> subsp. n.	170	— <i>syriacus</i>	105
— <i>dauidi dauidi</i>	166	<i>Dysstroma cinereata</i>	312
— <i>assamensis</i>	167	— <i>citrata dentifera</i>	312
— <i>malloryi</i> n. sp.	167	— <i>subapicaria</i>	312
— <i>zallorensis delineatus</i> ..	164, 166	<i>Ecliptopera lativittaria</i>	311
— <i>zallorensis</i>	165	— <i>relata</i>	311
<i>Delichon urbica alexandrovi</i>	105	— <i>silaceata</i>	311
— <i>meridionalis</i>	105	<i>Egretta garzetta</i>	377
— <i>urbica</i>	189, 191	<i>Elanus cæruleus cæruleus</i>	480
<i>Demiegretta sacra asha</i>	116	<i>Electrophaës aliena</i>	311
<i>Dendrocitta formosæ occidentalis</i>	459	<i>Elwesia lesliei</i>	429
— <i>pallida</i>	490	<i>Emballonura monticola</i>	45
— <i>rufa</i>	490	<i>Emberiza calandra calandra</i>	95
— <i>kinneari</i>	491	— <i>cia</i>	95
— <i>sclateri</i>	491	— <i>par</i>	95, 270
— <i>vagabunda</i>	263	— <i>stracheyi</i>	187, 193, 473
<i>Diaspis amygdali</i>	456	— <i>citrinella</i>	95
— <i>antiquorum</i>	456	— <i>erythrogenys</i>	95
— <i>bambusæ</i>	456	— <i>stracheyi</i>	372
— <i>barbatus</i>	456	— <i>fucata arcuata</i>	473
— <i>boisduvali</i>	456	— <i>hortulana</i>	96
— <i>calyptroides</i>	456	— <i>leucocephala</i>	473
— <i>grandilobis</i>	456	— <i>melanocephala</i>	95
— <i>heneratgoda</i>	456	— <i>schoenicius pallidior</i>	96
— <i>mihiriya</i>	456	— <i>stewarti</i>	473
— <i>orientalis</i>	456	— <i>maculatus maculatus</i>	465
— <i>phoenicis</i>	456	<i>Enicurus m. guttatus</i>	371
<i>Dicæum erythrorhynchum</i>	373	<i>Eoglaucmys fimbriatus</i>	403, 405
— <i>ignipectum</i>	475	<i>Eois rubridentata</i>	142
<i>Dicurus leucophæus longicaudatus</i>	470	<i>Eothenomys melanogaster</i>	400
— <i>stevensi</i>	187	<i>Epirrhoe (?) latifasciata</i>	309
— <i>longicaudatus longicaudatus</i> ..	371	<i>Erebia mani kamriana</i>	256
— <i>macrocerus</i>	469, 496	— <i>lorimeri</i>	257
— <i>macrocerus</i>	264, 284, 371	— <i>shandura</i>	257
<i>Dindica molleri</i>	132	— <i>yasina</i>	257
<i>Dipodilus indus</i>	42	<i>Eremophila alpestris bilopha</i>	97
<i>Dissura episcopus</i>	488	<i>Eriococcus bambusæ</i>	452
<i>Doicrava affinis</i>	313	— <i>nueræ</i>	452
<i>Drawida longatria</i>	182	— <i>osbeckiæ</i>	452
— <i>peguana</i>	182	— <i>paradoxa</i>	452
— <i>rangoonensis</i>	182	— <i>rhodomyrti</i>	452
— <i>rara</i>	182	— <i>tenius</i>	452
<i>Dremomys lokriah garonum</i>	394	— <i>transversus</i>	452
— <i>subflaviventris</i>	395	<i>Erioides cuneiformis</i>	452
<i>Dromas ardeola</i>	113, 377	— <i>rimulæ</i>	452
<i>Drymoipus insignis</i>	499	<i>Erionota alexandra</i>	440
		— <i>sybirita</i>	440

INDEX OF SPECIES

lxi

	PAGE		PAGE
<i>Erionota thrax acroleuca</i>	440	<i>Felis bengalensis</i>	4, 391, 405
— <i>grandis</i>	440	— <i>caracal</i>	7
— <i>thrax</i>	440	— <i>chaus</i>	7
<i>Erithacus rubecula</i>	104	— <i>lynx var. isabellina</i>	8
<i>Erolia ferruginea</i>	112	— <i>manul</i>	6
— <i>minuta minuta</i>	112	— <i>marmorata</i>	3
<i>Erythropsiza githaginea crassirostris</i> ..	269	— <i>nigrescens</i>	4
<i>Esacus recurvirostris</i>	376	— <i>nipalensis</i>	5
<i>Eschatarchia lineata</i>	143	— <i>ornata</i>	6
<i>Eudynamis scolopaceus scolopaceus</i> ..	477	— <i>pardus</i>	2
<i>Eumenes amedei</i>	246	— <i>rubiginosa</i>	5
— <i>conica</i>	241	— <i>temmincki</i>	4, 392
— <i>dimidiatipennis</i>	244	— <i>tigris</i>	1
<i>Eumenis lehana clarissima</i>	256	— <i>mongolica</i>	1
— <i>droshica</i>	256	— <i>torquata</i>	6
— <i>gilgitica</i>	256	— <i>uncia</i>	3
— <i>mniszeckii baldira</i>	256	— <i>viverinna</i>	4
<i>Eupetaurus cinereus</i>	203	<i>Ferisia virgatus</i>	453
<i>Euphyia scorteia</i>	146	<i>Fiorinia fenestrata</i>	456
<i>Eupithecia acyrtoterna</i>	317	— <i>kandyensis</i>	456
— <i>albispumata</i>	318	— <i>secreta</i>	456
— <i>anasticta</i>	317	<i>Francolinus francolinus</i>	376
— <i>consueta</i>	319	— <i>arabistanicus</i>	118
— <i>eupitheciata</i>	319	— <i>asiæ</i>	483
— <i>filicata</i>	319	— <i>francolinus</i>	118
— <i>hockingii</i>	308	<i>Franklinia buehanani</i>	265, 370
— <i>incurvata</i>	313	<i>Fringilla cœlebs cœlebs</i>	94
— <i>leucenthesia n. sp.</i>	318	— <i>montifringilla</i>	95
— <i>leucostaxis sp. n.</i>	318	<i>Fringillauda brandti hæmatopyga</i> ...	192
— <i>robiginascens</i>	316	— <i>nemoricola altaica</i>	188, 189, 193, 194
— <i>russeola n. sp.</i>	315	<i>Fulica atra atra</i>	526
— <i>sp. n.</i>	316	<i>Fuligula marila</i>	204
— <i>tricrossa n. sp.</i>	316	<i>Funambulus palmarum</i>	403
— <i>ustata</i>	319	— <i>brodiei</i>	239
<i>Eupoditis edwardsi</i>	526	— <i>favonicus</i>	239
<i>Eustroma inextricata</i>	312	— <i>kelaarti</i>	239
<i>Eutyphœus foveatus</i>	182	— <i>matugamensis</i>	239
— <i>peguanus</i>	182	— <i>olympius</i>	239
— <i>rarus</i>	182	— <i>palmarum</i>	239
<i>Exæretopus farinosus</i>	454	— <i>pennanti</i>	381
<i>Falco æsalon æsalon</i>	107	— <i>argentescens</i>	41, 405
— <i>cherrug</i>	107	<i>Galeopterus peninsulæ</i>	46
— <i>jugger</i>	375	<i>Galerida cristata chendoola</i>	273, 373
— <i>peregrinus</i>	481	<i>Gallinago gallinago</i>	112
— <i>peregrinator</i>	481	<i>Gallus g. gallus</i>	376
— <i>severus indicus</i>	481	— <i>sonnerati</i>	488
— <i>subbuteo subbuteo</i>	481	<i>Gangara saguinoculus</i>	439
— <i>tinnunculus interstinctus</i>	481	— <i>thyrsis clothilda</i>	439
<i>Faunis arcesilaus</i>	259	— <i>philippensis</i>	439
— <i>founuloides</i>	260	— <i>thyrsis</i>	439
<i>Felis affinis</i>	41, 381, 391, 403, 405		

	PAGE		PAGE
Gangara thyrsis yasodara	439	Gyps tenuirostris	472
Garrulax albogularis	370	Hadromys humei	400
———— whistleri	461	Halcyon smyrnensis	373
Garrulus glandarius	93	———— fusca	279
———— lanceolatus	370, 460	———— smyrnensis	106, 476
Gazella bennetti	407	Haliaëtus albicillus	479
Gelgeta	437	Haliastur indus indus	479, 526
Gelasma sp.	133	Halimococcus borassi	452
———— chromatocrossa n. sp.	133	Halpe hyrtacus	342
———— dysgenes	133	———— moorei	338
———— fuscifimbria	133	———— (Thoressa) astigmata	345
———— thetydaria	133	———— (Thoressa) honorei	348
Geochelidon anglica	116	Hasora alexis alexis	60
———— nilotica	116	———— badra badra	59
Gennæus hamiltoni	482	———— lanka	59
Geocichla citrina cyanotis	296	———— borneensis	59
———— wardi	467	———— leucospila parnia	60
Geococcus radicum	452	———— lizetta anura	59
Geometra silaceata	311	———— lizetta	59
Gerbillus gleadowi	406	———— moestissima coulteri... ..	60
Glareola lactea	377	———— mus pahanga n. sp.	59
———— nordmanni	112	———— myra	59
———— pratincola pratincola	112, 377	———— padma	61
Glaucidium cuculoides	375	———— proxissima	59
———— cuculoides	477	———— ribbei	60
Glaucion clangula	110	———— salanga	59
Glyphidrilus papillatus	182	———— schonherri schonherri	61
Golunda ellioti	382, 400	———— simplicissima lioneli	60
Gomalia elma albofasciata	79	———— subcœlestis	60
Gomphus bivittatus	425	———— taminatus almea	60
Gonepteryx rhamni burmensis	252	———— bharara	60
———— gilgitica	252	———— malayana	61
Grammatoptila striata striata	462	———— taminatus	60
Grandala cælicolor	466	———— thridas	60
Gunomys bengalensis	381, 397, 406	———— vitta	60
———— varius	47	Hastina azela stenozona	143
———— wardi	406	———— gemmifera	143
Gymnaspis affinis	456	Helaretos malayanus	394
———— bullata	456	Hemichelidon ferruginea	494
———— diospyros	456	———— fuliginosa	494
———— mangifere	456	———— sibirica gulmargi	188, 468
———— producta	456	Hemichionaspis bauhiniæ	455
Gymnopera rubroviridis	320	———— buxi	455
Gymnorhis flavicollis	270	———— ramakrishnæ	455
———— xanthocollis transfuga	95	Hemilecanium imbricans	454
———— xanthocollis	278, 372	Hemithea distinctaria	134
Gypætus barbatus	108	Hemitragus hylocrius	520
———— grandis	478	Herodias alba	488
Gyps fulvus	108, 478	Herpestes auropunctatus	11, 405
———— fulvescens	285	———— pallipes	41
———— himalayensis	478	———— edwardsi	41, 381

INDEX OF SPECIES

[xiii]

	PAGE		PAGE
<i>Herpestes edwardsi ferrugineus</i> ...	381	<i>Hirundo smithi</i> ...	473
----- <i>pallens</i> ...	405	----- <i>filifera</i> ...	271, 372
----- <i>mungo</i> ...	11	<i>Hodgsonius phœnicuroides</i> ...	187, 464
----- <i>nepalensis</i> ...	392	<i>Horisme flavofasciata</i> ...	314
----- <i>urva</i> ...	392	----- <i>hyperythra</i> ...	314
<i>Hesperia alpina alpina</i> ...	81	<i>Horornis albiventris</i> ...	498
----- <i>cashmirensis</i> ...	81	----- <i>pallidus</i> ...	187
----- <i>alveus</i> ...	81	----- <i>pallidus</i> ...	471
----- <i>antonia gigantea</i> ...	80	<i>Hurria rynchops</i> ...	205
----- <i>bieti</i> ...	82	<i>Hyarotis adrastus adrastus</i> ...	441
----- <i>cinaræ</i> ...	81	----- <i>praba</i> ...	441
----- <i>cribrellum</i> ...	80	----- <i>basiflava</i> ...	441
----- <i>evanidus</i> ...	79	----- <i>meluchus</i> ...	441
----- <i>galba</i> ...	79	----- <i>monteithii</i> ...	442
----- <i>maculatus maculatus</i> ...	82	----- <i>pria</i> ...	442
----- <i>thibetanus</i> ...	82	<i>Hydrelia sanguinipaga</i> ...	143
----- <i>malvæ</i> ...	81	<i>Hydrochelidon leucopareia leucopareia</i> ...	114
----- <i>oberthuri</i> ...	81	----- <i>leucoptera</i> ...	114
----- <i>phlomidis geron</i> ...	79	<i>Hyelaphus porcinus</i> ...	401
----- <i>phlomidis</i> ...	79	<i>Hylobates hoolock</i> ...	385
----- <i>poggei</i> ...	80	<i>Hypacanthus spinoides</i> ...	472
----- <i>proto</i> ...	80	<i>Hypenorhynchus erectilineata</i> ...	313
----- <i>sao lugens</i> ...	80	<i>Hypochroma danielaria</i> ...	132
----- <i>serratulæ</i> ...	81	<i>Hypocolius ampelinus</i> ...	99, 492
----- <i>sidæ</i> ...	81	<i>Hypothymis azurea styani</i> ...	495
----- <i>tessellum</i> ...	80	<i>Iambrix latifascia</i> ...	434
----- <i>zebra</i> ...	79	----- <i>salsala luteipalpus</i> ...	433
----- <i>zona</i> ...	82	----- <i>salsala</i> ...	433
<i>Heterophleps clityogramma</i> ...	321	----- <i>sindu</i> ...	434
<i>Hierætus fasciatus</i> ...	479	----- <i>stellifer</i> ...	433
----- <i>pennatus</i> ...	479	----- <i>tytleri</i> ...	433
<i>Himantopus h. himantopus</i> ...	113, 377	----- <i>unicolor</i> ...	433
<i>Hippolais icterina</i> ...	101	<i>Ianthia cyanura pallidiora</i> ...	467
----- <i>obsoleta</i> ...	497	----- <i>rufilata</i> ...	188, 194
----- <i>pallida elæica</i> ...	101, 497	----- <i>pallidiora</i> ...	371
----- <i>rama</i> ..	497	<i>Ibidorhynchus struthersi</i> ...	190, 377, 484
----- <i>scita</i> ...	497	<i>Icerya aegyptiaca</i> ...	451
<i>Hipposideros armiger</i> ...	387	----- <i>jacobsoni</i> ...	451
----- <i>cineraceus</i> ...	387, 404	----- <i>purchasi</i> ...	450
----- <i>fulvus pallidus</i> ..	40, 380, 404	----- <i>seychellarum</i> ...	451
----- <i>gentilis gentilis</i> ...	388	----- <i>var. nardi</i> ...	451
----- <i>leptophyllus</i> ...	388	<i>Ictinaëtus malayensis perniger</i> ...	479
<i>Hipposiderus atratus</i> ...	44	<i>Idæa butyroza</i> ...	138
<i>Hirundinapus caudacuta nudipes</i> ...	476	----- <i>falcipennis</i> ...	142
<i>Hirundo daurica erythropygia</i> ..	271, 372	----- <i>lacernigera</i> ...	308
----- <i>nepalensis</i> ...	225, 271, 473	----- <i>undulataria</i> ...	138
----- <i>rufula</i> ...	105	<i>Inessa ilion</i> ...	435
----- <i>striolata</i> ...	271	<i>Inglisæ bivalata</i> ...	455
----- <i>rustica</i> ...	105, 473	<i>Iodis argutaria</i> ...	135
----- <i>gutturalis</i> ...	270	----- <i>ctila</i> ...	134
----- <i>javanica</i> ...	372	----- <i>lara</i> ...	133

	PAGE		PAGE
<i>Iodis thetydaria</i>	134	<i>Karanasa regeli chitralica</i>	254
<i>Irania gutturalis</i>	104	<i>Kerana armata armata</i>	437
<i>Isma protoclea</i>	443	----- <i>niasana</i>	437
----- <i>sewa</i>	443	----- <i>fulgur</i>	438
----- <i>vulso bicolor</i>	443	----- <i>gimmifer</i>	437
----- <i>purpurascens</i>	443	----- <i>nigrita</i>	437
----- <i>vulso</i>	443	----- <i>vaijrada</i>	437
<i>Ismene amara</i>	63	<i>Kerivoula hardwickei</i>	390
----- <i>anadi</i>	62	<i>Ketupa zeylonensis leschenaulti</i>	106
----- <i>etelka</i>	62	----- <i>semenowi</i>	106
----- <i>formosana</i>	62	<i>Koruthaialos butleri avidha</i>	436
----- <i>gomata gomata</i>	63	----- <i>butleri</i>	436
----- <i>kanara</i>	63	----- <i>haraka</i>	436
----- <i>harisa harisa</i>	62	----- <i>niasicus</i>	436
----- <i>hemixanthus</i>	63	----- <i>focula</i>	436
----- <i>ilusca</i>	61	----- <i>nigerrima</i>	436
----- <i>jaina fergusonii</i>	62	----- <i>rubecula</i>	436
----- <i>jaina</i>	62	----- <i>xanites gopaka</i>	436
----- <i>margana</i>	62	----- <i>javanites</i>	436
----- <i>vasundhara</i>	62	----- <i>kerala</i>	436
----- <i>mahintha</i>	61	----- <i>xanites</i>	436
----- <i>nestor</i>	61	<i>Labioproctus polei</i>	451
----- <i>œdipodea ægina</i>	61	<i>Laccifer albizziaë</i>	451
----- <i>ataphus</i>	61	----- <i>ebrachiata</i>	451
----- <i>œdipodea</i>	61	----- <i>fici</i>	451
----- <i>tuckeri</i>	61	----- <i>lacca</i>	451
----- <i>œdipus</i>	62	<i>Lachnodius humboldtiaë</i>	453
----- <i>phul</i>	62	<i>Laiscopus c. whymperi</i>	372
----- <i>striata</i>	62	<i>Lampropteryx opistholasia</i>	311
----- <i>vasutana</i>	63	<i>Lanius aucheri</i>	495
----- <i>zonaras</i>	61	----- <i>collurio</i>	99
<i>Ithagenes cruentus</i>	503	----- <i>cristatus</i>	469
<i>Itys iadera</i>	442	----- <i>cristatus</i>	496
----- <i>microstictum</i>	442	----- <i>isabellinus</i>	266, 496
<i>Ixobrychus minutus minutus</i>	111	----- <i>phoenicuroides</i>	99, 496
<i>Ixulus flavicollis albicollis</i>	462	----- <i>erythronotus</i>	495
<i>Iynx torquilla japonica</i>	194	----- <i>excubitor</i>	495
<i>Janarda acuminata</i>	142	----- <i>aucheri</i>	99
<i>Jynx torquilla torquilla</i>	105	----- <i>lahtora</i>	495
<i>Karanasa huebneri astorica</i>	254	----- <i>isabellinus</i>	99
----- <i>balti</i>	255	----- <i>lahtora</i>	265
----- <i>calesia</i>	256	----- <i>nigriceps nigriceps</i>	469
----- <i>huebneri</i>	255	----- <i>nubicus</i>	99
----- <i>leechi</i>	255	----- <i>pallidirostris</i>	495
----- <i>modesta</i>	256	----- <i>schach</i>	495
----- <i>pallida</i>	255	----- <i>erythronotus</i>	266, 371, 469
----- <i>moorei gilgitica</i>	254	----- <i>senator niloticus</i>	495
----- <i>moorei</i>	254, 255	----- <i>s. tephronotus</i>	371
----- <i>pamirus pupilata</i>	254	----- <i>tephronotus</i>	495
----- <i>safeda</i>	254	----- <i>vittatus</i>	265, 284, 371, 469
----- <i>regeli bolonicus</i>	254	<i>Larentia ærata</i>	321

INDEX OF SPECIES

lxv

	PAGE		PAGE
<i>Larus argentatus vegæ</i>	114	<i>Limosa lapponica lapponica</i>	113
----- <i>cachinnans</i>	114	<i>Lioopicus mahrattensis</i>	475
----- <i>gelastes</i>	114, 377	<i>Liponycteris kachhensis</i>	380
----- <i>genei</i>	114	<i>Lithotis rupicola</i>	447
<i>Larvivora brunnea</i>	464	----- <i>tumida</i>	449
<i>Leachia festiva</i>	451	<i>Lobivanellus indicus indicus</i>	377, 483
<i>Lecaniodiaspis mimusopis</i>	455	<i>Locustella lanceolata</i>	497
<i>Lecanium bicruciatum</i>	454	----- <i>nævia straminea</i>	497
----- <i>desolatum</i>	454	<i>Lonicera spinosa</i>	186
----- <i>fusiforme</i>	454	<i>Lophomackia albiradiata</i>	132
----- <i>illuppalamæ</i>	454	----- <i>semialba</i>	132
----- <i>imbricans</i>	454	<i>Lophophanes melanolophus</i>	460
----- <i>ixoræ</i>	454	----- <i>rufonuchalis rufonucha-</i>	
----- <i>latio perculatum</i>	454	<i>lis</i>	461, 491
----- <i>limbatum</i>	454	<i>Lophophorus impejanus</i>	188, 482
----- <i>mancum</i>	454	<i>Lotongus avesta</i>	446
----- <i>magniferæ</i>	454	----- <i>calathus</i>	446
----- <i>marsupiale</i>	454	----- <i>onara</i>	446
----- <i>piperis var. namunakuli</i>	454	----- <i>sarala</i>	446
----- <i>ramakrishnæ</i>	454	----- <i>schoedia</i>	446
----- <i>tessalatum</i>	454	<i>Loxia curvirostra himalayana</i>	471
----- <i>var. obsoletum</i>	454	<i>Lullula arborea pallida</i>	96
----- <i>trifasciatum</i>	454	<i>Luscinia luscinia</i>	104
----- <i>tripartitum</i>	454	----- <i>megarhyncha apicana</i>	104, 493
<i>Lecanopsis ceylonica</i>	454	----- <i>golzii</i>	493
<i>Lefroyia castenea</i>	452	----- <i>suecica magna</i>	104
<i>Leggada booduga</i>	382, 399	<i>Lusciniola melanopogon</i>	497
----- <i>jacksoniæ</i>	399	----- <i>mimica</i>	100
----- <i>nagarum</i>	399	<i>Lutra lutra nair</i>	393
----- <i>pahari</i>	400	<i>Lutrogale barang tarayensis</i>	393
<i>Leggadilla sadhu</i>	42, 382	<i>Lygranoa cinerea</i>	322
<i>Leioptila c. pallida</i>	370	----- <i>erectilineata</i>	313
----- <i>p. psaroides</i>	370	<i>Lyroderma lyra</i>	380, 388, 402
<i>Lepidosaphes albizziæ</i>	456	<i>Macaca adusta</i>	43
----- <i>dilatilobis</i>	456	----- <i>assamensis</i>	385
----- <i>leucooplææ</i>	456	----- <i>irus</i>	43
----- <i>punctatissima</i>	456	----- <i>mulatta</i>	385
----- <i>punicæ</i>	456	<i>Machetes pugnax</i>	113
<i>Lepidosiren</i>	503	<i>Machlolophus xanthogenys xanthogenys</i>	461
<i>Leptocoma asiatica asiatica</i>	282	<i>Maniola cheena cheena</i>	258
<i>Lepus dayanus</i>	42, 407	----- <i>chitralica</i>	258
----- <i>ruficaudatus</i>	382, 401	----- <i>kashmirica</i>	258
<i>Lerwa lerwa</i>	483	----- <i>pulchra astorica</i>	258
<i>Lethe armandii armandii</i>	259	----- <i>baroghila</i>	257
----- <i>fusca</i>	259	----- <i>chitralica</i>	257
----- <i>khasiana</i>	259	----- <i>pulchra</i>	258
----- <i>lacticolora</i>	259	<i>Manis pentadactyla</i>	402
----- <i>oberthur</i>	259	<i>Mareca penelope</i>	485
----- <i>saggitata</i>	259	<i>Marmaronetta angustirostris</i>	485
<i>Leuconoe</i>	389	<i>Marmota himalayana</i>	403
<i>Limnocyba indica</i>	447, 449	<i>Matapa aria</i>	441

	PAGE		PAGE
Matapa celsina	441	Mirafra erythroptera	273
----- druna	441	Molpastes hæmorrhous... ..	492
----- purpurascens	441	----- hæmorrhous	291
----- sasivarna	441	----- pallidus	264, 279
----- shalgrama	441	----- leucogenys leucogenys	463
Matapoides smaragdinus	440	----- leucotis	279, 492
Megaderma spasma majus	388	----- leucotis leucotis	464
----- spasma medium	44	Monophlebus phyllanthi	450
Megalæma virens marshallorum	475	----- zeylanicus	451
Megalornis antijone	378	Monticola cinclorhyncha	372, 463
----- grus	112	----- erythrogastra	467
----- leucogeranus	378	----- longirostris	494
Megascolex mauritii	182	----- pandoo	494
Melanippe artemidora	321	----- saxatilis	103, 494
----- latifusata	144	----- solitaria	494
Melanocorypha calandra calandra	96	----- pandoo	190, 269, 468
----- hebraica	96	----- solitarius	372
----- psammochroa	96	----- longirostris	103
Melia azadarach	262	----- transcaspicus	103
----- azadirachta	262	Montifringilla brandti hæmatopygia... ..	473
Melogale personata	393	----- n. adamsi	273, 472
Melophus melanicterus	473	----- remoncola altaica	472
Menetes berdmorei	47	Mooreana trichoneura pralaya	73
Mergus albellus	110	----- trichoneura	73
----- serrator	110	Moschus moschiferus	401
Merops apiaster	106	Motacilla alba	98
----- javanicus... ..	373	----- alboides	474
----- orientalis... ..	475	----- dukhunensis	98, 271, 473
----- persicus persicus	106, 373	----- hodgsoni	191
Metacardia conchiferata	451	----- persica	98
Metallolophia arenaria	132	----- personata	98, 473
Metoponia pusilla	189, 190, 192	----- cinerea	98
Microcichla scouleri scouleri	465	----- caspica	98
Microloba bella diacena	321	----- melanope	372, 474
Micropus affinis	476	----- citreola citreola	192, 474
----- a. galilejensis	373	----- citreoloides	188
----- apus pellinensis	476	----- c. calcarata	372
----- melba	476	----- feldegg melanogrisea	271
----- murinus murinus	106	----- flava thunbergi	98
----- pacificus pekinensis	476	----- maderaspatensis	271, 474
Microscelis psaroides ganeesa	295	----- melanope	98
----- psaroides	462	Muntiacus vaginalis	401
Microtarsus melanocephalus	492	Murina cyclotis	389
Microtus (Hyperacrius) wynnei	407	----- tubinaris	389
----- melanogaster	400	Mus dubius	399
Millardia meltada	382, 406	----- gerbillus	406
Milvus lineatus	108, 479	----- gliroides	400
----- migrans govinda	375, 479, 524	----- homourus	382, 407
----- migrans	108	----- musculus	399
Mirafra assamica	373	----- urbanus	382, 407
----- cantillans cantillans	287	Muscicapa hypoleuca semitorquata	100

	PAGE		PAGE
Muscicapa striata neumanni ...	100, 494	Nyroca ferina	485
Mycalesis suaveolens duguidi ...	259	----- fuligula	485
----- sebonga ...	259	----- rufa	485
Mycerobas carnpes	471	Ochotona roylei	403
----- melanoxanthus	471	Octochaetus birmanicus... ..	182
Myiophoneus horsfieldii	293	Ochromela nigrorufa	300
----- temmincki	188	Ochus subvittatus	430
----- temmincki	372	Odina decoratus	73
----- temmincki	468	----- hieroglyphica ortygia	73
Myotis hasselti	45	Odontoptilum angulata	77
----- muricola	45	----- pygela	78
----- siligorensis	389	Ænanthe (see Ænanthe.)	
Naia hanna	226	----- alboniger	492
Nelumbium speciosum	262	----- deserti	465
Nemorhædus goral	203	----- oreophila	493
Neobalbis elæaria	131	----- isabellina	268
Neolecanium crustutiforme	455	----- melanoleuca melanoleuca	492
Neoleucaspis parallela	455	----- picata	268, 492
Neophron percnopterus	478	----- xanthoprymna chrysopygia.	493
----- ginginianus... ..	284	Oeneis garhwalia n. sp.	253
----- percnopterus.	108	Oerane neæra	441
Neritina perottetiana	448	Opuntia dillenii	269
Nesokia indica	42	Oreicola ferrea ferrea	187, 464
Netta rufina	485	Oreocinclæ dauma dauma	467
Nettion crecca	485	----- mollissima	467
Nettopus coromandelianus	484	Oreocorys sylvanus	474
Nietnera pundaluoya	451	Organopoda brevipalpis	135
Niltava sundara sundara	469	Oriolus luteolus luteolus	471
Nisoniades marloyi marloyi	83	----- oriolus kundoo	266, 471
----- pelias	83	----- oriolus	94
----- montanus montanus	83	----- s. kundoo	371
----- nigrescens	83	Ortholitha latifusata	144
----- tages	83	Orthopætus lalita	64
Notocrypta curvifascia... ..	439	----- lidderdali	64
----- feisthamelii avattana	439	----- phanæus	64
----- paralysos alysia n. sp.	438	Orthotomus sutorius sutorius	264, 470, 497
----- devadatta	439	Otis tarda	112
----- paralysos	439	----- tetrax	112
----- sidha	439	Otocompsa emeria emeria	492
----- quadrata	438	----- fuscicaudata	291, 370
----- renardi	439	Otoconys alpestris elwesi	474
----- volux niasana	438	----- longirostris	189, 191, 373, 474
----- volux	438	Otogyps calvus	375
Nucifraga caryocatactes hemispila	460	Otus brucei	107
----- hemispila	370	Ovis vignei	174
Numenius arquata lineatus	43	----- cycloceros	401
Nycticebus coucang	43	Pachyodes arenaria	132
Nycticejus castaneus	45	Pachyura hodgsoni	301
----- ornatus	389	----- sp.	46, 380, 391, 404
Nycticorax nycticorax nycticorax	111, 204	Padraona gola	334
Nyctocleptes cinereus	48		

	PAGE		PAGE
<i>Paduka lebadea andamanica</i> ..	440	<i>Parus major blanfordi</i> ...	98
————— <i>glandulosa</i> ...	440	————— <i>melanolophus</i> ...	370
————— <i>lebadea</i>	440	————— <i>monticola</i> ...	460
————— <i>subfasciata</i> ...	440	————— <i>m. intermedius</i> ...	491
————— <i>toradja</i>	440	————— <i>m. kashmiriensis</i> ...	187, 460
<i>Paguma grayi</i>	10, 392, 403	————— <i>m. mahrattarum</i> ...	301
<i>Paludomus</i>	449	————— <i>m. planorum</i> ...	491
<i>Pandion haliaëtus</i>	477	————— <i>palustris korejewi</i> ...	491
<i>Panicum javanicum</i>	530	<i>Passer domesticus biblicus</i> ...	95
<i>Papilio noblei haynei</i>	249	————— <i>indicus</i> ...270, 372, 472	
<i>Paradoxurus crossi</i>	9	————— <i>parkini</i>	190
————— <i>hermaphroditus</i> ...	9, 10	————— <i>hispaniolensis transcaspicus</i> ...	95
————— <i>niger</i>	9, 381	————— <i>moabiticus moabiticus</i> ...	95
————— <i>strictus</i>	10, 392	————— <i>pyrrhonotus</i>	372
<i>Paræchinus blanfordi</i>	404	————— <i>rutilans debilis</i> ...	188, 472
<i>Paranticopsis xenodes prontis</i> ...	249	<i>Pastor roseus</i>	94, 267
<i>Pararge menava mœroides</i>	253	<i>Pedestes bononoides</i>	432
<i>Parascaptor leucurus</i>	390	————— <i>masuriensis</i>	432
<i>Parazoma</i> Gen. n.	314	————— <i>pandita</i>	432
————— <i>ferax</i> n. sp.	314	<i>Pedronia strobilanthi</i>	452
————— <i>semifusca</i>	315	<i>Prionodon pardicolor</i>	392
<i>Parlatoria blanchardii</i>	457	<i>Pelecanus crispus</i>	522
————— <i>camelliæ</i>	457	————— <i>onocrotalus onocrotalus</i> 108,	522
————— <i>cingala var. namunakuli</i> ...	457	<i>Pellorneum ruficeps granti</i>	293
————— <i>cinnamoma</i>	457	————— <i>ruficeps</i>	294
————— <i>cristifera</i>	457	<i>Perdicula asiatica</i>	376
————— <i>limoniæ</i>	457	<i>Pericrocotus brevirostris brevirostris</i> . 266,	469
————— <i>mesuæ</i>	457	————— <i>erythropygius</i>	266
————— <i>proteus var. crotonis</i> ...	457	————— <i>peregrinus malabaricus</i> . 284	
————— <i>rutherfordi</i>	457	————— <i>peregrinus</i> 266, 496	
————— <i>zeylanica</i>	457	————— <i>pallidus</i>	496
<i>Parnassius delphius atkinsoni</i> ...	251	————— <i>roseus roseus</i>	469
————— <i>chitralica</i>	250	————— <i>speciosus</i>	469
————— <i>florenciæ</i>	251	<i>Perionyx excavatus</i>	180, 182
————— <i>rileyi</i>	251	————— <i>fulvus</i> (... ..	180, 182
————— <i>stoliczkana</i>	250	<i>Perizoma albofasciata</i>	309
————— <i>zogilaica</i>	251	————— <i>bipartaria</i>	310
————— <i>discobulus</i>	249	————— <i>cerva</i>	308
————— <i>baroghila</i>	250	————— <i>conjuncta</i>	309
————— <i>hunzaica</i>	250	————— <i>decorata quadrinotata</i> ...	309
————— <i>dongalaica</i>	249	————— <i>hockingii</i>	308
————— <i>jacquemontii shandura</i> ...	250	————— <i>lacer nigra</i>	308
————— <i>simo lorimeri</i>	252	————— <i>latifasciata</i>	309
<i>Parus coeruleus georgicus</i>	99	————— <i>lucifrons</i>	309
————— <i>cinereus</i>	491	————— <i>maculata</i>	309
————— <i>c. kashmiriensis</i>	370	————— <i>puerilis</i>	308
————— <i>kashmiriensis</i>	491	————— <i>schistacea</i>	308
————— <i>lugubris</i>	98	————— <i>seriata</i>	310
————— <i>anatolix</i>	98	————— <i>variabilis</i>	310
————— <i>dubius</i>	98	<i>Pernis apivorus apivorus</i>	108
————— <i>mahrattarum</i>	491	<i>Perrisospiza icteroides</i>	471

	PAGE		PAGE
<i>Petaurista inornatus</i>	403, 405	<i>Phylloscopus nitidus nitidus</i>	265
----- <i>mergulus</i> n. sp.	46	----- <i>viridanus</i>	470
----- <i>primrosei</i> , subsp. n. 22, 46		----- <i>occipitalis occipitalis</i>	470
----- <i>reguli</i> , subsp. n. 22, 46		----- <i>proregulus simlaensis</i>	470
----- <i>yunnanensis</i>	394	----- <i>pulcher kangræ</i>	470
<i>Petronia petronia exigua</i>	95	----- <i>sindianus</i>	498
<i>Petrophila cinclorhyncha</i>	187	----- <i>tristis</i>	498
<i>Phalacrocorax carbo</i>	109, 484	----- <i>trochiloides trochiloides</i>	470
----- <i>pygmæus</i>	109	----- <i>trochilus trochilus</i>	100
<i>Phasianus humiæ</i>	204	----- <i>tytleri</i>	188, 498
<i>Phenacoccus ballardi</i>	453	<i>Physetobasis dentifascia</i>	315
----- <i>glomeratus</i>	453	<i>Pica pica bactriana</i>	93, 490
----- <i>hirsutus</i>	453	<i>Picumnus innominatus innominatus</i>	475
----- <i>ornatus</i>	453	<i>Picus canus occipitalis</i>	475
----- <i>spinosus</i>	453	----- <i>occipitalis</i>	373
<i>Pheretima anomala</i>	182	----- <i>squamatus squamatus</i>	475
----- <i>elongata</i>	182	<i>Pinnaspis bauhiniæ</i>	455
----- <i>houlletti</i>	182	----- <i>buxi</i>	455
----- <i>insolita</i>	182	----- <i>ramakrishnæ</i>	455
----- <i>lignicola</i>	182	<i>Pinus longifolia</i>	225
----- <i>peguana</i>	182	<i>Pipistrellus babu</i>	380, 388
----- <i>planata</i>	182	----- <i>coromandra</i>	45, 388
----- <i>posthuma</i>	182	----- <i>kuhli lepidus</i>	40
<i>Phibalapteryx eupitheciata</i>	319	----- <i>mimus</i>	41, 380, 388, 404
----- <i>hyperythra</i>	314	<i>Pithecus johnii</i>	223
<i>Phœnicopterus ruber antiquorum</i>	110, 522	----- <i>obscurus</i>	43
<i>Phœnicurus erythrogaster grandis</i>	493	----- <i>pileatus</i>	386
----- <i>frontalis</i>	194	----- <i>saturatus</i>	386
----- <i>ochrurus phœnicuroides</i>	104, 188, 192, 268, 371, 465	----- <i>tenebricus</i>	386
----- <i>phœnicurus</i>	104	<i>Planesticus maximus</i>	196
----- <i>ph. mesoleuca</i>	104	<i>Plastingia aurantiaca</i>	444
<i>Phoenix sylvestris</i>	262	----- <i>callineura burmana</i>	444
<i>Photoscotosia miniosata</i>	312	----- <i>callineura</i>	444
<i>Phyletis similata</i>	135	----- <i>perakana</i>	444
<i>Phyllorhina leptophyllus</i>	388	----- <i>corissa corissa</i>	445
<i>Phylloscopus affinis</i>	192, 371, 470	----- <i>tytleri</i>	445
----- <i>collybita</i>	189, 498	----- <i>helena fruhstorferi</i>	444
----- <i>abietina</i>	100	----- <i>helena</i>	444
----- <i>tristis</i>	265	----- <i>latoia latoia</i>	444
----- <i>griseolus</i>	498	----- <i>margherita</i>	444
----- <i>humii</i>	194, 497	----- <i>naga</i>	445
----- <i>humii</i>	265	----- <i>noëmi noëmi</i>	445
----- <i>præmium</i>	498	----- <i>tavoyana</i>	445
----- <i>superciliosus</i>	498	----- <i>pugnans</i>	445
----- <i>indicus</i>	265	----- <i>sala</i>	445
----- <i>inornatus</i>	498	----- <i>submaculata kanara</i>	445
----- <i>humii</i>	470	----- <i>submaculata</i>	445
----- <i>maculipennis</i>	470	----- <i>tessellata</i>	445
----- <i>mandellii</i>	498	----- <i>vermiculata</i>	444
----- <i>neglectus</i>	498	<i>Platalea leucorodia major</i>	111
		<i>Platygomphus dolobratus</i>	416
		----- <i>feæ</i>	417

	PAGE		PAGE
<i>Plecotus wardi</i>	404	<i>Pseudococcus kandyensis</i>	453
<i>Ploceus bengalensis</i>	269	————— <i>maritimus</i>	453
<i>Ploceus manyar peguensis</i>	288	————— <i>monticola</i>	453
————— <i>flaviceps</i>	269, 372	————— <i>pulverarius</i>	453
————— <i>philippinus philippinus</i> 269, 284, 372		————— <i>bambusæ</i>	453
<i>Pnœpyga albiventer pallidior</i>	463	————— <i>virgatus</i>	453
<i>Podiceps cristatus cristatus</i>	116	<i>Pseudogyps bengalensis</i> ... 285, 375, 478	
————— <i>nigricollis nigricollis</i>	116	<i>Pseudoparlatoria pusilla</i>	457
————— <i>ruficollis capensis</i>	116	<i>Pseudopulvinaria castenea</i>	452
————— <i>iraquensis</i>	116	<i>Pseudotantalus leucocephalus</i>	488
————— <i>ruficollis</i>	116	<i>Pseudoterpra elæaria</i>	131
<i>Polea ceylonica</i>	452	<i>Psittacula cyanocephala cyanocephala</i> 477	
<i>Polioaëtus humilis</i>	479	————— <i>cupatria nepalensis</i>	477
<i>Pollinia ceylonica</i>	452	————— <i>schisticeps schisticeps</i>	477
<i>Polyphasia dentifera</i>	312	<i>Pterocles alchata caudacutus</i> ... 117, 376	
<i>Polyplectron bicalcaratum</i>	204	————— <i>senegalensis</i>	375
<i>Pomatorhinus erythro-</i>		————— <i>senegallus</i>	117, 376
<i>genys</i>	370, 462	<i>Pteromys (Hylopetes) alboniger</i> ... 394	
<i>Pontoscolex corethrurus</i>	182	————— <i>inornatus</i>	195
<i>Porzana parva</i>	117	<i>Pteropus edulis</i>	44
————— <i>porzana</i>	116	————— <i>giganteus</i> ... 379, 386, 402	
<i>Prinia gracilis</i>	102	————— <i>hypomelanus geminorum</i> ... 44	
————— <i>irakensis</i>	102	————— <i>intermedius</i>	44
————— <i>lepida</i> ... 102, 265, 371, 499		————— <i>nicobaricus</i>	44
————— <i>palestinæ</i>	102	————— <i>vampyrus</i>	44
————— <i>inornata inornata</i> . 265, 284, 291, 371		————— <i>malaccensis</i>	44
————— <i>jerdoni</i>	291	<i>Pteruthius xanthochloris occidentalis</i> 462	
————— <i>socialis stewarti</i>	265, 371	<i>Ptychopoda aeqinsinuata</i>	141
————— <i>sylvatica insignis</i>	499	————— <i>complanata</i>	138
————— <i>rufescens</i>	499	————— <i>quingestriata</i>	138
————— <i>sylvatica</i>	499	————— <i>muricolor</i>	140
<i>Propasser rhodochrous</i> 194, 195, 196		————— <i>semilinea</i>	140
————— <i>thura</i>	194, 196	————— <i>semisericea</i>	140
<i>Prosopis spicigera</i>	262	<i>Ptyonoprogne concolor</i> ... 270, 287	
<i>Protopterus</i>	503	————— <i>obsoleta obsoleta</i> ... 105	
<i>Prunella blanfordi</i>	104	————— <i>rupestris</i>	189, 191
————— <i>collaris</i>	104, 468	<i>Pucrasia macrolopha</i>	482
————— <i>caucasicus</i>	104	<i>Pudicitia pholus</i>	440
————— <i>himalayanus</i>	468	<i>Pulvinaria cellulosa</i>	455
————— <i>modularis</i>	104	————— <i>maxima</i>	455
————— <i>orientalis</i>	104	————— <i>polygonata</i>	455
————— <i>rubeculoides</i> ... 189, 192, 372		<i>Pycnonotus erythrophthalmus</i> ... 492	
————— <i>strophciata jerdoni</i>	494	————— <i>h. hæmorrhous</i>	370
<i>Psara phœopteralis</i>	529	————— <i>h. intermedius</i>	370
<i>Pseudantonina bambusæ</i>	452	————— <i>l. leucogenys</i>	370
<i>Pseudococcus citriculus</i>	453	————— <i>leucotis mesopotamiæ</i> ... 100	
————— <i>comstocki</i>	453	<i>Pyctorhis sinensis</i>	370
————— <i>corymbatus</i>	453	<i>Pyrrhocorax graculus</i> ... 460, 491	
————— <i>crotonis</i>	453	————— <i>pyrrhocorax</i> ... 93, 191, 460	
————— <i>debregeasiæ</i>	453	<i>Pyrrhospiza punicea humii</i> ... 192, 193, 477	
————— <i>detoquens</i>	453	<i>Pyrrhula aurantiaca</i> ... 188, 195, 196	

	PAGE		PAGE
<i>Pyrrhula erythrocephala</i>	471	<i>Rhopalocampta benjaminii benjaminii</i>	63
<i>Pyrrhulauda frontalis</i>	96	————— <i>crawfurdi</i> .	63
<i>Pyrrhulauda grisea grisea</i>	273, 284, 373	<i>Rhopocichla atriceps atriceps</i> ...	293
<i>Python curtus</i>	84	<i>Rhyacornis fuliginosus</i>	371, 466
———— <i>molurus</i>	84	<i>Runeca ferrilineata</i>	136
———— <i>reticulatus</i>	84	<i>Rusa unicolor</i>	402
———— <i>timorensis</i>	84	<i>Saccharum arundinaceum</i>	262
<i>Querquedula crecca</i>	110	<i>Salpornis spilonota</i>	264
———— <i>querquedula</i>	485	<i>Salvadora persica</i>	262
<i>Rallus aquaticus aquaticus</i>	234	<i>Sancus pulligo pulligo</i>	437
———— <i>indicus</i>	234	———— <i>subfasciatus</i>	437
———— <i>korejeiwi</i>	234	<i>Sarangesa dasahara albicilia</i>	76
———— <i>indicus</i>	234	———— <i>dasahara</i>	76
<i>Rana tigrina</i>	228	———— <i>davidsoni</i>	76
<i>Rattus alexandrinus</i>	406	———— <i>purendra</i>	76
———— <i>berdmorei</i>	47	———— <i>sati hopkinsi</i>	76
———— <i>concolor</i>	47	———— <i>sati</i>	76
———— <i>fulvescens</i>	398	<i>Sarcidiornis melanota</i>	284, 287
———— <i>listeri</i>	398	<i>Sarcogrammus indicus aigneri</i>	113
———— <i>garonum</i>	398	<i>Satarupa gopala gopala</i>	70
———— <i>manipulus</i>	398	———— <i>nymphalis</i>	70
———— <i>nitidus</i>	398	———— <i>sambara affinis</i>	70
———— <i>rattus arboreus</i>	403	———— <i>dohertyi</i>	70
———— <i>rufescens</i>	381, 406	———— <i>formosana</i>	70
———— <i>sikkimensis</i>	398	———— <i>javanensis</i>	70
———— <i>tikos</i>	48	———— <i>niphates</i>	70
———— <i>tistæ</i>	399	———— <i>sambara</i>	70
———— <i>wellsi</i>	399	———— <i>strigata</i>	70
———— <i>surifer</i>	48	<i>Saxicola caprata</i>	492
———— <i>vicereus</i>	406	———— <i>atrata</i>	291
———— <i>vociferans</i>	48	———— <i>bicolor</i>	268, 371, 464
<i>Ratufa gigantea</i>	394	———— <i>burmanica</i>	492
———— <i>melanopepla celænopepla</i>	46	———— <i>rubetra noskœ</i>	104
<i>Regulus regulus himalayensis</i> ...	499	———— <i>torquata</i>	104
<i>Remiz coronatus</i>	491	———— <i>indica</i>	268, 371, 464
<i>Riparia brevicaudata</i>	270	<i>Saxicoloides fulicata cambaiensis</i> .	269, 282, 467
———— <i>chinensis</i>	372	<i>Sciurus assamensis</i>	396
<i>Rhinolophus affinis</i>	386	———— <i>blythi</i>	396
———— <i>himalayanus</i>	386	———— <i>domelicus</i>	47
———— <i>blythi szechuanus</i>	386	———— <i>erythræus</i>	395
———— <i>ferum</i>	387	———— <i>lokroïdes</i>	396
———— <i>lepidus</i>	387	———— <i>subflaviventris</i>	395
———— <i>pearsoni</i>	387	———— <i>sullivanus</i>	47
———— <i>perniger</i>	387	<i>Scobura cephalæ</i>	435
———— <i>yunanensis</i>	387	———— <i>cephaloides</i>	435
<i>Rhinopoma hardwickei</i>	380, 404	———— <i>coniata</i>	435
———— <i>xanthocomes</i> n. sp.	320	———— <i>pheditia</i>	435
<i>Rhipidura aureola aureola</i>	268, 284, 371	<i>Scolopax rusticola</i>	112, 377, 484
———— <i>pectoralis</i>	289	<i>Scops plumipes</i>	477
<i>Rhizomys pruinosus</i>	400	<i>Scopula bispurcata</i>	137
<i>Rhodospiza obsoleta</i>	94	———— <i>brachypus</i>	139
<i>Rhodostophia similata</i>	135		

	PAGE		PAGE
Scopula butyrosea	138	Sphenocercus sphenurus	486
——— complanata	138	Spheg flavipennis	151
——— dentata	138	——— lobatus	147, 157
——— ferrilineata	136	Spizaetus nipalensis nipalensis	479
——— marcidaria	137	Stachyridopsis pyrrkops	370, 461
——— patularia	139	Stercorarius parasiticus	114
——— quinquestriata	138	Sterna albifrons albifrons	115
——— segregata	136	——— albigena	115
——— straminea	139	——— anæthetus fuligula	116
——— undulataria	138	——— a. saundersi	377
——— unisignata	137	——— bengalensis bengalensis	115, 377
——— sp. n.	137	——— bergii velox	115
Scotomanes ornatus imbreus	389	——— caspia caspia	115, 377
Scotophilus castaneus	45	——— hirundo hirundo	191, 192
——— kuhli ... 41, 380, 389, 402, 404		——— melanogaster	377
——— temminckii	45	——— minuta	115
——— wroughtoni ... 380, 389, 402		——— nilotica	377
Scotosia indistincta	322	——— repressa	115
——— lativittaria	311	Sterrhia actiosaria	140
——— miniosata	312	——— acuminata	142
Seicercus burkii whistleri	470	——— acquisinuata	141
——— xanthoschistus albosuperciliaris		——— castelli	141
	470, 498	——— falcipennis	142
Sepa bononia	444	——— ingloria n. sp.	140
——— cronus	443	——— lamprotis sp. n.	142
——— dawna n. sp.	443	——— muricolor	140
——— fenestrata	443	——— ocnerya sp. n.	142
——— feralia	443	——— rubridentata	142
——— inarime	444	——— semilinea	140
——— miosticta	443	——— smisericea	140
Serinus pusillus	94, 372, 472	——— spp.	140
——— syriacus	94	Stoparola albicaudata	298
Sesquiptera inaequata	320	——— melanops melanops...268, 371, 469	
Silybura sp.	528	Streptopelia chinensis suratensis	281, 482
Siphia parva hyperythra	187	——— decaocto decaocto. 117, 281, 482	
——— parva	268	——— orientalis meena ... 191, 375, 482	
——— strophata	468	——— senegalensis cam-	
Sitta castaneiventris castaneiventris	264, 370	bayensis ... 117, 281, 482	
——— frontalis	491	——— suratensis	375
——— ——— corallina	491	——— turtur arenicolor... ..	117
——— himalayensis	461	Strix aluco nivicola	477
——— kashmiriensis	188	——— sanctinicolai	107
——— leucopsis leucopsis	194, 461	Struthio camelus syriacus	118
——— neumayer	98	Sturnopastor contra contra	267, 371
——— ——— dresseri	98	Sturnus sophiæ	93
——— ——— kurdistanica	98	——— vulgaris	93
——— rupicola tschitscherini	98	——— nobilior	93
Siva strigula strigula	462	——— oppenheimi	93
Soriculus caudatus	390	——— poltaratzskii... ..	261
——— radulus n. sp.	390	——— purpurascens	93
Spatula clypeata	485	Suada swerga cataleucos	438

	PAGE		PAGE
<i>Suada swerga swerga</i>	435	<i>Tagiades obscurus alica</i>	72
— — — — — <i>triplex</i>	435	— — — — — <i>athos</i>	71
<i>Suastus everyx everyx</i>	434	— — — — — <i>distans</i>	71
— — — — — <i>scopas</i>	435	— — — — — <i>meetana</i>	72
— — — — — <i>gremius gremius</i>	434	— — — — — <i>obscurus</i>	72
— — — — — <i>subgrisea</i>	434	— — — — — <i>patimoka</i>	72
— — — — — <i>minuta</i>	434	— — — — — <i>sangarava</i>	72
— — — — — <i>plana</i>	435	— — — — — <i>sumbawana</i>	72
— — — — — <i>rama aditus</i>	434	— — — — — <i>toba</i>	72
— — — — — <i>bipunctus</i>	434	— — — — — <i>waterstradti</i>	72
— — — — — <i>rama</i>	434	<i>Takahashia japonica</i>	455
<i>Succinea arboricola</i>	447	<i>Talpa micrura</i>	390
<i>Suya crinigera crinigera</i>	471	<i>Tamarindus indica</i>	262
— — — — — <i>striatula</i>	499	<i>Tamarix dioica</i>	262
<i>Sylvia affinis</i>	497	<i>Tamiops maclellandi</i>	397
— — — — — <i>althea</i>	191	— — — — — <i>barbei</i>	47
— — — — — <i>atricapilla atricapilla</i>	101	<i>Tapena atilia</i>	77
— — — — — <i>communis icterops</i>	101, 497	— — — — — <i>palawana</i>	77
— — — — — <i>conspicillata conspicillata</i>	101	— — — — — <i>thwaitesi hampsoni</i>	77
— — — — — <i>curruca affinis</i>	265, 497	— — — — — <i>minuscule</i>	77
— — — — — <i>minula</i>	497	— — — — — <i>thwaitesi</i>	77
— — — — — <i>hortensis</i>	101	<i>Taphozous longimanus</i>	380
— — — — — <i>crassirostris</i>	101	<i>Tarsiger chrysæus whistleri</i>	467
— — — — — <i>melanocephala</i>	101	<i>Tatera indica</i>	381, 406
— — — — — <i>mystaica</i>	101	— — — — — <i>sherrini</i>	41
— — — — — <i>nana nana</i>	101, 497	<i>Tchitrea paradisi</i>	371
— — — — — <i>nisoria</i>	101	<i>Telicota augias</i>	326
— — — — — <i>simplex</i>	101	— — — — — <i>bambusæ</i>	325
— — — — — <i>pallida</i>	101	— — — — — <i>mæsoides</i>	327
<i>Synegiodes hyriaria</i>	136	— — — — — <i>palmarum</i>	326
<i>Syrnium biddulphi</i>	196	<i>Temenuchus pagodarum</i>	267, 286, 371, 471
— — — — — <i>nivicola</i>	196	<i>Temnogomphus bivittatus</i>	425
<i>Syrrhaptis paradoxus</i>	204, 522	<i>Tephrodornis pondicerianus</i>	266
<i>Tachardina decorella</i>	451	<i>Terekia cinerea</i>	112
— — — — — <i>lobata</i>	451	<i>Terpna mölleri</i>	132
— — — — — <i>minuta</i>	451	— — — — — <i>vigil n. sp.</i>	131
— — — — — <i>ternata</i>	451	<i>Terpsiphone paradisi paradisi</i> 268, 469, 495	
— — — — — <i>theæ</i>	451	<i>Tesia castaneocoronata castaneocoro-</i>	
<i>Tadorna tadorna</i>	109, 224	— — — — — <i>nata</i>	463
<i>Tagiades atticus atticus</i>	71	<i>Tetraogallus himalayensis</i>	376, 483
— — — — — <i>helferi</i>	71	— — — — — <i>tibetanus</i>	503
— — — — — <i> khasiana</i>	71	<i>Thalassodes aptifimbria</i>	134
— — — — — <i>rajaghra</i>	71	— — — — — <i>distinctaria</i>	134
— — — — — <i>calligana</i>	73	<i>Thalera argutaria</i>	135
— — — — — <i>gana gana</i>	71	<i>Thamnobia f. cambayensis</i>	371
— — — — — <i>menanto</i>	71	<i>Thera cyphoschema n. sp.</i>	312
— — — — — <i>japetus</i>	72	<i>Threskiornis æthiopicus</i>	111
— — — — — <i>lavata</i>	71	<i>Tichodroma muraria</i>	98, 463
— — — — — <i>litigiosa litigiosa</i>	72	<i>Timandra correspondens</i>	136
— — — — — <i>vajuna</i>	72	— — — — — <i>oligoscia</i>	136
— — — — — <i>menaka</i>	72	<i>Tomeutes blythi</i>	396

	PAGE		PAGE
<i>Tomeutes bellona</i>	397	<i>Turdus merula</i>	493
————— <i>mearsi</i>	397	————— <i>intermedius</i>	102
————— <i>stevensi</i>	397	————— <i>maximus</i>	188
————— <i>virgo</i>	397	————— <i>syriaca</i>	102
————— <i>lokroides</i>	396	————— <i>obscurus subobscurus</i>	493
<i>Torgos calvus</i>	285, 478	————— <i>philomelos philomelos</i>	102
<i>Totanus stagnatilis</i>	113	————— <i>simillimus</i>	493
————— <i>totanus eurhinus</i>	113	————— <i>unicolor</i>	467
<i>Tragopan melanocephalus</i>	482	————— <i>bonapartei</i>	194, 371
————— <i>satyra</i>	376	<i>Tylococcus formicarii</i>	453
<i>Tragulus ravus lampensis</i>	48	————— <i>simplicior</i>	453
————— <i>mergatus</i>	48	<i>Tylonycteris meyeri</i>	44
<i>Trappa bispinosa</i>	262	————— <i>pachypus</i>	44
<i>Tribulus terrestris</i>	262	<i>Typha latifolia</i>	262
<i>Tribura major</i>	189, 190, 191, 193	<i>Tyto alba alba</i>	107
<i>Trichoclada opsinaria</i>	139	————— <i>erlangeri</i>	107
<i>Trichoplites cuprearia</i>	310	<i>Udaspes folus</i>	438
————— <i>latifasciaria</i>	310	————— <i>stellatta</i>	438
<i>Tringa hypoleuca</i>	187, 188, 484	<i>Uliocnemis albiradiata</i>	132
————— <i>ochropus</i>	192, 484	<i>Upupa epops epops</i>	106, 476
————— <i>totanus</i>	484	————— <i>ceylonensis</i>	298
————— <i>eurhinus</i>	192	————— <i>e-nigripennis</i>	373
<i>Triphosa consona</i> n. sp.	313	<i>Urocissa flavirostris cucullata</i>	224, 459
<i>Trochalopteron cachinans cachinans</i>	301	————— <i>melanocephala occipitalis</i>	225, 459
————— <i>erythrocephalum erythrocephalum</i>	370, 461	<i>Uroloncha malabarica</i>	269, 284, 372
————— <i>lineatum griseicentior</i>	462	————— <i>p. punctulata</i>	372, 471
————— <i>lineatum</i>	195, 370, 462	<i>Vandeleuria dumeticola</i>	400
————— <i>variegatum simile</i>	195	————— <i>oleracea</i>	382
————— <i>variegatum</i>	462	<i>Vanellus vanellus</i>	113
<i>Troglodytes troglodytes</i>	105	<i>Vesperugo pachypus</i>	44
————— <i>hyrcanus</i>	105	<i>Viverra zibetha</i>	8, 392
————— <i>neglectus</i>	188, 194, 463	<i>Viverricula malaccensis</i>	9, 392
————— <i>pallidus</i>	105	<i>Vulpes bengalensis</i>	381, 403
<i>Troides ferrari</i>	248	<i>Vulpes leucopus</i>	405
<i>Turbinicola saxeae</i>	449	<i>Walkeria ovilla</i>	451
<i>Tupaia belangeri assamensis</i>	390	<i>Watsoniella swinhoei</i>	437
————— <i>brunetta</i>	45	<i>Xantholæma hæmacephala indica</i>	281
————— <i>versuræ</i>	390	<i>Xanthorhœ placida</i>	144
————— <i>clarissa</i>	45	<i>Xenococcus annandalei</i>	454
<i>Turdoides griseus griseus</i>	292	<i>Xenoclystia nigroviridata</i>	145
————— <i>terricolor terricolor</i>	263	————— <i>phæoloma</i>	146
<i>Turdus albocinctus</i>	226, 493	————— <i>unijuga</i>	145
————— <i>atroregularis</i>	269, 467	<i>Zizyphus jujuba</i>	262
————— <i>boulboul</i>	371	————— <i>nummularia</i>	262
————— <i>bourdilloni</i>	493	<i>Zographetus orgygia andamana</i>	442
————— <i>castaneus</i>	188, 226	————— <i>orgygia</i>	442
————— <i>maximus</i>	493	————— <i>satwa</i>	442
		<i>Zoothera monticola</i>	225, 226
		<i>Zosterops palpebrosa</i>	373
		————— <i>elwesi</i>	273, 475
		————— <i>palpebrosa</i>	300

GEOMETRIDÆ FROM UPPER BURMA

Explanation of Plate I

To face Plate I on page 129 of this Volume.

Fig.	1.	<i>Gelasma chromatocrossa</i> ,	Prout, sp. n.
„	2.	<i>Nothomiza aegriviridis</i> ,	„ „
„	3.	<i>Comibæna swanni</i> ,	„ „
„	4.	<i>Pæcilasthena burmensis</i> ,	„ „
„	5.	<i>Hydrelia opedogramma</i> ,	„ „
„	6.	<i>Thera cyphoschema</i> ,	„ „
„	7.	<i>Organopoda brevipalpis</i> ,	„ „
„	8.	<i>Sterrha lamprotis</i> ,	„ „
„	9.	„ <i>castelli</i> ,	„ „
„	10.	<i>Heterostegania thamina</i> ,	„ „
„	11.	<i>Triphosa consona</i> ,	„ „
„	12.	<i>Terpna vigil</i> ,	„ „
„	13.	<i>Opisthograptis swanni</i> ,	„ „
„	14.	<i>Hypochrosis eurynota</i> ,	„ „
„	15.	<i>Lampropteryx opistholasia</i> ,	„ „
„	16.	<i>Swannia marmarea</i> ,	„ „
„	17.	<i>Bapta aluta</i> ,	„ „
„	18.	<i>Lomographa tenebrimedia</i> ,	„ „
„	19.	<i>Cænotephria mononyssa</i> ,	„ „
„	20.	<i>Chiasmia levata</i> ,	„ „
„	21.	<i>Tasta argozona</i> ,	„ „
„	22.	<i>Nothomiza ithyterpa</i> ,	„ „
„	23.	<i>Hydrelia enisaria</i> ,	„ „
„	24.	<i>Rhinoprora xanthocomes</i> ,	„ „

NOTES ON THE BIRDS OF THE SIKKIM HIMALAYAS

Additions and Corrections

Volume XXIX

P. 504, line 20 from the bottom. Additional matter having been inserted in my MS., this sentence should read 'Since the first volume of Oates's *Fauna* was published in 1889, our knowledge has advanced from several bases, excepting in regard to Pterylography, i.e., the study of the Pterylosis (the distribution of the feather-tracts (*pterylæ*), in opposition to the featherless interspaces (*apteria*), which is of importance in nestling birds as an aid to classification, a branch of Ornithology.'

P. 507, line 10 from the bottom. *Delete* 'which' *after* 'Suthoras'; and *delete* the 'semicolon' *after* 'and.' My MS reads: 'Crow-Tits and Suthoras are just as dependent on reed and bamboo-growth, — as Nutcrackers, Crossbills, and are generally considered as occurring exclusively in the pine forests.'

Delete the brackets enclosing the name of the describer of the bird in Nos. 1, 2, 3, 4, 8, 11, 14, 18, 22, 24, 26, 27, 39, 41, 46, 47, 48, 49, 51, 53, 55, 56, 57, 59, 60, 62, 67, 68, 70, 71, 72, 74, 75, 76, 77, 81, 83, 85, 86, 89, 91, 92, 93, 96, 131, 133, 143, as the generic name employed is that in his original description.

P. 515, line 5 from the top. *Delete* 'A.' and read: 'Chelura'. Transfer 'A' to p. 517, line 4 from the top, and *read* 'A flock.'

No. 11. *Nucifraga hemispilla* is possibly best treated as a distinct species with its own group of forms; especially as 'multipunctata' occupies an intervening area between 'caryocatactes' and 'hemispila', and is a good species.

No. 15. *Read* 'PARUS MONTICOLUS LEPCHARUM, Meinertzhagen.
Bull. B.O.C. vol. xlv, p. 96, April 1926.'

No. 16. *Read* 'AEGITHALISCUS CONCINNA RUBRICAPILLUS, Ticehurst.
Bull. B.O.C. vol. xlv, p. 22, October 1925.'

No. 17. In the original description IOUSCHISTOS.

No. 24. *Read* 'AEMODIUS'.

P. 726, line 20 from the bottom. *Delete* '14.6.'

P. 727, line 6 from the bottom. *For* 'Second Reference "3,600 feet"', *read* "3,800 feet."

No. 32. CÆRULATUS.

No. 39. *Substitute* 'Hodgson' *for* 'Oates.'

No. 41. *Substitute* 'Blyth' *for* 'Hodgson.'

No. 50. XIPHIRHYNCHUS should be valid yet not confused with the genus *Xiphorhynchus*.

No. 67. *Substitute* 'Blyth' *for* 'Hodgson.'

P. 735, line 1, SIBIINÆ.

No. 72. The Hoary Bar-wing. *Substitute* 'Gould' *for* 'Hodgson.'

No. 82. LETHOTRIX LUTEA CALIPYGA.

P. 737, line 1 from the bottom, 'Calipyga.'

P. 739, line 16 from the bottom. *Insert after* 'under', 'each'; line 7, 'Flower-pecker.'

- No. 93. In original description. *IGNOTINCA*.
- No. 104. *SITTA CASTANEIVENTRIS CINNAMOVENTRIS*, Blyth.
- No. 108. *MACROCERCUS*.
- P. 1009, line 7 from the bottom. *Delete* 'many' and *substitute* 'and'.
- P. 1010, line 21 from the top. *P. rufonuchalis beavani* and *P. d. dichrous*.
- No. 125. *Read* 'PNŒPYGA ALBIVENTER' Hodgson, *J. A. S. B.*, February 1837. See Kinnear, *Bull. B. O. C.*, vol. xlv, p. 9.'
- No. 126. PNŒPYGA.
- No. 127. *Read* 'REGULUS REGULUS SIKKIMENSIS, Meintz. *Bull. B. O. C.* vol. xlvi, p. 97, April 1926.'
- No. 128. *Read* 'CEPHALOPYRUS FLAMMICEPS SATURATUS, Whistler. *Bull. B. O. C.* vol. xlv, p. 15, October 1924.
- No. 133. *ATROGULARIS*.
- No. 137. *AEDON*.
- P. 1021, bottom line. *Delete* 'hills' and *insert*, 'this.'
- P. 1022, line 11 from the top. *Delete* 'Ibis' and *insert*, 'Ibid.'
- P. 1023, line 14 from the bottom. *Delete* 'is' and *insert* 'was'.
- P. 1026, line 15 from the bottom. *Delete* 'steel transparent.'
- P. 1029, line 5 from the top. *After* 'Sikkim' *add*, 'in the National Collection.' Line 13, ♂, April 4, 1912.
- vol. xxx, page 56, line 6 from the top. *Delete* '5,' *Read* '♂ March 30. 1918.'
- P. 57, line 3 from the top: *Caryota*.
- No. 182. Most probably best treated as a good species. In Tonkin, both *indicus* and *tenuirostris* occur at similar levels during the breeding season *indicus* being confined to the heavily forested region, while *tenuirostris* occupies the surrounding tracts of a more open character.
- Page 60, line 5 from the bottom. *Read* '♀ † October 2, 1919.'
- Page 61, line 7 from the top. '† Four'
- No. 203. *Read* 'CYORNIS MELANOLEUCA, Blyth. = *Muscicapa collini*, Roths *Bull. B. O. C.*, vol. xlv, pp. 89, 90.'
- Page 62. For '*C. blythi*', read '*C. melanoleuca*.'
- Page 65, line 30 from the top. *Read* '10,160 feet.'
- Page 66, line 10 from the top. TERPSIPHONE.
- No. 222. *Read* 'This Chat which ascends the hills of the outer ranges in Sikkim, to breed at moderate elevations, is provisionally retained under *indica*; but my two specimens are very diminutive: wing, ♂ 66.5; ♀ 64 mm. It is represented in the National Collection by many other similar specimens from Sikkim.'
- No. 223. *Read* 'The Turkestan Bush-Chat. *SAXICOLA PRZEWALSKII* (Pleske).' I relegate all specimens of this cold weather migrant to the Plains of Assam and the outer hills of Sikkim under *przewalskii*. My specimens are certainly not *stejnegeri*. Both *przewalskii* and *stejnegeri* breed in the mountains of Tonkin with somewhat similar zonal distributions, or at all events of a none too excessive variation, *stejnegeri* being confined mainly to lower limits. For this reason I retain *przewalskii* as a good species. I regard the breeding Chat from Yunnan as nearer to *stejnegeri* than *indica*, wherealso *przewalskii* is certain to breed as the Tonkin specimen with eggs which I obtained was collected on the Yunnan frontier.
- No. 229. *IMMACULATUS*.
- No. 238. *Güldenstadt*.

Page 355, line 22 from the bottom. To '*Oriolus trailii*' also add 'the male of *Irena puella*.'

Nos. 281, 282. LAISCOPUS.

No. 284. *Substitute* 'Moore' for 'Hodgson.'

No. 290. I follow Whistler and treat this Grosbeak as a good species. *Ibid.* vol. xxx, pp. 701-2.

No. 297. LOXIA.

Page 372, line 24 from the bottom. *Delete* 'more' and *insert* 'were.'

Page 377, line 26 from the top. *alboides*.

No. 411. FUCIPHAGA.

No. 421. SPARVERIOIDES.

No. 459. GYPÆTUS.

No. 461. HIERAÆTUS.

No. 464. SPIZÆTUS.

Page 884, line 25 from the bottom. *Ilerda*.

Page 887, line 30 from the bottom. 'diminishing'.

Line 26, 'depredations'.

Line 17, 'smashed.'

Page 888, line 5 from the top. 'rocky.'

Line 2 from the bottom: 'has.'

No. 520. *Read* 'TETRAOGALLUS TIBETANUS AQUILONIFER, Meintz. *Bull. B. O. C.*, vol. xlvi, pp. 99, 100, April 1926.'

H. S.

THE
JOURNAL

OF THE

BOMBAY NATURAL HISTORY SOCIETY

INDEX AND TITLE PAGE

VOL. XXXI

PARTS 3 & 4

MADRAS

PRINTED AT THE DIOCESAN PRESS

1927

For convenience in binding it has been found advisable to issue the Index in two parts. The present issue includes the Index to Parts 3 and 4, Vol. XXXI, pp. 533-1042. The Index to Parts 1 and 2 is issued separately.

INSTRUCTIONS TO BINDER.

The contents of these two parts should be arranged in the following order when they are being bound :—

Title page	Frontispiece.
Contents of Parts 3 and 4, Vol. XXXI.	}		To follow the frontispiece in this order.
List of Contributors	
List of Plates	
Index to Illustrations	
Errata	To go at the end of the two numbers.
Index of species	

THE
JOURNAL
OF THE
BOMBAY NATURAL HISTORY SOCIETY

EDITED BY

SIR R. A. SPENCE, KT., F.Z.S., S. H. PRATER, C.M.Z.S. and SALIM A. ALI

VOL. XXXI

Parts 3 & 4

Containing 2 Coloured Plates, 21 Black and White Plates,
2 Maps and 16 Text-figures

Dates of Publication

Part III. (Pages 533 to 832) ... 1st November, 1926.
„ IV. („ 833 to 1042) ... 20th February, 1927.

LONDON AGENTS

DULAU & Co., Ltd., 32, Old Bond Street, London, W. I.

PRINTED AT THE DIOCESAN PRESS, MADRAS.

CONTENTS OF VOLUME XXXI.

No. 3

	PAGE
THE GAME BIRDS OF THE INDIAN EMPIRE. (<i>With a coloured plate.</i>) The Purple Coot. (<i>Porphyrio p. poliocephalus.</i>)...	533
REVISION OF THE FLORA OF THE BOMBAY PRESIDENCY.	
Part I. By E. Blatter, S.J., Ph.D. F.Z.S.....	547
SNAKES COLLECTED IN BURMA IN 1925. (<i>With a block.</i>) By Col. F. Wall, C.M.G., K.H.S.....	558
A CONTRIBUTION TO THE ORNITHOLOGY OF DELHI. Part II. By the Late S. Basil-Edwardes, M.B.O.U., R.A.O.U.....	567
NOTES ON SOME NEW AND INTERESTING BUTTERFLIES FROM INDIA AND BURMA. Part II. (<i>With three plates.</i>) By Major-General H. C. Tytler, C.B., C.M.G., C.I.E., D.S.O., F.E.S.....	579
BOMBAY NATURAL HISTORY SOCIETY'S MAMMAL SURVEY OF INDIA, BURMA AND CEYLON. Report Nos. 43, 44 and 45 and Supplement to Report Nos. 23 and 31. By Mrs. Helen M. Lindsay, M.A., B.Sc.....	591
THE IDENTIFICATION OF INDIAN BUTTERFLIES. Part XI. (<i>With a plate.</i>) By Col. W. H. Evans, D.S.O., F.Z.S., F.E.S.....	615
A SUMMARY OF SCIENTIFIC RESEARCH IN THE PERSIAN GULF. By Sir A. T. Wilson, K.C.I.E., C.S.I., C.M.G., D.S.O., etc.....	638
THE COMMON BUTTERFLIES OF THE PLAINS OF INDIA. Part XXXVII. (Including those met with in the Hill Stations of the Bombay Presidency.) By T. R. Bell, C.I.E., I.F.S. (Retd.).....	655
THE BIRDS OF BRITISH BALUCHISTAN. Part I. (<i>With two plates and a map.</i>) By C. B. Ticehurst, M.A., M.R.C.S., M.B.O.U., F.R.G.S.....	687
NOTES ON INDIAN BUTTERFLIES. By Col. W. H. Evans, D.S.O., F.Z.S., F.E.S.....	712
AN ELEPHANT SHOOT ON THE BARAGUR HILLS. Coimbatore District. (<i>With a plate.</i>) By R. C. Morris.....	720
A NEW ICHNEUMON FLY OF THE GENUS <i>Aphanteles</i> from the Death's Head Moth <i>Acherontia styx</i> . (<i>With a plate.</i>) By C. F. W. Muesebeck, with a note on its Life History by C. McCann.....	726

	PAGE
NOTES ON SMALL GAME SHOOTING IN THE KHASI HILLS. By E. McCulloch.....	728
INDIAN DRAGONFLIES. Part XXV. (<i>With one plate and five text-figures.</i>) By Lt.-Col. F. C. Fraser, I.M.S., F.E.S.....	733
LUMINESCENCE IN PLANTS AND ANIMALS. By E. Blatter, S.J., Ph.D., F.Z.S.....	748
THE MASON WASP (<i>Eumenes conica</i>). Part II. By Major R. W. G. Hingston, I.M.S.....	754
DRAGONFLIES—A REVISION OF THE GENUS <i>Zygonyx</i> , Selys. (<i>With a text-figure.</i>) By Lt.-Col. Major F. C. Fraser, I.M.S., F.E.S.....	762
NOTES ON FISHES FROM BOMBAY. By H. W. Fowler.....	770
ON A COLLECTION OF MOTHS OF THE FAMILY <i>Geometridæ</i> from Upper Burma made by Capt. A. E. Swann. Part III. By L. B. Prout, F.E.S.....	780
REVIEWS :—GOLD FISH CULTURE FOR AMATEURS	800
BIG GAME HUNTING IN THE HIMALAYAS AND TIBET.....	800
‘THE JOURNAL OF THE DARJEELING NATURAL HISTORY SOCIETY.’.....	801
THE PALMS OF BRITISH INDIA AND CEYLON.....	801
A HANDBOOK OF THE BIRDS OF EASTERN CHINA	802
EDITORIAL.....	803
MISCELLANEOUS NOTES :—	
I. A Panther Shoot at Sea. (<i>With a photo.</i>) By N. Kirwan.....	808
II. The Breeding Habits of the Panther (<i>Felis pardus</i>). By C. H. Donald.....	809
III. A Tigress with Five Cubs. By R. C. Morris.....	810
IV. Wild Dogs and Further Jungle Tragedies. By R. C. Morris.....	811
V. The Breeding Seasons of the Gooral (<i>Nemorhædus goral</i>) and Himalayan Tahr (<i>Hemitragus jemlaicus</i>). By Capt. E. E. G. L. Searight, R.E.	812
VI. Jackal and Hare. By Salim A. Ali.....	812
VII. How do Wild Dogs kill their Prey. By Salim A. Ali.....	813
VIII. Twin Elephant Calves. (<i>With a photo.</i>) By Major C. H. Stockley.....	813
IX. Notes on the Kashmir Stag (<i>Cervus hanglu</i>). (<i>With a photo.</i>) By Major C. H. Stockley.....	814
X. List of Cetaceans taken in Travancore from 1902–19. By R. Shunkara Narayan Pillay.....	815

	PAGE
XI. List of Birds observed in the Bheling Valley, Tehri Garhwal. By Capt. E. E. G. L. Searight, R.E.....	817
XII. Assamese Names of certain Birds used in Nowgong District. By J. C. Higgins, I.C.S.....	819
XIII. Occurrence of the Indian Courser. (<i>Cursorius coromandelicus</i>) at Panchgani. By C. McCann...	820
XIV. Note on the Breeding of the Genus <i>Caprimulgus</i> (Nightjars) in the Punjab Salt Range. (<i>With a plate.</i>) By H. W. Waite, F.Z.S., M.B.O.U.....	821
XV. Occurrence of the Sheldrake (<i>Tadorna tadorna</i>) and Lapwing (<i>Vanellus vanellus</i>) in Upper Burma. By J. K. Stanford, I.C.S., M.B.O.U.....	823
XVI. The Mating of Crows. By Salim A. Ali.....	823
XVII. A Note on the Nidification of the Western Reef Egret (<i>Leptorodius asha</i>) in Karachi City, Sind. (<i>With a plate.</i>) By K. R. Eates.....	823
XVIII. Breeding of the Great Himalayan Barbet. (<i>Megalæma virens marshallorum</i>) in the Punjab Salt Range. By H. W. Waite, F.Z.S., M.B.O.U...	825
XIX. Extraordinary Tameness of the Garganey Teal (<i>Querquedula querquedula</i>). By Capt. J. Mc. C. Clive.....	826
XX. The Red-legged Falconet (<i>Microhierax eutolmus</i>) Hawking Butterflies. By S. F. Hopwood, I.F.S.	826
XXI. House Gecko (<i>Hemidactylus</i> sp.) shows a Sweet Tooth. By C. McCann.....	827
XXII. Comments on Fr. Leigh's Notes on Snakes. By C. McCann.....	827
XXIII. Parturitions of Electric Rays and a Sea Snake in the Marine Aquarium, Madras. By B. Sundara Raj.....	828
XXIV. Fishing in Lonavla. (<i>With a block.</i>) By F. V. Evans.....	828
XXV. Death from the Sting of a Scorpion. By G. S. Shukla, L.M. & S.....	803
PROCEEDINGS	831

No. 4

THE MOGHUL EMPERORS OF INDIA AS NATURALISTS AND SPORTSMEN. Part I. (<i>With three plates.</i>) By Salim A. Ali.....	833
---	-----

	PAGE
THE BIRDS OF BRITISH BALUCHISTAN. Part II. (<i>With two plates.</i>) By C. B. Ticehurst, M.A., M.R.C.S., M.B.O.U., F.R.G.S.....	862
INDIAN DRAGONFLIES. Part XXVI. (<i>With three text-figures.</i>) By Lt.-Col. F. C. Fraser, I.M.S., F.E.S.....	882
THE MASON WASP (<i>Eumenes conica</i>). Part III. By Major R. W. G. Hingston, I.M.S.....	890
REVISION OF THE FLORA OF THE BOMBAY PRESIDENCY. Part II. By E. Blatter, S.J., Ph.D., F.L.S.....	897
IMPRESSIONS OF PACHMARHI BIRDS. (<i>With four plates.</i>) By Capt. R. S. P. Bates.....	918
ON A COLLECTION OF MOTHS OF THE FAMILY <i>Geometridæ</i> from Upper Burma made by Capt. A. E. Swann. Part IV. By L. B. Prout, F.E.S.....	932
THE COMMON BUTTERFLIES OF THE PLAINS OF INDIA (Including those met with in the Hill Stations of the Bombay Presidency). Part XXXVIII. By T. R. Bell, C.I.E., I.F.S. (Retd.).....	951
NOTES ON THE BIRDS OF KASHMIR. Part I. (<i>With a plate and a map.</i>) By B. B. Osmaston, C.I.E., M.B.O.U.....	975
FURTHER NOTES ON THE BIRDS OF AMBALA DISTRICT. By A. E. Jones.....	1000
A CRITICAL REVIEW OF SIR J. C. BOSE'S 'NERVOUS MECHANISM OF PLANTS.' By R. H. Dastur.....	1009
REVIEWS :—BIRD STUDY IN INDIA.....	1015
IN UNKNOWN ARABIA.....	1015
NOMOGENESIS OF EVOLUTION DETERMINED BY LAW.....	1017
AN INTRODUCTION TO EXPERIMENTAL EMBRYOLOGY.....	1018
A SHIKARI'S POCKET-BOOK.....	1019
EDITORIAL.....	1020
MISCELLANEOUS NOTES :—	
I. A Comment on 'Tigers and Elephants' and Notes on Tigers and Buffaloes. By Maharaj Kumar Victor N. Narayen of Cooch Behar.....	1025
II. Notes on Tiger preferring Carrion to live Bait. By A Beginner.....	1025
III. Black Leopards. By Salim A. Ali.....	1027
IV. Ships' Cats, and an Instance of their Homing Instinct. By J. H. Hughes, M.B.E.....	1027
V. Wild Dogs in Mysore. By E. G. Phythian-Adams, Major I.A. (Retd.).....	1028

	PAGE
VI. Notes on the Long-armed Sheath-tailed Bat (<i>Taphozous longimanus</i> Hardw.). By C. McCann. (<i>With a photograph</i>).....	1030
VII. The Breeding of the Indian Rhinoceros (<i>Rhinoceros unicornis</i>) in Captivity. By Salim A. Ali...	1031
VIII. Extension of Habitat of Hume's Scimitar Babbler. (<i>Pomatorhinus horsfieldi obscurus</i>) to Orissa. By Salim A. Ali.....	1032
IX. Indian Cuckoo Notes—Koel (<i>Eudynamis s. scolopaceus</i>) parasitising nest of Indian Oriole. (<i>Oriolus o. kundoo</i> .) By E. A. D'Abreu, F.Z.S....	1032
X. Occurrence of the Wood Snipe (<i>Gallinago nemoricola</i>) in Burma. By A. Macdonald.....	1033
XI. Arrival of Snipe in Mysore. By G. V. R. Frend.	1033
XII. Migration of Wildfowl. By G. B. Pauri, Private Secretary to H. H. The Maharaja of Dhar.....	1034
XIII. The Breeding of the Indian Lesser-Whitethroat (<i>Sylvia curruca affinis</i>) at Quetta. By Brig.- Genl. R. M. Betham.....	1035
XIV. The Effects of a Recent Cyclonic Storm on Bird- Life in Karachi and its Environs. By K. R. Eates.....	1035
XV. Strange Behaviour of a Truncated Fly. By Major C. W. Sanders.....	1037
XVI. The Pairing of Sea-Snakes. (<i>With a photo.</i>) By Commander A. R. Poyntz.....	1038
XVII. Occurrence of the Worm-like Batrachian (<i>Ichthyophis monochrous</i> Bouleng.) at Khandala, Poona District. By C. McCann.....	1039
XVIII. Viviparity in a Thistle. (<i>With a plate.</i>) By E. Blatter.....	1039
XIX. The Use of Patent Bullets in Shot Guns. By E. L. McCulloch.....	1040

ALPHABETICAL LIST OF CONTRIBUTORS

VOLUME XXXI

Nos. 3 and 4

	PAGE		PAGE
ALI, SALIM A.; Jackal and Hare	812	of the Plains of India including those met with in the Hill Stations of the Bombay Presidency, Part XXXVII	655
—————; How do Wild Dogs kill their Prey	813	—————; Part XXXVIII	951
—————; The Mating of Crows	823	BETHAM, BRIGADIER-GENERAL, R. M.; The Breeding of the Indian Lesser Whitethroat (<i>Sylvia curruca affinis</i>) at Quetta	1035
—————; The Moghul Emperors of India as Naturalists and Sportsmen. Part I (with three plates)	833	BLATTER, E., S. J., Ph. D., F.L.S.; Revision of the Flora of the Bombay Presidency, Part I	547
—————; Black Leopards	1027	—————; Part II	897
—————; The Breeding of the Indian Rhinoceros (<i>Rhinoceros unicornis</i>) in Captivity.	1031	—————; Luminescence in Plants and Animals	748
—————; Extension of Habitat of Hume's Scimitar Babbler (<i>Pomatorhinus horsfieldi obscurus</i>) to Orissa	1032	—————; Viviparity in a Thistle (with a plate)	1039
BAKER, E. C. STUART, J. P., F. Z. S., Th. S., M. B. O. U., C. F. A. O. U.; The Game Birds of the Indian Empire, Vol. V. The Waders and other Semi-sporting Birds, Part II (with a coloured plate) of <i>Porphyrio poliocephalus poliocephalus</i> . The Indian Purple Moorhen, The Kora or Water-Cock, The Indian Moorhen, The Coot	533	CLIVE, CAPT. J. McC.; Extraordinary Tameness of the Garganey Teal (<i>Querquedula querquedula</i>)	826
BASIL-EDWARDES, The late S., M. B. O. U., R. A. O. U.; A Contribution to the Ornithology of Delhi, Part II	567	D'ABREU, E. A., F. Z. S.; Indian Cuckoo Notes—Koel (<i>Eudynamis s. scolopaceus</i>) parasitizing nest of Indian Oriole (<i>Oriolus o. kundoo</i>)	1032
BATES, CAPT. R. S. P.; Impressions of Pachmarhi Birds (with 4 plates)	918	DASTUR, R. H.; A Critical Review of Sir J. C. Bose's 'Nervous Mechanism of Plants' (communicated by E. Blatter)	1009
BEGINNER, A.; Notes on Tiger preferring Carrion to Live Bait	1025	DONALD, C. H.; The Breeding Habits of the Panther (<i>Felis pardus</i>)	809
BELL, T. R., C.I.E., I.F.S., (Retd.) The Common Butterflies		EATES, K. R.; A Note on the Nidification of the Western Reef Egret (<i>Leptorodius asha</i>) in Karachi City, Sind (with a plate)	823
		—————; The Effects of a recent Cyclonic Storm on Bird-Life in Karachi and its Environs	1035

	PAGE		PAGE
EDITORIAL;—Game Preservation	803	India, Burma and Ceylon,	
—————; Mr. B. C. Ellison ...	804	Report No. 43, Nelliampathy	
—————; Mr. Prater in		Plateau and Palni Hills (<i>with</i>	
America	805	<i>Field-notes by the Collectors</i>).	591
—————; Shikar Stories	807	MCCANN, C.; Note on the Life	
—————; Sir Sasson David,		History of an Ichneumon Fly	
Bart., K. C. S. I.	807	of the Genus <i>Aphanteles</i> (<i>with</i>	
EVANS, F. V.; Fishing in Lon-		<i>a plate</i>)	726
avla (<i>with a text-figure</i>)	828	—————; The Occurrence of	
EVANS, COL. W. H., D.S.O.,		the Indian Courser (<i>Cursorius</i>	
F.Z.S., F.E.S.; The Identifi-		<i>coromandelicus</i>) at Panchgani.	820
cation of Indian Butterflies,		—————; House Gecko	
Part XI (<i>with a plate</i>)	615	(<i>Hemidactylus</i> sp.) shows a	
—————; Notes on		Sweet Tooth	827
Indian Butterflies	712	—————; Comments on Fr.	
FRASER, LT.-COL. F. C., I.M.S.,		Leigh's Notes on Snakes	827
F.E.S.; Indian Dragonflies,		—————; Notes on the	
Part XXV (<i>with 1 plate and 5</i>		Long-armed Sheath-tailed Bat	
<i>text-figures</i>)	733	(<i>Taphozous longimanus</i> ,	
—————; Part XXVI (<i>with 3 text-</i>		Hardw.) (<i>with a photo</i>)	1030
<i>figures</i>)	882	—————; Occurrence of the	
—————; A Revision of the genus		Worm-like Batrachian (<i>Ichthyo-</i>	
<i>Zygonyx</i> , Selys. (<i>with a text-</i>		<i>phis monochrous</i> , Bouleng.) at	
<i>figure</i>)	762	Khandala, Poona District	1039
FOWLER, H. W.; Notes on		MACDONALD, A.; Occurrence of	
Fishes from Bombay	770	the Wood-Snipe (<i>Gallinago</i>	
FREND, G. V. R.; Arrival of		<i>nemoricola</i>) in Burma	1033
Snipe in Mysore	1033	MCCULLOCH, E.; Notes on Small	
HIGGINS, J. C., I.C.S.; Assamese		Game Shooting in the Khasi	
names of certain Birds used in		Hills	728
Nowgong District	819	MCCOLLOCH, E. L.; The Use	
HINGSTON, MAJOR R. W. G.;		of Patent Bullets in Shot	
The Mason Wasp (<i>Eumenes</i>		Guns	1040
<i>conica</i>) Part II	754	MORRIS, R. C.; An Elephant	
—————; Part III	890	Shoot on the Baragur Hills,	
HOPWOOD, S. F., I.F.S.; The		(Coimbatore District) (<i>with a</i>	
Red-legged Falconet (<i>Mic-</i>		<i>plate</i>)	720
<i>rorhierax leutolmus</i>) Hawking		—————; A Tigress with	
Butterflies	826	five Cubs	810
HUGHES, COMMANDER JOHN H.,		—————; Wild Dogs and Fur-	
M.B.E.; Ships' Cats, and an		ther Jungle Tragedies	811
instance of their Homing		MUESEBECK, C. F. W.; A new	
Instinct	1027	Ichneumon Fly of the Genus	
JONES, A. E.; Further Notes on		<i>Aphanteles</i> from the Death's	
the Birds of the Ambala		Head Moth (<i>Acherontia styx</i>)	
District	1000	with a Note on its Life History	
KIRWAN, N.; A Panther Shoot at		by C. McCann (<i>with a plate</i>)	726
Sea (<i>with a photo</i>)	808	NARAYEN, VICTOR; A Com-	
LINDSAY, MRS. HELEN M., M.A.,		ment on 'Tigers and Ele-	
B.Sc.; Bombay Natural History		phants' and Notes on 'Tigers	
Society's Mammal Survey of		and Buffaloes	1025

	PAGE		PAGE
OSMASTON, B. B., M.B.O.U. ; Notes on the Birds of Kashmir, Part I (<i>with a plate and a map</i>)	975	SEARIGHT, CAPT. E. E. G. L., F.R. ; The Breeding-seasons of the Gooral (<i>Nemorhædus gooral</i>) and Himalayan Tahr (<i>Hemitra- gus jemlaicus</i>)	812
PAURI, G. B. ; Migration of Wild Fowl	1034	— ; A List of Birds observed in the Bheling Valley, Tehri Garhwal, April 1, 1926 to May 25, 1926, from 2,000 feet to 13,000 feet ...	817
PHYTHIAN-ADAMS, MAJOR E. G., I. A. (Retd.) ; Wild Dogs in Mysore	1028	SHUKLA, G. S., L.M. & S. ; Death from the Sting of a Scorpion...	830
PILLAY, R. ; Shunkara Narayan, C. M. Z. S. ; List of Cetaceans taken in Travancore from 1902 to 1925	815	STANFORD, J. K., M. B. O. U. ; Occurrence of the Sheldrake (<i>Tadorna tadorna</i>) and Lap- wing (<i>Vanellus vanellus</i>) in Upper Burma	823
POYNTZ, A. R. ; The Pairing of Sea Snakes (<i>with a photo</i>) ...	1038	STOCKLEY, MAJOR C. H. ; Twin Elephant Calves (<i>with a photo</i>)	813
PROCEEDINGS	831	— ; Notes on the Kash- mir Stag (<i>Cervus hanglu</i>) (<i>with a photo</i>)	814
PROUT, L. B., F.R.S. ; On a Col- lection of Moths of the Family <i>Geometridæ</i> from Upper Burma made by Captain A. E. Swann, Part III	780	TICEHURST, C. B., M. A., M. R. C. S., M.B.O.U., F.R.G.S. ; The Bird of British Baluchistan, Part I (<i>with two plates and a map</i>)	687
— ; Part IV	932	— — — — — Part II (<i>with two plates</i>)	862
RAJ, B. SUNDARA ; Parturitions of Electric Rays and a Sea Snake in the Marine Aqua- rium, Madras	828	TYTLER, MAJOR-GENERAL H. C., C. B., C.M.G., C.I.E., D.S.O., F.E.S. ; Notes on some new and interesting Butterflies from India and Burma, Part II (<i>with one coloured plate and two black and white plates</i>) ...	579
REVIEWS:—Gold Fish Culture for Amateurs	800	WAITE, H. W. ; Note on the Breeding of the Genus <i>Capri- mulgus</i> (Nightjars) in the Punjab Salt Range	821
— ; Big Game Hunting in the Himalayas and Tibet	800	— ; Breeding of the Great Himalayan Barbet (<i>Me- galæma virens marshallorum</i>) in the Punjab Salt-range ...	825
— ; <i>The Journal of the Darjeeling Natural History Society</i> , Vol. I, No. 1	801	WALL, COL. F., C. M. G., K. H. S. ; Snakes collected in Burma in 1925 (<i>with a block</i>)	558
— ; The Palms of British India and Ceylon	801	WILSON, SIR ARNOLD T., K.C. I.E., C.S.I., C.M.G., D.S.O., etc. ; A Summary of Scientific Research in the Persian Gulf...	638
— ; A Handbook of the Birds of Eastern China	802		
— ; Bird Study in India .	1015		
— ; In Unknown Arabia	1015		
— ; Nomogenesis or Evo- lution determined by Law	1017		
— ; An Introduction to Experimental Em- bryology	1018		
— ; A Shikari's Pocket Book	1019		
SANDERS, MAJOR C. W. ; Strange Behaviour of a Truncated Fly...	1037		

LIST OF PLATES

VOLUME XXXI

No. 3

	PAGE
The Game Birds of the Indian Empire—	
Plate—The Purple Coot (<i>Porphyrio poliocephalus poliocephalus</i>) ...	533
New and Interesting Butterflies from India and Burma—	
Plate III—Fig. 1. <i>Liminitis trivena pallida</i> ♂ ...	581
Fig. 2. „ „ <i>trivena</i> ♂ ...	581
Fig. 3. „ „ <i>ligyes</i> ♂ ...	581
Fig. 4. „ <i>lepichini hydaspes</i> ♂ ...	581
Fig. 5. <i>Maniola pulchra astorica</i> ♀ ...	581
Fig. 6. <i>Liminitis lepichini gilgitica</i> ♂ ...	581
Fig. 7. <i>Satarupa splendens</i> ♂ ...	581
Fig. 8. „ <i>zulla</i> ♂ ...	581
Fig. 9. <i>Aemona lena haynei</i> ♂ ...	581
Plate II—Fig. 1. <i>Neptis mackwoodi</i> ♂ ...	583
Fig. 2. „ <i>suffusa</i> ♂ ...	583
Fig. 3. <i>Theda doni</i> ♂ ...	583
Fig. 4. <i>Apatura cooperi</i> ♂ ...	583
Fig. 5. <i>Neptis zaida bhutanica</i> ♂ ...	583
Fig. 6. „ „ <i>manipuriensis</i> ♂ ...	583
Fig. 7. „ „ <i>drummondi</i> ♂ ...	583
Fig. 8. <i>Virachola smilis smilis</i> ♂ ...	583
Fig. 9. <i>Apatura ulupi kalaaurica</i> ♂ ...	583
Plate V—Fig. 1. <i>Dophla cooperi</i> ♂ ...	586
Fig. 2. <i>Charaxes durnfordi merguia</i> ♂ ...	586
Fig. 3. <i>Parnassius acco pundata</i> ♀ ...	586
Fig. 4. <i>Oeneis garhwalia</i> ♀ ...	586
The Identification of Indian Butterflies—	
Plate XXXII HESPERIIDÆ <i>Pamphilinæ</i> (Part) ...	615
The Birds of British Baluchistan—	
1 Map of Baluchistan ...	687
2 (a) Hills from the Quetta Plain with Orchards ...	696
(b) Low bare hills, Sind—Las Belas Frontier ...	696
3 (a) Khirthar Range, Sind, Baluchi Frontier ...	697
(b) A 'Tangi' in the Juniper Forest dotted with wild Rose, etc., ...	697
An Elephant Shoot on the Baragur Hills—	
1 (a) Mr. R. C. Morris's Tusker ...	723
(b) The Trophies ...	723
A New Ichneumon Fly of the Genus <i>Apanteles</i> from the Death's Headmoth (<i>Acherontia styx</i>)—	
1 (a) Caterpillar with Ichneumon Cocoon on underside ...	726
(b) Lateral view of a. ...	726

	PAGE
Indian Dragonflies—	
Dorso-lateral view of <i>Macrogomphus annulatus</i> ♂	736
" " <i>seductus</i> ♂	736
" " <i>montanus</i> ♂	736
" <i>Perissogomphus stevensi</i> ♂	736
" <i>Acrogomphus fraseri</i> ♂	736
" <i>Leptogomphus gestroi</i> ♂	736
" " <i>inclitus</i> ♂	736
Note on the Breeding of the Genus <i>Caprimulgus</i> (Nightjars) in the Punjab Salt Range.	
(a) Typical Breeding Ground of <i>Caprimulgus monticolus</i> and <i>C. europæus uniwini</i> . Sethi, Jhelum District, Punjab Salt Range (altitude 2,600 feet)	821
(b) <i>Caprimulgus europæus uniwini</i> , sitting on eggs Sethi, Jhelum District, Punjab Salt Range (altitude 2,600 feet)	821
Note on the Nidification of the Western Reef Egret (<i>Leptorodius asha</i>) in Karachi City, Sind—	
(a) Nest of <i>Leptorodius asha</i> depicting leafy type with leaves still fairly fresh	823
(b) The same, showing leaf curtain gradually falling off owing to leaves withering and dropping	824
No. 4	
Mogul Emperors of India as Naturalists and Sportsmen—	
Plate 1. The Emperor Jehangir shoots a Large Lion	833
2. The Close of a Qamargah Hunt	842
3. A Lion Hunt with Buffaloes	843
Birds of British Baluchistan—	
Plate I.—(a) In the Juniper Forest, Ziarat	868
(b) On the edge of the Juniper Forest	868
Plate II.—(a) Tangi in Ziarat Forest	869
(b) Khirthar Range, Sind-Baluchi Frontier	869
Pachmarhi Birds—	
1. Indian Purple Sunbird	920
2. Northern Green Barbet	921
3 (a) Common Indian House-Crow	922
(b) Smaller White Scavenger Vulture	922
4 (a) Common Hawk-Cuckoo	923
(b) Indian Magpie Robin	923
Birds of Kashmir—	
Map showing Dominions of the Maharaja of Kashmir	975
1 (a) On the Dal Lake	990
(b) In Camp—Sind Valley	990
Viviparity in a Thistle	1039

INDEX TO ILLUSTRATIONS

VOLUME XXXI

	PAGE		PAGE
<i>Acherontia styx</i> , Pl.	... 726	<i>Leptogomphus gestroi</i> , <i>inclitus</i> , Pl.	
<i>Acrogomphus fraseri</i> , fig. 7,		fig. 7. Dorso-lateral view of male	736
Genitalia	735	————— <i>lansbergei</i> , figs.	
Fig. 9, anal appendage	735	3 and 4, anal appendage	886
Pl. fig. 5. Dorso-lateral view of		<i>Liminitis lepichini gilgitica</i> ,	
male	736	male, Pl. fig. 6	581
Wings, fig. 3	739	————— <i>lepichini hydaspes</i> ,	
<i>Æmona lena haynei</i> , male, Pl.		male Pl. fig. 4	581
fig. 9	581	————— <i>trivena ligyes</i> , male,	
<i>Anormogomphus heteropterus</i> ,		Pl. fig. 3	581
fig. 12, anal appendage.		————— <i>pallida</i> , male,	
Dorso-lateral view	735	Pl. fig. 1	581
—————, Wings of male, fig. 5 <i>b</i> .	744	————— <i>trivena</i> , male,	
————— <i>kiritschenkoi</i> ,		Pl. fig. 2	581
fig. 11, anal appendage. Dorso-		<i>Macrogomphus annulatus</i> , figs.	
lateral view	735	1 and 2, anal appendage	735
—————Wings, fig. 5 <i>a</i> ...	744	—————fig.	
<i>Apatura cooperi</i> , male, Pl. fig. 4...	583	5, Genitalia	735
————— <i>ulupi kalaurica</i> , male,		—————Pl. fig. 1. Dorso-lateral	
Pl. fig. 9	583	view of male	736
<i>Barilius bola</i> , fig.	828	————— <i>montanus</i> , Pl.	
<i>Caprimulgus europæus unwini</i> ,		fig. 3. Dorso-lateral view of male	736
Photo (bird sitting)	820	————— <i>seductus</i> , fig.	
<i>Cervus hanglu</i> , Photo	815	wings of male	733
<i>Charaxes durnfordi merguia</i> ,		Figs. 3 and 4, anal appendage	735
male, Pl. fig. 2	586	—————fig. 6,	
<i>Copsychus saularis saularis</i> , Pl.		Genitalia	735
fig. 2	923	—————Pl. fig. 2. Dorso-lateral	
<i>Corvus splendens splendens</i> , Pl.		view of male	736
fig. 1	922	<i>Maniola pulchra astorica</i> , female,	
<i>Dophla cooperi</i> , male, Pl. fig. 1 ...	586	Pl. fig. 5	581
<i>Elephas indicus</i> , Pl. (Tusks) ...	723	<i>Naia hannah</i> , male sexual organ, fig	565
—————Photo of twin-calves	814	<i>Neophron percnopterus gingini-</i>	
<i>Felis pardus</i> , fig.	808	<i>anus</i> , Pl. fig. 2	922
<i>Hesperiidæ</i> , spp. Pl.	615	<i>Neptis mackwoodi</i> , male, Pl. fig. 1	583
<i>Hierococyx varius</i> , Pl. fig. 1 ...	923	————— <i>suffusa</i> , male, Pl. fig. 2 ...	583
<i>Lepterodius asha</i> , Pl. nests ...	824	————— <i>zaida bhutanica</i> , male,	
<i>Leptocoma asiatica asiatica</i> , Pl.	920	Pl. fig. 5	583
<i>Leptogomphus gestroi</i> , Pl. fig. 6.		————— <i>drummondi</i> , male	
Dorso-lateral view of male... ..	736	Pl. fig. 7	583
—————, wings fig.	882	————— <i>manipurensis</i> , male,	
—————Figs. 1 and 2, anal appendage	886	Pl. fig. 6	583

	PAGE		PAGE
<i>Oeneis garhwalica</i> , female, Pl.		<i>Satarupa splendens</i> , male Pl. fig. 7	581
fig. 4	586	—————, male Pl. fig. 8	581
<i>Pamphilinae</i> , spp. Pl.	615	<i>Sieboldius nigricolor</i> , figs. 5 and 6,	
<i>Parnassius acco punctata</i> , female,		anal appendage	886
Pl. fig. 3	586	—————Wings	887
<i>Perissogomphus stevensi</i> , fig. 8.		<i>Taphozous longimanus</i> , fig. ...	1030
Genitalia, fig. 10, anal appen-		<i>Theda doni</i> , male, Pl. fig. 3 ...	583
dage	735	<i>Virachola smilis smilis</i> , male,	
—————, Pl. fig.		Pl. fig. 8	583
4. Dorso-lateral view of male.	786	<i>Zygonyx iris malabarica</i> ,	
Wings, fig. 4	742	fig. Larva	768
<i>Porphyrio p. poliocephalus</i> ,			
Pl. col.	533		

ERRATA

Vol. XXXI, Nos. 3 and 4

Page 585, line 14 for *Lycæne* read *Lycæna*.

„ 688 „ 9 from bottom for *phillipinus* read *philippinus*.

„ 688 „ 11 „ „ „ *Rhiphidura* read *Rhipidura*.

„ 689 „ 7 for *craytal* read *raytal*. -

„ 692 „ 44 for *Dendroocitta* read *Dendrocitta*.

„ 696 „ 37 „ *Acrocephalcus* read *Acrocephalus*.

Plate opp. p. 696, fig. 1 for *Kippolais* read *Hippolais*.

Page 698 line 1 for *langiuda* read *languida*.

„ 700 „ 47 „ *ceottoides* read *cettoides*.

„ 819 „ 44 for *Calophaps* read *Chalcophaps*.

„ 819 „ 57 „ *Gennæes* read *Gennæus*.

„ 870 „ 9 from bottom for *Barkhaw* read *Barkhan* and for *Fort Sandeman District* read *Loralai District*.

„ 877 „ 44 for *Lynx* read *lynx*.

„ 881 „ 23 „ *Toccacua* read *Taccocua*.

„ 988 „ 31 „ *Lanuis* read *Lanius*.

„ 1001 „ 8 from bottom for *Hororins* read *Horornis*.

„ 1006 „ 20 for *Hierautes* read *Hieraetus*.

„ 1029 „ 20 „ *mature* read *immature*.

INDEX OF SPECIES

	PAGE		PAGE
<i>Abraxas aphorista</i> n. sp.	947	<i>Agrobates galactodes familiaris</i>	696
——— <i>dichostata</i>	950	<i>Alæmon alaudipes cinerea</i>	873
——— <i>illuminata</i>	948	<i>Alauda arvensis cinerascens</i>	1004
——— <i>martaria</i>	948	——— <i>intermedia</i>	874
——— <i>metabasis</i>	949	——— <i>schach</i>	874
——— <i>metamorpha sinicaria</i>	948	——— <i>cinerea</i>	874
——— <i>picaria</i>	949	——— <i>dulcivox</i>	874
——— <i>pusilla</i>	948	——— <i>gulgula gulgula</i>	1004
——— <i>suffusa</i>	948	——— <i>inconspicua</i>	874
<i>Acanthion leucurus</i>	596, 605	<i>Alcedo atthis bengalensis</i>	568, 818
<i>Acanthis cannabina fringillirostris</i>	864	——— <i>pallasii</i>	568, 879, 1005
——— <i>flavirostris ladacensis</i>	996	<i>Alseonax ruficaudus</i>	988
<i>Acanthopneuste magnirostris</i>	991	<i>Alsodeia zeylanica</i>	909
——— <i>occipitalis occipitalis</i> 992, 1001		<i>Amandava amandava amandava</i>	862
<i>Acasis viretata</i>	782	<i>Amathusia phidippus binghami</i>	579
<i>Acerbas albimedia</i>	615	——— <i>fredericki</i>	579
——— <i>anthea</i>	615	<i>Amauornis akool</i>	1007
——— <i>duris duris</i>	615	——— <i>fuscus bakeri</i>	1007
——— <i>martini</i>	615	——— <i>phoenicura</i>	1007
——— <i>nitidifasciata</i>	615	——— <i>chinensis</i>	820
<i>Acherontia styx</i>	726	<i>Amblonyx cinerea</i>	594
<i>Acidalia ærata</i>	787	<i>Amblycephalus macularius</i>	566
——— <i>tephrosaria</i>	792	<i>Amblypodia camdeo sebonga</i> sub. sp. nov. 588	
<i>Acridotheres tristis tristis</i> 704, 817, 819, 993		——— <i>maymyoia</i> n. sp.	588
<i>Acrocephalus agricola agricola</i> 697, 1001		——— <i>suffusa atarana</i> sub. sp. nov. 588	
——— <i>arundinaceus zarudnyi</i> 696		<i>Ammomanes deserti iranica</i>	876
——— <i>concineus concineus</i>	989	——— <i>phoenicuroides</i>	876
——— <i>dumetorum</i>	696, 697	——— <i>phoenicura zarudnyi</i>	876
——— <i>stentoreus brunnescens</i> 696, 989, 1001		<i>Anas platyrhyncha platyrhyncha</i> 577, 820	
——— <i>streperus</i>	697	——— <i>pœcilorhyncha pœcilorhyncha</i> 577, 820	
<i>Acrogomphus fraseri</i>	739, 740	<i>Anamirta cocculus</i>	550
<i>Actinor radians</i>	623	<i>Anastomus oscitans</i>	819, 820, 1007
<i>Adelura cœruleocephala</i>	984	<i>Anaxa kaluga</i>	797
<i>Adolias intermedia merguia</i>	580	<i>Anhinga melanogaster</i>	576, 820
<i>Ægithaliscus concinna iredalei</i> ... 663, 817, 978		<i>Anonychia diversilinea</i>	790
——— <i>leucogenys</i>	693	——— <i>grisea</i>	790
——— <i>niveogularis</i>	978	——— <i>latifasciaria</i>	790
<i>Ægithina nigrolutea</i>	1001	<i>Anormogomphus heteropterus</i> 744, 745	
——— <i>tiphia tiphia</i>	1001	——— <i>kiritschenkoi</i> 744, 746	
<i>Æthopyga siparaja seheriæ</i>	1004	<i>Anser anser</i>	577, 820

	PAGE		PAGE
Anser indicus	577, 820	Augiades siva	629
Anthracoseros coronatus albirostris...	819	———— subhyalina subhyalina ...	629
Anthus campestris griseus	873	———— thibetana	629
———— cervinus	873	———— sylvanoides	629
———— hodgsoni	872	———— sylvanus majuscula...	629
———— jerdoni	872	———— sylvanus	629
———— maculatus	872	———— venata	629
———— richardi rufulus	872	Autallacta subobliquaria	781
———— rosaceus... ..	873, 1004	Auzea apicata	797
———— similis	872	Balænoptera indica	815
———— sordidus decaptus	872	Baoris agna	633
———— spinoletta blakistoni	873	———— alicia	634
———— coutelli	873	———— amalia	635
———— trivialis trivialis	872	———— apostata	636
Anthyperrythra hermearia	795	———— assamensis	634
Apatura cooperi	579	———— aurociliata	633
———— ulupi florenciæ	580	———— austeni	633
———— kalaurica	580	———— bada	681
———— ulupi	580	———— balarama	633
Aphanteles acherontiae	726	———— bevani bevani	636, 676
Aplochlora viridis	786	———— ——— thyone	636
Apodemus sylvaticus griseus	605, 611	———— bipunctata	636
Apolectus niger	774	———— brunnea	632
Aquila hastata	571	———— cahira austeni	633
———— nipalensis nipalensis	571	———— ——— bromus	633
———— rapax vindhiana... ..	571	———— ——— cahira	633
Arboricola torqueola torqueola	818	———— ——— carna	633
Arceuthornis viscivorus bonapartei ...	985	———— ——— nirwana	633
Arctomys littledalei	612	———— ——— sirius	633
Ardea cinerea cinerea	576	———— canaraica	636, 684
Ardeola grayii	577, 820	———— ceramica	633
Argya caudata caudata	694	———— chaya	632
———— eclipses	694	———— chimdroa	633
———— huttoni	694	———— cinnara	636
———— earlii	694	———— coere	632
Arichanna plagifera	947	———— cœrulescens	636
———— ramosa	947	———— colaca	675
———— transectata	947	———— conjuncta conjuncta	634, 672
Artamus fuscus	827	———— ——— narooa	634
Asio flammeus flammeus	1005	———— connasa	634
Aspilates falconaria	790	———— contigua	635
———— obliquaria	791	———— cormasa	634
———— trilinearia	795	———— discreta discreta... ..	635
Asthena lassa	780	———— ——— himalaya	635
Astrape dipterygia	828	———— ella... ..	633
Astur badius dussumieri	572	———— eltola	635
Augiades boudhdha	629	———— flava	632
———— brahma	630	———— flexilis	636
———— crataeis	630	———— fortunet	636
———— ochracea	629	———— guttatus bada	636
———— similis... ..	629	———— ——— guttatus	636, 676

	PAGE		PAGE
Baoris haga	636	Bapta aluta	785
— intermedia	636	— distans	785
— jansonis	636	— ectiptica	785
— jolanda	634	— platyleucata	785
— juliana	633	Baracus hampsoni	655
— kolantus	636	Baxa orellana	909
— kumara kumara	634, 667	Blepharoctenucha virescens	944
— — lanka	634	Boarmia conjunctaria	798
— — malaya	634	— cyclophora	799
— — moorei	634	— fimbriata	942
— mangala	636	— melanosticta catotæniaria ..	942
— mathias mathias	632	— retrahens	942
— — midea	632	— virguncula	942
— — thrax	632	Bombycilla garrulus centralasiæ	867
— mencia	633	Brachypternus aurantius aurantius	567
— minuta	635	— — bengalensis dilutus	877
— monilis	633	Bubo benghalensis	570
— nascens	633	— — bubo benghalensis	1006
— neophytes	632	— — coromandus	570, 1006
— niasica	633	Bubulcus ibis coromandus	577, 820
— nondoa	636	Buchanetes githaginea crassirostris	863
— oceia farri... ..	632, 663	— — mongolicus	863
— — leechi	632	Bungarus fasciatus	564
— — oceia... ..	632	— — multicinctus	564
— — penicillata	632	Burhinus oedinemus indicus	574
— — simillima	632	Butastur teesa	571
— pagana	633	Buteo buteo rufiventer	1006
— pellucida	635	— — ferox	572
— philino	636	Butorides striatus javanicus	577, 1007
— philippina philippina	634, 667	Cacomantis passerinus	1032
— — seriata	634, 669	Cadaba farinosa	901
— — philotas	636	— — indica	901
— — plebeia	632, 667	Calandrella acutirostris acutirostris	875
— — prominens	632	— — — tibetana	1004
— — quinigera	636	— — — brachydactyla longipennis	874, 1004
— — scopulifera	632	— — — raytal adamsi	875
— — scortea	635	— — — rufescens leucophæa	875
— — seriata	669	— — — persica	875
— — sidata	636	— — — seebohmi	875
— — sifa	636	— — — scistanica	875
— — sikkima... ..	632	Callacanthis burtoni	996
— — similis	632	Callerinmys obliquilinea	792
— — sinensis sinensis	632	Calliope pectoralis pectoralis	983
— — sodalis	632	— — — tschbaiewi	984
— — subochracea	632, 679	Callophis maclellandi	566
— — toona	635	Collix lentiginosaria	796
— — tulusi	634	Caloris assamensis	663
— — unicolor	632, 663	— — aurociliata	663
— — zelleri colaca	636	— — austeni	663
— — — zelleri	636	— — bevani	663
Bapta alba	785		

	PAGE		PAGE
<i>Caloris cahira</i>	663	<i>Caprimulgus europæus plumipes</i> ...	880
———— canaraica	684	—————— unwini	822, 880
———— colaca	663	—————— indicus jataka	1005
———— conjuncta	663	—————— macrurus nipalensis ...	1005
———— contigua	663	—————— mahrattensis	821, 880
———— discreta... ..	663	—————— monticolus	821, 1005
———— eltola	663	<i>Carangoides malabaricus</i>	774
———— kumara... ..	663, 667	<i>Caranx djeddaba</i>	774
———— moolata	663	———— kalla	774
———— onchisa	663	<i>Carduelis caniceps caniceps</i> ...	864, 995
———— pagana... ..	663	—————— orientalis	864
———— plebeia	663	—————— paropanisi	864
———— seriata	663	—————— subcaniceps	864
———— tulsi	663	—————— subulata	864
———— uma	663	———— carduelis major	864
<i>Canis aureus</i>	601, 852	<i>Carine noctua brama</i>	570
———— dukhunensis	1029	<i>Carpodacus erythrinus roseatus</i> ...	863, 994
———— indicus	610	—————— rhodochlamys grandis] ...	863
———— naria	594, 614	—————— rubicilla severtzovi ...	995
———— pallipes	852	—————— rubicilloides	995
<i>Capella stenure</i>	1033	<i>Carpospiza brachydactyla</i>	866
<i>Capparis aphylla</i>	904	<i>Casearia ferruginea</i>	577, 820
———— bisperma	905	—————— esculenta	916
———— brevispina	903	—————— graveolens	915
———— cleghormii	905	—————— rubescens	916
———— dalzellii	907	—————— tomentosa	916
———— decidua	904	<i>Catreus wallichi</i>	818
———— divaricata	904	<i>Centropus sinensis sinensis</i> ...	569, 819, 881
———— diversifolia	905	<i>Cephalopyrus flammiceps</i>	992
———— galeata	903	<i>Certhia familiaris hodgsoni</i>	980
———— grandis	905	———— himalayana	980
———— heyneana	904	—————— limes	695
———— horrida	906	<i>Cervus hanglu</i>	814
———— incanescens	906	<i>Ceryle lugubris guttulata</i>	818
———— leucophylla	903	———— rudis leucomelanura	568, 879
———— longispina	909	<i>Cettia cetti cettoides</i>	700
———— moonii	905	<i>Chaimarrhornis leucocephala</i>	709, 983
———— murrayana	903	<i>Chalcophaps indica</i>	819
———— pedunculosa	906, 907	<i>Chapra brunnea</i>	663
———— rotundifolia	903, 907	———— mathias	663, 678
———— roxburghii	905	———— midea	663
———— sepiaria	906	———— sinensis	663
———— soddada	904	———— subochracea	663, 679
———— spinosa	903	<i>Charaxes durnfordi merguia</i>	579
———— stylosa... ..	904	<i>Charronia flavigula</i>	602
———— tenera	907	<i>Chaulelasmus streperus</i>	577, 820
———— tomentosa	905	<i>Chiasmia levata</i>	787
———— vulgaris	903, 906	———— strigata	787
———— zeylanica	906	<i>Chibia hottentotta hottentotta</i> ...	1001
<i>Caprilia specularia</i>	791	<i>Chlamydotis undulata macqueeni</i> ...	574, 1007
<i>Caprimulgus asiaticus</i>	569	<i>Choriotes edwardsi</i>	574

	PAGE		PAGE
<i>Chrysopela ornata</i>	564	<i>Coluber oxycephalus</i>	563
<i>Cimicodes castanearia</i>	797	----- <i>porphyraceus</i>	562
----- <i>cruentaria</i>	797	----- <i>prasinus</i>	562
<i>Cinclus cinclus kashmiriensis</i>	993	----- <i>radiatus</i>	562
----- <i>pallasi tenuirostris</i>	817, 993	<i>Columba leuconota leuconota</i>	818
<i>Cinnyris asiatica brevirostris</i>	877	----- <i>livia intermedia</i>	573
<i>Circus macrourus</i>	572	<i>Colymbus arcticus</i>	1008
<i>Cissampelos pareira</i>	555	<i>Copsychus saularis saularis</i>	819, 923
<i>Cisticola exitis courtoisi</i>	801	<i>Coracias benghalensis affinis</i>	819
<i>Clamator jacobinus</i>	1005	----- <i>benghalensis benghalensis</i>	567, 877
----- <i>jacobinus</i>	569	----- <i>garrula semenowi</i>	878
<i>Cleome angustifolia</i>	899	<i>Corvus collaris</i>	692
----- <i>aspera</i>	899	----- <i>corax laurencei</i>	691
----- <i>brachycarpa</i>	899	----- <i>ruficollis</i>	691
----- <i>burmanni</i>	899	----- <i>tibetanus</i>	976
----- <i>chelidonii</i>	900	----- <i>cornix sharpei</i>	692
----- <i>felina</i>	900	----- <i>corone orientalis</i>	691, 976
----- <i>glauca</i>	899	----- <i>coronoides andamanensis</i>	819
----- <i>longipetiolata</i>	899	----- <i>intermedius</i>	817, 976
----- <i>monophylla</i>	898	----- <i>frugilegus</i>	692
----- <i>papillosa</i>	898	----- <i>monedula soemmeringii</i>	692, 977
----- <i>pentaphylla</i>	900	----- <i>splendens splendens</i>	819, 823, 922
----- <i>quinquenervia</i>	898	----- <i>zugmayeri</i>	692, 977
----- <i>separatum</i>	899	----- <i>tschusii</i>	692
----- <i>simplicifolia</i>	898	<i>Corycia alba</i>	785
----- <i>stocksiana</i>	898	<i>Corymica immaculata</i>	791
----- <i>tinella</i>	899	----- <i>specularia</i>	791
----- <i>viscosa</i>	899	<i>Cossypha gutturalis</i>	709
<i>Cleora aagostigma</i>	939	<i>Coturnix coturnix coturnix</i>	574
----- <i>alienaria</i>	938	<i>Crataeva religiosa</i>	901
----- <i>cinnina</i>	939	<i>Cremnomys australis siva</i>	596
----- <i>coniozona</i>	940	<i>Creteus cyrina cyrina</i>	617
----- <i>cryptogonia</i>	941	----- <i>parca</i>	617
----- <i>decussata</i>	939	----- <i>meleagrina</i>	617
----- <i>euphiles</i>	940	<i>Crocallis acutaria</i>	797
----- <i>eurydiscaria</i>	939	----- <i>bilinearia</i>	796
----- <i>imbecilis</i>	939	<i>Crocidura kingiana</i>	601
----- <i>incongruaria</i>	940	----- <i>rubricosa</i>	600, 609
----- <i>latifasciata</i>	938, 939	<i>Crocopus phoenicopterus</i>	573
----- <i>pammicra</i>	939	----- <i>phoenicopterus</i>	801, 818, 1006
----- <i>perspicuata</i>	940	<i>Crypsicometes homœma</i>	788
----- <i>semiclarata</i>	940	<i>Ctenognophos methoria</i>	798
----- <i>sericea</i>	942	<i>Cuculus canorus</i>	880
----- <i>trigrapta</i>	940	----- <i>telephonus</i>	818
----- <i>variegata</i>	938	----- <i>subtelephonus</i>	881
----- <i>vatia</i>	941	----- <i>telephonus</i>	881, 1005
----- <i>venustularia</i>	939	<i>Culicicapa ceylonensis ceylonensis</i>	988
<i>Coccothraustes coccothraustes humii</i>	862	<i>Cuncuma leucorypha</i>	571
<i>Cocculus hirsutus</i>	552	<i>Cuon dukhenensis</i>	852
----- <i>pendulus</i>	552	<i>Cupitha alara</i>	623
<i>Cochlospermum gossypium</i>	910		

	PAGE		PAGE
<i>Cupitha lycorias</i>	623	<i>Dyspteris asiatica</i>	782
——— <i>purreea</i>	623, 659	<i>Echinops echinatus</i>	1039
——— <i>tympanifera</i>	623	<i>Ectropis æthregenes</i>	933
——— <i>verruca</i>	623	<i>Ectropis albibasis</i>	934
<i>Cursorius coromandelicus</i> ...	575, 820	——— <i>bisinuata dignampta</i> ...	936
<i>Cynopterus sphinx</i>	593	——— <i>chrysoteucta</i>	799
<i>Cyanosylvia cyanecula abbotti</i> ...	983	——— <i>conjunctaria</i>	798
——— <i>suecica pallidogularis</i> ...	709, 983	——— <i>conspicuata</i>	936
——— <i>suecica</i>	709	——— <i>cyclophora</i>	799
<i>Cyclea fissicalyx</i>	557	——— <i>defervescens</i>	933
——— <i>peltata</i>	556	——— <i>dentilineata</i>	937
<i>Cylindrophis rufus</i>	559	——— <i>duplexa</i>	932
<i>Cynælurus jubatus</i>	850	——— <i>embolochroma</i>	935
<i>Cynoglossus macrolepidotus</i> ...	772	——— <i>enormis</i>	932
<i>Cyornis superciliaris superciliaris</i> ...	987	——— <i>hiulca</i>	935
——— <i>tricolor tricolor</i>	987	——— <i>impos</i>	936
<i>Cyrtostomus asiaticus asiaticus</i> ...	817	——— <i>inceptaria</i>	936
<i>Dafila acuta</i>	577, 820	——— <i>indistincta</i>	936
<i>Decetia tinctaria</i>	789	——— <i>insecura</i>	936
<i>Delichon urbica</i>	869, 999	——— <i>intractabilis</i>	937
——— <i>cashmeriensis</i>	999	——— <i>irrorata</i>	933
<i>Delphinus delphis</i>	816	——— <i>leucozona</i>	933
<i>Dendrocitta rufa pallida</i>	692	——— <i>lophomeris</i>	933
<i>Dendrocitta rufa rufa</i>	817	——— <i>marmorata</i>	932
<i>Dendrocitta sinensis himalayensis</i> ...	817	——— <i>nigrosparsa</i>	799
<i>Dendrocygna javanica</i>	577, 820, 1008	——— <i>ochrifasciata</i>	932
<i>Dendrophis pictus</i>	563	——— <i>planaria</i>	932
<i>Dendrotreron hodgsoni</i>	818	——— <i>pulvicopia</i>	937
<i>Dicrurus leucophæus longicaudatus</i> ...	989	——— <i>simplaria</i>	933
——— <i>macrocerus</i>	819	——— <i>sp.</i>	799
——— <i>albirictus</i>	817	——— <i>zotica</i>	934
——— <i>macrocerus</i>	695	<i>Eetion ayankara</i>	616
<i>Dilophodes elegans</i>	950	——— <i>eburus</i>	616
<i>Diploclisia glaucescens</i>	551	——— <i>elia</i>	616
<i>Diplurodes vestita</i>	937	——— <i>magniplaga</i>	616
<i>Dipodillus indus</i>	610	<i>Egretta intermedia intermedia</i> ...	577
<i>Dipsadomorphus hexagonatus</i> ...	564	——— <i>spp.</i>	577
——— <i>multimaculatus</i>	564	<i>Elanus coeruleus coeruleus</i>	572
<i>Dipterygium glamum</i>	900	<i>Elephas indicus</i>	720, 813
<i>Discoloxia nigrifurca</i>	782	——— <i>maximus</i>	853
——— <i>purpuraria</i>	782	<i>Emberiza auricola</i>	868, 1003
<i>Dissemurus paradiseus grandis</i> ...	819	——— <i>buchanani</i>	867
<i>Dissoura episcopa episcopa</i>	576, 820	——— <i>cia par</i>	867, 1003
<i>Distira cyanocinctus</i>	828	——— <i>fucata arcuata</i>	998
<i>Dophla cooperi</i>	580	——— <i>icterica</i>	868
<i>Drepane punctata</i>	779	——— <i>leucocephala</i>	867, 998
<i>Dryobates himalayensis</i>	817	——— <i>luteola</i>	868
——— <i>macei macei</i>	817, 1004	——— <i>melanocephala</i>	868
——— <i>scindeanus</i>	877	——— <i>rutila</i>	996
<i>Dupetor flavicollis flavicollis</i> ...	820	——— <i>schoeniclus pallidior</i> ...	867
<i>Duranta ellisi</i>	727	——— <i>stewarti</i>	867, 998

	PAGE		PAGE
<i>Emberiza stracheyi</i>	868, 998	<i>Funambulus sublineatus</i> ...	595, 614
— — — — — <i>striolata striolata</i> ...	868, 1003	— — — — — <i>wroughtoni</i>	595
<i>Enantiodes consanguinea</i>	798	<i>Gabaria subalbataria</i>	794
<i>Enicurus maculatus maculatus</i> ...	982	<i>Galerida cristata chendoola</i> ...	875
<i>Eogenes ahrimani</i>	637	— — — — — <i>cristata magna</i>	875
— — — — — <i>alcides</i>	637	— — — — — <i>magna</i>	875
<i>Eoglaucmys fimbriatus</i>	603, 610	<i>Gallicrex cinerea</i>	537, 1007
<i>Epione adustata</i>	792	<i>Gallinago gallinago gallinago</i> ...	575
— — — — — <i>obliquilinea</i>	792	— — — — — <i>nemoricola</i>	728, 1033
<i>Erolia temminckii</i>	575	— — — — — <i>solitaria</i>	729, 818
<i>Eudynamis scolopaceus malayanus</i> ...	819	<i>Gallinula chloropus</i>	541
— — — — — <i>scolopaceus</i>	569, 1032	— — — — — <i>parvifrons</i>	541, 574
<i>Eumenes conica</i>	754, 890	— — — — — <i>poliocephalus</i>	533
— — — — — <i>dimidiatipennis</i>	758	<i>Gallus gallus gallus</i>	819
<i>Eupetaurus cinereus</i>	612	<i>Garæus apicata</i>	797
<i>Falco</i> sp.	572	— — — — — <i>cruentatus</i>	797
— — — — — <i>chiquera chiquera</i>	572	— — — — — <i>specularis</i>	797
— — — — — <i>peregrinus peregrinator</i> ...	1006	<i>Garrulax albogularis whistleri</i> ...	817
— — — — — <i>tinnunculus tinnunculus</i> ...	572, 818	— — — — — <i>leucolophus leucolophus</i> ...	817, 1000
<i>Fascellina chromataria</i>	795	— — — — — <i>bispecularis bispecularis</i> ...	817
— — — — — <i>inornata</i>	796	— — — — — <i>lanceolatus</i>	817
— — — — — <i>plagiata</i>	796	<i>Gazella bennettii</i>	612
<i>Felis affinis</i>	594, 601	<i>Gegenes karsana</i>	637, 661
— — — — — <i>bengalensis</i>	594, 609	— — — — — <i>lefebvrei</i>	637
— — — — — <i>domesticus</i>	601	— — — — — <i>nostrodamus</i>	637, 661, 951
— — — — — <i>leo</i>	847	— — — — — <i>pumilio</i>	637
— — — — — <i>ornata</i>	609	— — — — — <i>pygmæus</i>	637
— — — — — <i>pardus</i>	601, 808, 809, 849	<i>Gehenna abima abima</i>	617
— — — — — <i>tigris</i>	810, 831, 848	— — — — — <i>angulifera</i>	617
<i>Flacourtis inermis</i>	914	— — — — — <i>dawna</i>	617
— — — — — <i>integrifolia</i>	914	— — — — — <i>greæ</i>	617
— — — — — <i>latifolia</i>	913	<i>Gennæus hamiltoni</i>	818, 1006
— — — — — <i>montana</i>	912	— — — — — <i>horsfieldi horsfieldi</i> ...	730, 819
— — — — — <i>occidentalis</i>	914	<i>Geometra dentisignata</i>	786
— — — — — <i>ramontchi</i>	912	— — — — — <i>lutea</i>	795
— — — — — <i>sepiaria</i>	914	— — — — — <i>viretata</i>	782
<i>Francolinus francolinus asiæ</i>	574, 818	<i>Gerbillus gleadowi</i>	610
— — — — — <i>melanonotus</i>	819	<i>Gerres filamentosus</i>	776
— — — — — <i>gularis</i>	819	<i>Glaucidium brodiei</i>	1006
— — — — — <i>pondicerianus interpositus</i> ...	574	<i>Gnophos approximaria</i>	798
<i>Franklinia buchanani</i>	697	— — — — — <i>cinerea</i>	798
<i>Fringilla montifringilla</i>	865	<i>Golunda ellioti</i>	605
<i>Fringillauda brandti hæmatopyga</i> ...	998	— — — — — <i>myothrix</i>	605, 611
— — — — — <i>nemoricola altaica</i>	997	<i>Gonodela khasiana</i>	793
<i>Fulica atra atra</i>	544, 574	<i>Gonodontis acutaria</i>	797
— — — — — <i>cinerea</i>	537	— — — — — <i>bilinearia</i>	796
<i>Funambulus gossei</i>	614	— — — — — <i>insulata</i>	797
— — — — — <i>palmarum comorinus</i>	614	— — — — — <i>variegata</i>	797
— — — — — <i>palmarum</i>	595	<i>Graucalus macei</i>	1002
— — — — — <i>pennanti</i>	603	<i>Gunomys kok</i>	596
— — — — — <i>argentescens</i>	610	— — — — — <i>wardi</i>	604

	PAGE		PAGE
<i>Gymnorhis xanthocollis transfuga</i> ...	865	<i>Halpe perfossa</i> ...	621
<i>Gynandropsis pentaphylla</i> ...	900	— <i>selangora</i> ...	621
<i>Gypætus barbatus grandis</i> ...	818	— <i>separata</i> ...	618
<i>Gyps fulvus fulvescens</i> ...	570	— <i>sikkima palawea</i> ...	621
— <i>himalayensis</i> ...	818	— <i>sikkima</i> ...	621
<i>Halcyon smyrnensis</i> ...	568	— <i>sitala</i> ...	619
— <i>fusca</i> ...	818	— <i>subflava</i> ...	618
— <i>smyrnensis</i> ...	879	— <i>submacula</i> ...	620
<i>Haliastur indus indus</i> ...	819	— <i>sulphurifera</i> ...	622
<i>Halpe aina</i> ...	619	— <i>teliga</i> ...	621
— <i>albipectus</i> ...	618	— <i>thandaunga</i> ...	619
— <i>albipectus</i> ...	618	— <i>varia</i> ...	619
— <i>albivitta</i> ...	619	— <i>wantona</i> ...	622
— <i>astigmata</i> ...	619	— <i>zema ormenes</i> ...	621
— <i>auriferus auriferus</i> ...	622	— <i>zema</i> ...	621
— <i>peletronix</i> ...	622	<i>Harpiocephalus harpia madrassius</i> ...	593
— <i>bazilana</i> ...	621	<i>Harpodon nehereus</i> ...	772
— <i>beturina</i> ...	621, 622	<i>Hemerophila cupsearia</i> ...	944
— <i>bivitta</i> ...	619	— <i>retractaria</i> ...	944
— <i>blanchardi</i> ...	619	<i>Hemichelidon sibirica gulmergi</i> ...	705, 981
— <i>brunnea</i> ...	622	<i>Hemidactylus sp.</i> ...	827
— <i>burmana</i> ...	620	<i>Hemitragus jemlaicus</i> ...	812
— <i>cerata</i> ...	620	<i>Herodias alba alba</i> ...	820
— <i>debitis</i> ...	619	— <i>garzetta garzetta</i> ...	820
— <i>decorata</i> ...	620	<i>Herpestes auropunctatus pallipes</i> ...	610
— <i>egena egena</i> ...	622	— <i>edwardsi carnaticus</i> ...	594, 613
— <i>evershedii</i> ...	620	— <i>ferrugineus</i> ...	601
— <i>fasciata</i> ...	621	— <i>fuscus</i> ...	594
— <i>flava</i> ...	622	— <i>viticollis</i> ...	594
— <i>grahami</i> ...	618	<i>Heterogomphus pars</i> ...	733
— <i>gupta</i> ...	619	<i>Heterolocha falconaria</i> ...	790
— <i>hazis</i> ...	622	— <i>patalata</i> ...	791
— <i>hieron</i> ...	622	— <i>segregis</i> ...	791
— <i>homolea ceylonica</i> ...	622	<i>Heterostegania nigrofusa</i> ...	788
— <i>homolea</i> ...	622	— <i>thamia</i> ...	788
— <i>honorei</i> ...	620	<i>Hidari bhawani</i> ...	616
— <i>hyrie</i> ...	620	— <i>doesœna</i> ...	616
— <i>hyrtacus</i> ...	618	— <i>hypœpa</i> ...	616
— <i>insignis</i> ...	620	— <i>irava</i> ...	616
— <i>joloana</i> ...	621	— <i>ozias</i> ...	616
— <i>knyvetti</i> ...	621	— <i>thrax</i> ...	616
— <i>kumara</i> ...	621	<i>Hieraetus fasciatus</i> ...	571
— <i>latris</i> ...	619	— <i>pennatus</i> ...	1006
— <i>lucasi</i> ...	620	<i>Hierococcyx varius</i> ...	569, 818, 923
— <i>luteisquama</i> ...	622	<i>Hilsa blochii</i> ...	771
— <i>majuscula</i> ...	621, 622	<i>Himantopus himantopus himantopus</i> ...	575
— <i>marta</i> ...	622	<i>Hippolais caligata</i> ...	698
— <i>masoni</i> ...	620	— <i>languida</i> ...	698
— <i>moorei</i> ...	621	— <i>pallida</i> ...	692
— <i>nephele</i> ...	622	— <i>rama</i> ...	697, 1001
— <i>perara</i> ...	622	— <i>scita</i> ...	698

INDEX OF SPECIES

xxv

	PAGE		PAGE
Hirasa contubernalis	944	Ixulus flavicollis flavicollis	1000
-----plagiochorda	944	Iyngipicus hardwickii	1005
-----scripturaria	944	-----pygmaeus	1004
Hirundo daurica erythropyga	1003	lynx torquilla torquilla	567, 877
-----nepalensis 578, 817, 1003		Johnius aeneus	776, 777
-----rufula	870	-----carutta	778
-----striolata	578	-----diacanthus	777
-----rustica rustica	870, 999	-----glaucus	778
-----smithi filifera	870	Kogia breviceps	817
Hodgsonius phoenicuroides phoeni-		Krananda orthotmeta	792
curoides	981	Laiscopus collaris whympersi	986
Holorista usta	783	-----himalayanus	986
Homalium zeylanicum	917	Lanius colluris	702
Horornis pallidus pallidus	992, 1001	-----cristatus phoenicuroides	702, 1035
Horsfeldia anita gigantea	587	-----excubitor	701
Hyaena striata	852	-----aucheri	701
Hyarotis adrastus	967	-----lahtora	701
Hydnocarpus wightiana	915	-----pallidirostris	701
Hydrelia bicolorata	781	-----przewalskii	702
-----enisaria	781	-----isabellinus	703, 988
-----opedogramma	781	-----minor	702
-----purpuraria	782	-----schach erythronotus	702, 988
-----rufinota	781	-----senator niloticus	702
Hydrochelidon leucopareia indica	575	-----vittatus	702, 817, 988
Hydrophasianus chirurgus	1007	Lantana camera	727
Hylobates lar	846	Larvivora brunnea	981
Hypacanthis spinoides spinoides	999	Laticauda laticauda	564
Hyperythra calcearia	796	Laticilla burnesi	697
-----spurcatoria	795	Leggada booduga	596, 605
Hypocolius ampelinus	694	-----palnica	596
Hyprochrosis ancylotoxa	790	Leggadilla platythrix	596, 604
-----eurynota	789	Lepterodius asha	823
-----hyadaria	689	Leptocoma asiatica asiatica	920
-----mixticolor	789	Leptogomphus gestroi	884
-----tinctaria	789	-----inclitus	883
Hyria bicolorata	781	-----lansbergei	885
Iambrix salsala	971	-----maculivertex	886
Ianthia cyanura rufilata	984	Leptomiza calcearia	796
Ianthocincla lineata ziaratensis	694	Leptoptilus javanicus	820
Ichthyophis monochrous	1039	Leptostegna asiatica	782
Icterodes jaguarinaria	947	Lepus dayanus	612
Ictinaetus malayensis perniger	571	-----nigricollis	597, 614
Idaea hyalinata	787	-----oiostolus	613
Ilisca indica	771	-----ruficaudatus	605
Inocotis papillosus papillosus... ..	576, 820	Liminitis trivena gilgitica	581
Ionidium suffruticosum	908	-----hydaspes	581
Iton azona	631	-----lepichini	581
-----semamora	631	-----ligyes	581
-----watsonii	631	-----pallida	581
Ixobrychus cinnamomeus	1008	-----trivena	581
-----minutus minutus	1008	Liopeltis frenatus	563

	PAGE		PAGE
<i>Liopicus mahrattensis</i>	567	<i>Mærua ascabra</i>	901
<i>Lioptila capistrata capistrata</i>	817	——— <i>ovalifolia</i>	900, 901
<i>Liponycteris kachhensis</i>	608	<i>Manis crassicaudatus</i>	614
<i>Lobivanellus indicus</i>	820	<i>Marcala obliquaria</i>	789
——— <i>indicus</i>	575, 818	<i>Mareca penelope</i>	577, 820
<i>Locustella certhiola</i>	801	<i>Marmota himalayana</i>	603
<i>Lomographa tenebrimedia</i>	787	——— <i>littledalei</i>	612
<i>Lophobates ochreicostata</i>	943	<i>Martes foina</i>	602
<i>Lophoceros birostris</i>	568	<i>Medasina albidaria</i>	946
<i>Lophophanes rufonuchalis rufonuchalis</i>	817	——— <i>basistrigaria</i>	946
<i>Lophophorus impejanus</i>	818	——— <i>combustaria</i>	945
<i>Loxaspilates obliquaria</i>	791	——— <i>lasiochora</i>	946
——— <i>trilinearia</i>	795	——— <i>livida</i>	946
<i>Loxotephria perileuca</i>	793	——— <i>objectaria</i>	945
<i>Luscinia megarhyncha golzii</i>	709	——— <i>parallela</i>	946
<i>Lusciniola melanopogon mimica</i>	697	——— <i>tephrosiaria</i>	945
<i>Lutjanus johnii</i>	774	<i>Megaderma spasma horsfieldi</i>	593
<i>Lutra lutra nair</i>	595, 603	<i>Megalæma virens marshallorum</i>	817, 825, 1005
<i>Luxiaria amasa fasciosa</i>	792	<i>Megalornis antigone antigone</i>	574, 820
——— <i>mitorrhaphes</i>	792	<i>Melanocorypha bimaculata</i>	874, 1004
——— <i>tephrosaria</i>	792	<i>Melitæa balba balbina</i>	583
<i>Lycæna astorica</i>	584	——— <i>saxitalis lunulata</i>	583
——— <i>baroghila</i>	583	<i>Melophus melanicterus</i>	817
——— <i>christophi lesliei</i>	584	<i>Mergus albellus</i>	578
——— <i>florenciæ</i>	583	<i>Merops apiaster</i>	878
——— <i>galathea chitralica</i>	585	——— <i>orientalis beludschicus</i>	878
——— <i>depreei</i>	585	——— <i>orientalis</i>	567
——— <i>metallica</i>	584	——— <i>persicus persicus</i>	567, 878
——— <i>chitralensis</i>	584	——— <i>philippinus</i>	878
——— <i>orbitulus astorica</i>	585, 586	——— <i>superciliosus javanicus</i>	567, 1005
——— <i>ellisi</i>	585, 586	<i>Metaponia pusilla</i>	996, 1002
——— <i>jaloka</i>	585, 586	<i>Micrabraxas anisonoma</i>	946
——— <i>leela</i>	586	<i>Microcichla scouleri scouleri</i>	982
——— <i>walli</i>	586	<i>Microhierax eutolmus</i>	826
<i>Lycodon aulicus</i>	561	<i>Micropus affinis affinis</i>	569
——— <i>fasciatus</i>	562	——— <i>galilegensis</i>	880
<i>Lymnocryptes minimus</i>	575	——— <i>apus pekinensis</i>	880
<i>Lyroderma lyra</i>	593, 608	——— <i>melba</i>	879, 1005
<i>Macaca ferox</i>	592	——— <i>murinus murinus</i>	880
——— <i>mulatta</i>	599, 608	——— <i>pacificus leuconyx</i>	818
——— <i>radiata</i>	592	<i>Microscelis psaroides psaroides</i>	980
<i>Macacus rhesus</i>	845	<i>Microtus (Hyperacrius) wynei</i>	605, 611
<i>Macaria monticolaria</i>	794	<i>Millardia meltada</i>	596, 611
——— <i>myandaria</i>	794	——— <i>pallidior</i>	604
<i>Macrogomphus annulatus</i>	734	<i>Milvus migrans govinda</i>	572, 879
——— <i>montanus</i>	734, 738	<i>Mirafra assamica</i>	1004
——— <i>robustus</i>	734, 737	——— <i>cantillans</i>	875
——— <i>seductus</i>	733, 734, 737	——— <i>erythroptera sindianus</i>	875
——— <i>wynaadicus</i>	734, 736	<i>Molpastes hæmorrhous intermedius</i>	817, 1001
<i>Mærua arenaria</i>	900	——— <i>pallidus</i>	1001
——— <i>var. scabra</i>	900	——— <i>leucogenys</i>	817

	PAGE		PAGE
Molpastes leucogenys leucogenys ...	979	Natrix clerki ...	560
Monticola cinclorhyncha ...	985	—— gilhodesi ...	559
—— erythrogaster... ..	1002	—— khasiensis ...	559
—— saxatilis ...	711	—— trianguligera ...	560
—— solitaria longirostris ...	711	—— venningi ...	560
—— pandoo ...	710, 985	Nectogale sikkimensis ...	613
—— transcaspicus ...	711	Nemorhœdus goral ...	606, 812
Montifringilla nivalis adamsi ...	997	Neomeris phocænoides ...	816
—— alpicola ...	867	Neophron percnopterus ...	571
Moringa concanensis ...	907	—— percnopterus ginginianus ...	818, 922
—— oleifera ...	907	Neptis mackwoodi ...	582
—— pterygosperma ...	907	—— suffusa ...	582
Motacilla alba dukhunensis ...	870	—— zaida bhutanica ...	582
—— hodgsoni ...	1003	—— drummondi ...	583
—— personata ...	871	—— manipurensis ...	583
—— cinerea caspica ...	871	—— zaida ...	582
—— melanope ...	817, 819	Nerodia modesta ...	560
—— citreola calcaratus ...	872	—— piscator... ..	561
—— citreola ...	872, 1003	Nesokia griffithi ...	611
—— feldegg melanogriseus ...	871, 1003	Netta rufina ...	578, 820
—— flava beema ...	871, 1003	Nettion crecca crecca ...	577, 820
—— thunbergi ...	871, 1003	Nicevillea akar ...	623
—— melanope ...	871	—— alfurus ...	623
—— raii ...	871	—— concinna ...	623
Mungos mungo ellioti ...	613	—— fabriolata ...	623
Munia atricapilla ...	1002	—— gola ...	623
Muntiacus vaginalis ...	606	—— goloides ...	623
Mus gerbillinus ...	611	—— grandis ...	623
—— homourus ...	604	—— locus ...	623
—— musculus ...	596, 604	—— naranata ...	623
—— myothrix ...	605	—— nipata ...	623
Muscicapa striata neumanni ...	704	—— paragola ...	623
Mustela alpina ...	602	—— pseudolus ...	623
—— canigula ...	602	—— rajagriha ...	623
—— foina ...	602	—— taxilus... ..	623
—— hodgsoni ...	602	—— trishna ...	623
—— temon ...	602, 603	Nieuhria arenaria ...	900
—— whiteheadi ...	602	—— oblongifolia ...	901
Mycerobas carneipes speculigerus ...	862	Niphanda cymbia ...	587
Myiophonus horsfieldi temminckii ...	711, 817, 986	—— marcia ...	587
Myotis caliginosus ...	600	—— tessellata ...	587
—— moupinensis ...	612	Nothomiza ægriviridis ...	786
—— muricola ...	600	—— cinerascens ...	786
Myrteta fuscolineata ...	784	—— dentisignata ...	786
—— icuncula ...	783	—— ithyterma ...	786
—— ocernaria ...	783	—— obscuristrigata ...	786
Nadagara epopsioneura ...	792	—— simpliciaris... ..	786
Naia hannah ...	564	—— viridis ...	786
—— naia ...	565	Notocrypta feisthamelii ...	959
Natrix bitæniata ...	560	—— restricta ...	956
		Nucifraga caryocatactes hemispila ...	817

	PAGE		PAGE
<i>Nucifraga multipunctata</i> ...	977	<i>Otolithus ruber</i> ...	776
<i>Numenius arquatus lineatus</i> ...	575	<i>Otus sunia sunia</i> ...	570
<i>Nyctalus labiatus</i> ...	600	<i>Ourapteryx ebuleata</i> ...	783
—— <i>leisleri</i> ...	600	—— <i>multistrigaria</i> ...	783
<i>Nycticorax nycticorax nycticorax</i> ...	577, 1008	<i>Ovis ammon</i> ...	800
<i>Nyroca ferina ferina</i> ...	578, 820	—— <i>hodgsoni</i> ...	800
—— <i>fuligula</i> ...	578, 820	<i>Pachydia vexillaria</i> ...	795
—— <i>rufa</i> ...	1034	<i>Pachyura hodgsoni</i> ...	600
—— <i>rufa</i> ...	578, 820	—— <i>saturation</i> ...	612
—— <i>rufina</i> ...	1034	—— <i>sp.</i> ...	594, 600, 609
<i>Ochotona roylei</i> ...	605, 613	<i>Padraona ahastina</i> ...	627
—— <i>wollastoni</i> ...	613	—— <i>ahrendti</i> ...	627
<i>Ocœlophora agana</i> ...	796	—— <i>angustata</i> ...	627
—— <i>lentiginosaria</i> ...	796	—— <i>dara confucius</i> ...	627
<i>Oenanthe alboniger</i> ...	706	—— <i>dara</i> ...	627
—— <i>capistrata</i> ...	707	—— <i>dilutior</i> ...	627
—— <i>deserti atrogularis</i> ...	708	—— <i>pseudomoesa</i> ...	627
—— <i>oreophila</i> ...	708, 981	—— <i>scrina</i> ...	672
—— <i>finschii barnesi</i> ...	707	—— <i>dushta</i> ...	627
—— <i>isabellina</i> ...	708	—— <i>fettingi fettingi</i> ...	625
—— <i>leucomela</i> ...	707	—— <i>orfitus</i> ...	625
—— <i>monacha</i> ...	706	—— <i>ganda</i> ...	627
—— <i>œnanthe</i> ...	708	—— <i>hetærus</i> ...	627
—— <i>opistholeuca</i> ...	706	—— <i>chariyawa</i> ...	626
—— <i>picata</i> ...	706, 707	—— <i>hetærus</i> ...	626
—— <i>xanthoprymna chrysopygia</i> ...	708	—— <i>zatilla</i> ...	626
<i>Oenopopelia tranquebarica</i> ...	819	—— <i>kobros rectifasciata</i> ...	626
—— <i>tranque-</i>		—— <i>phellus phellus</i> ...	626
<i>barica</i> ...	573	—— <i>tanya</i> ...	626
<i>Oligodon albocinctus</i> ...	563	—— <i>mæsia</i> ...	627
—— <i>herberti</i> ...	563	—— <i>moesoides moesoides</i> ...	626
—— <i>purpurascens</i> ...	563	—— <i>ottala</i> ...	626
—— <i>splendidus</i> ...	563	—— <i>ottalina</i> ...	626
—— <i>theobaldi</i> ...	563	—— <i>nala</i> ...	627
<i>Onryza meiktila</i> ...	622	—— <i>nikaja</i> ...	627
—— <i>siamica</i> ...	622	—— <i>nitida</i> ...	627
<i>Onychogomphus circularis</i> ...	741	—— <i>palnia</i> ...	627
<i>Opisthograptis molleri</i> ...	791	—— <i>pava</i> ...	627
—— <i>swanni</i> ...	791	—— <i>pseudomoesa</i> ...	627
<i>Oreicola ferrea ferrea</i> ...	981	—— <i>purpura</i> ...	625
<i>Oreocincla dauma dauma</i> ...	1002	—— <i>sapitana</i> ...	627
<i>Oreocorys sylvanus</i> ...	873	—— <i>sunias satra</i> ...	627
<i>Oriolus indicus</i> ...	819	—— <i>sunias</i> ...	627
—— <i>oriolus kundoo</i> ...	703, 817, 992, 1002, 1032	—— <i>tropica</i> ...	627
—— <i>oriolus</i> ...	703	—— <i>trachala</i> ...	628
<i>Orogomphus campioni</i> ...	741	—— <i>yojana</i> ...	627
<i>Orsonoba æthocrypta</i> ...	794	—— <i>zebra</i> ...	627
<i>Orthomiella putaoica</i> ...	587	<i>Paguma grayi</i> ...	601
<i>Ostracion gibbosus</i> ...	779	<i>Palumbus palumbus casiotus</i> ...	1006
<i>Otocorys alpestris albigula</i> ...	874	<i>Pamphila comma comma</i> ...	630
—— <i>penicillata</i> ...	874	—— <i>demila</i> ...	630

	PAGE		PAGE
<i>Pamphila comma florinda</i> ...	630	<i>Phoenicurus hodgsoni</i> ...	709
————— <i>mixta</i> ...	630	————— <i>ochrurus phoenicuroides</i> ...	709, 817, 982
<i>Paradarisa azyx</i> ...	937	<i>Phthonoloba decussata</i> ...	782
————— <i>chloauges</i> ...	937	<i>Phylloscopus affinis</i> ...	990
<i>Paradoxurus jerdoni</i> ...	594	————— <i>collybita collybita</i> ...	699
————— <i>niger</i> ...	601	————— <i>tristis</i> ...	699
<i>Paræchinus blanfordi</i> ...	609	————— <i>collybitus sindianus</i> ...	991
<i>Parasynergia suffusa</i> ...	788	————— <i>tristis</i> ...	699, 991
<i>Parnara bada</i> ...	663	————— <i>griseolus</i> ...	991
————— <i>canaraica</i> ...	663	————— <i>humei</i> ...	700, 991
————— <i>flexilis</i> ...	663	————— <i>indica</i> ...	699
————— <i>guttata</i> ...	663	————— <i>neglectus neglectus</i> ...	700
————— <i>vaika</i> ...	663	————— <i>nitidus nitidus</i> ...	700
<i>Parus major intermedius</i> ...	693	————— <i>proregulus simlaensis</i> ...	991
————— <i>kashmiriensis</i> ...	817, 978	————— <i>tytleri</i> ...	900
————— <i>melanolophus</i> ...	693, 978	<i>Pica pica bactriana</i> ...	692
————— <i>monticolus</i> ...	817, 978	<i>Picus gorii</i> ...	877
————— <i>palustris korejewi</i> ...	693	————— <i>squamatus flavirostris</i> ...	877
————— <i>rufonuchalis rufonuchalis</i> ...	693, 979	————— <i>squamatus</i> ...	817
<i>Passer domesticus indicus</i> ...	819, 865	————— <i>striolatus</i> ...	1004
————— <i>parkini</i> ...	865, 997, 1003	<i>Pipistrellus babu</i> ...	606
————— <i>griseigularis</i> ...	866	————— <i>ceylonicus indicus</i> ...	593
————— <i>hispaniolensis transcaspicus</i> ...	866	————— <i>coromandra</i> ...	593
————— <i>montanus dilutus</i> ...	866	————— <i>kuhli lepidus</i> ...	608
————— <i>pallidus</i> ...	866	————— <i>mimus</i> ...	608
————— <i>pyrrhonotus</i> ...	866	<i>Pirdana albicornis</i> ...	617
————— <i>rutilans debitis</i> ...	997	————— <i>albidiscus</i> ...	617
<i>Pastor roseus</i> ...	703	————— <i>distanti</i> ...	617
<i>Pavo cristatus</i> ...	574, 819	————— <i>hyela</i> ...	617
<i>Pelecanus</i> sp. ...	576	————— <i>ismene</i> ...	617
<i>Pellorneum ruficeps jonesi</i> ...	1000	————— <i>niasica</i> ...	617
<i>Pennahia æneus</i> ...	776	————— <i>pavona</i> ...	617
<i>Peratophyga hyalinata</i> ...	787	————— <i>rudolphii</i> ...	617
<i>Pernia felinaria</i> ...	950	————— <i>sargon</i> ...	617
<i>Pericrocotus brevirostris brevirostris</i> ...	703, 989	————— <i>scanda</i> ...	617
————— <i>erythropygius</i> ...	1002	<i>Pithauria aithchisoni</i> ...	618
————— <i>peregrinus</i> ...	703	————— <i>glauca</i> ...	618
————— <i>speciosus speciosus</i> ...	817	————— <i>marsena</i> ...	618
<i>Perissogomphus stevensi</i> ...	742, 743	————— <i>murdava</i> ...	618
<i>Perissospiza icteroides icteroides</i> ...	993	————— <i>ornata</i> ...	618
<i>Petaurista fulvinus</i> ...	603	————— <i>straumineipennis</i> ...	618
————— <i>inornatus</i> ...	603, 610	————— <i>subornata</i> ...	618
————— <i>philippensis</i> ...	595, 614	<i>Pithecus entellus</i> ...	599
<i>Petelica vexillaria</i> ...	795	————— <i>anchises</i> ...	592
<i>Petronia petronia intermedia</i> ...	866	————— <i>johni</i> ...	592
<i>Phalacrocorax javanicus</i> ...	576	<i>Pitta brachyura</i> ...	1004
<i>Phalæna lutea</i> ...	795	<i>Plataela leucorodia</i> ...	819
<i>Philomachus pugnax</i> ...	1007	————— <i>major</i> ...	576, 820
<i>Phænicurus erythronotus</i> ...	709, 982	<i>Ploceus manyar flaviceps</i> ...	862, 1002
————— <i>erythrogaster grandis</i> ...	982	————— <i>philippinus philippinus</i> ...	862
<i>Phoenicurus frontalis</i> ...	982		

	PAGE		PAGE
Plutodes warreni	787	Pseudotantalus leucocephalus leucocephalus	1007
Pnoëpyga squamata	801	Psittacula cyanocephala cyanocephala	569, 818
Pœcilasthena burmensis	780	Psittacula schisticeps schisticeps	1005
Polanisia chelidonii	900	—————torquata	569, 818, 881
—————felina	900	Psyra spurcatoria	795
—————viscosa	899	Pterocles indicus	573
Polioaëtus ichthyaëtus	572	—————orientalis	573
Polydactylus plebejus	773	—————senegalensis erlangeri	574
—————sextarius	773	Pteromys albiventer	603
Polyodontophis collaris	559	—————inornatus	603
Polyommatus gigantea gilgitica	586	Pteropus giganteus	600
—————sarta gooraisica	586	Ptyas carinatus	562
—————rapala	586	—————mucosus	562
Polyplectron bicalcaratum bicalcaratum	819	Ptynoprogne obsoleta pallida	869
Polyscia ochrilenia	797	—————rupestris	869, 999
Pomadasis hasta	775	Pucrasia macrolopha macrolopha	818
—————maculatus	775	Putorius alpinus	602
Pomatorhinus erythrogenys erythro-		—————canigula	602
genys	1000	Putrius erminea	602
—————horsfieldi	1032	Pycnonotus leucotis humii	695
Porphyrio poliocephalus edwardsi	534	—————leucotis	694
—————poliocephalus	533, 820	—————mesopotamiae	695
—————seistanicus	534	Pycnorhis sinensis sinensis	1000
Porzana porzana	1007	Pyrhacorax alpinus	693
—————pusilla	1006	—————graculus	693, 978
Prinia gracilis lepida	700	—————pyrrhacorax	692, 817, 978
—————inornata inornata	701	—————himalayanus	692
—————socialis socialis	1032	Pyrrhospiza punicea humii	994
—————sylvatica sylvatica	1001	Pyrrhula aurantiaca	993
Prochasma albimonilis	943	Pyrrhulauda frontalis affinis	876
—————dentilinea	943	—————grisea siccata	876
Propasser rhodochrous	994	Python molurus	559
—————thurus blythi	994	—————reticulatus	559
Prunella atrogularis	711, 986	Querquedula querquedula	820, 826
—————fulvescens fulvescens	986	Rapala rubida prorosacea	590
—————ocularis	711	Rattus blanfordi	596, 614
—————rubeculoides	986	—————fulvescens	604
—————strophiatu jerdoni	711, 986	—————nitidus	612
Psammodynastes pulverulentus	564	—————rattus gangutrianus	604
Pseudasthena cinerea	798	—————rufescens	596, 604, 611, 614
Pseudochliarsa virgo	589	—————sikkimensis	612
Pseudogyps bengalensis	571	—————wroughtoni	596, 614
Pseudomacromia torrida	769	—————vicerex	604, 611
Pseudomija argentilinea eugraphes	797	Ratufa indica bengalensis	614
—————castanearia	797	—————maxima	595
—————cruentaria	797	—————macroura dandolena	595
Pseudorca crassidens	816	Regulus regulus himalayensis	992
Pseudorhombus russellii	772	—————tristis	696
Pseudosciaena amblyceps	777	Remodes fasciata	783
Pseudostegania plurilinearia	780		

	PAGE		PAGE
<i>Rhabdophis bicolor</i>	561	<i>Semiothisa clivicola</i>	794
————— <i>speciosus</i>	561	————— <i>khasiana</i>	793
————— <i>stolatus</i>	561	————— <i>monticolaria</i>	794
————— <i>subminiatus</i>	561	————— <i>myandaria</i>	794
<i>Rhinobatus rhinobatus</i>	770	————— <i>subalbataria</i>	794
<i>Rhinoceros indicus</i>	859	<i>Semnopithecus entellus</i>	845
————— <i>sondaicus</i>	860	<i>Serinus pussillus</i>	864
————— <i>sumatrensis</i>	860	<i>Seriola nigrofasciata</i>	773
————— <i>unicornis</i>	1032	<i>Sieboldius nigricolor</i>	888
<i>Rhinolophus beddomei</i>	593	<i>Siphia parva parva</i>	705, 987
————— <i>ferrum equinum</i> <i>tragatus</i>	600	————— <i>hyperythra</i>	705, 987
————— <i>rouxi</i>	593, 613	————— <i>strophia</i>	1002
<i>Rhinopoma hardwickei</i>	594, 609	<i>Sitta castaneiventris cinnamoventris</i> ...	817
————— <i>kinneari</i>	609	——— <i>kashmiriensis</i>	695, 979
<i>Rhipidura albicollis stanleyi</i>	817	——— <i>leucopsis leucopsis</i>	979
<i>Rhodospiza obsoleta</i>	864	——— <i>neumayer tephronota</i>	695
<i>Rhyacornis fuliginosa</i>	709, 983	<i>Somatina plurilineata</i>	780
<i>Rhynchops albicollis</i>	576	<i>Sotalia lentiginosa</i>	816
<i>Riparia chinensis</i>	869	<i>Sparus berda</i>	775
——— <i>riparia diluta</i>	869, 1003	——— <i>spinifer</i>	776
——— <i>riparia</i>	869	<i>Spatula clypeata</i>	578, 820
<i>Rostratula benghalensis benghalensis</i>	575, 730, 1007	<i>Sphyræna barracuda</i>	772
<i>Sancus subfasciatus</i>	962	<i>Stephania andamanica</i>	553
<i>Sarcidiornis melanota</i>	577	——— <i>elegans</i>	553
<i>Sarciophorus malabaricus</i>	575, 1007	——— <i>glabra</i>	553
<i>Satarupa splendens</i>	590	——— <i>gracilenta</i>	553
——— <i>zulla</i>	590	——— <i>grandulifera</i>	553
<i>Saurida tumbet</i>	771	——— <i>hernandifolia</i>	553, 554
<i>Sauris decussata</i>	782	——— <i>japonica</i>	553, 555
——— <i>fasciata</i>	783	<i>Clypea wightii</i>	553
——— <i>ignobilis</i>	783	<i>Sterna melanogaster</i>	576
——— <i>usta</i>	783	——— <i>scena</i>	576
<i>Saxicola caprata bicolor</i>	705, 817	<i>Streptopelia chinensis suratensis</i> 513, 818, 819	
——— <i>rossorum</i>	706	——— <i>decaocto decaocto</i> 573, 818, 819	
——— <i>insignis</i>	1002	——— <i>orientalis agricola</i>	819
——— <i>macrorhyncha</i>	706	——— <i>meena</i>	818
——— <i>maura</i>	706	——— <i>senegalensis cambayensis</i> 573,	
——— <i>torquata indica</i>	706, 981	818, 819	
<i>Scotiodon acutus</i>	770	<i>Strix ocellata ocellata</i>	1006
<i>Scolopia crenata</i>	910	<i>Sturnopastor contra contra</i>	819, 1002
<i>Scolopax rusticola rusticola</i>	818	<i>Sturnus humii</i>	704, 993
<i>Scotocerca inquietus striatus</i>	700	——— <i>purpurascens</i>	704
<i>Scotophilus kuhli</i>	593, 608	——— <i>vulgaris dresseri</i>	704
<i>Sciuropterus fimbriatus</i>	603	——— <i>nobilior</i>	704
——— <i>inornatus</i>	603	——— <i>poltaratzskii</i>	704
<i>Sebastonyma dolopia</i>	623	——— <i>porphyronotus</i>	704
<i>Seiaena dussumieri</i>	778	<i>Suya crinigera striatula</i>	709
<i>Seicerens xanthoschistus albosuperci-</i>		<i>Swannia marmarea</i>	784
<i>lians</i>	992	<i>Sylvia althæa</i>	698, 699
<i>Semiothisa ageta</i>	793	——— <i>cinerea</i>	690
		——— <i>communis icterops</i>	698

	PAGE		PAGE
<i>Sylvia curruca affinis</i>	698, 699, 990, 1035	<i>Tephrodornis pondicerianus pallidus</i> ...	703
—mimila	699, 1001	<i>Terpsiphone paradisi leucogaster</i> ...	988
—hortensis crassirostris ...	698, 1035	—paradisi ...	817
—indica	700	<i>Thamnobia fulicata cambayensis</i> ...	710
—nana nana... ..	698	<i>Thecla doni</i>	587
<i>Sylviparus modestus simlaensis</i> ...	979	<i>Therapon jarbua</i>	775
<i>Synegia suffusa</i>	788	<i>Thereiceryx zeylanicus caniceps</i> ...	921
<i>Sypheotides bengalensis</i>	820	<i>Threlicornis melanocephalus mela-</i>	
<i>Syrrhodia lutea</i>	795	nocephalus	820
<i>Sysstema aulotis</i>	944	<i>Thymelicus actaeon</i>	630
—longiplaga	945	—hyrax	630
—pauilla... ..	945	—leonina	630
—semicirculata	945	—lineola	630
<i>Taccocua leschenaulti sirkee</i> ...	881	—nervulata astigmata ...	631
—sirkee sirkee	569	—nervulata	631
<i>Tadorna tadorna</i>	823	—stigma	630
<i>Tanaoctenia haliaria</i>	786	—sylvatica occidentalis ...	631
<i>Taphozous kachhensis</i>	608	—sylvatica	631
—longimanus	1030	—tenebrosa	631
<i>Taractrocera aliena</i>	625	—thaumas	630
—archias quinta	625	<i>Tichodroma muraria</i>	696, 980, 1001
—ardonia	624	<i>Tiliacora acuminata</i>	551
—ceramas atropunctata ...	624	<i>Timanda subobliquaria</i>	781
—ceramas	624	<i>Tinospora cordifolia</i>	549
—nicèvillei	624	—malabarica	549
—oberthuri	624	<i>Torgos calvus</i>	570, 878
—danna	624	<i>Torpedo sinus-persici</i>	770
—dhshalia	625	<i>Tribura major</i>	989
—dischaka	625	<i>Trimeresurus gramineus</i>	566
—flavoides	624	—monticola	566
—lineata	624	<i>Tringa erythropus</i>	575
—maevius flaccus	624	—glareola	575
—maevius	624	—hypoleuca	575
—myconius	625	—ochropus	575
—sagara	624	<i>Trirhinopholis nuchalis</i>	561
—sangira	624	<i>Trochalopterum lineatum griseicentior</i>	979
—silhetica	624	—variegatum simile	979
—sumatrensis	624	<i>Troglodytes t. neglectus</i>	696, 980
—talantus	625	<i>Turdus atrogularis</i>	710, 985
—ziclea samadha	624	—boulboul	1002
—tissara	624	—castaneus castaneus	984
<i>Tasta argozana</i>	784	—merula intermedia	710
<i>Tatera cuvieri</i>	595, 614	—maxima	710, 984
—indica	603, 611	—unicolor	710, 985
<i>Tchitrea paradisi turkestanica</i> ...	705	—viscivorus bonapartei ...	710, 817, 985
<i>Telicota augias augias</i>	629	<i>Turnera ulmifolia</i>	897
—palmarum	628	<i>Turnix javanica taigoor</i>	1006
—prusias insularis	628	—tanki tanki	1006
—pythias bambusae	629	<i>Tursiops catalanta</i>	816
—pythias	629	—dawsoni	816
—upadhana	629	—fergusoni	816

			PAGE				PAGE
<i>Tursiops gilli</i>	816	<i>Virachola smilis nicevillei</i>	589
<i>Typhlops braminus</i>	558	----- <i>smilis</i>	589
----- <i>diardi</i>	558	<i>Viverricula malaccensis</i>	601, 613
<i>Tyto alba javanica</i>	570, 1005	<i>Vulpes bengalensis</i>	602
<i>Udaspes folus</i>	964	----- <i>leucopus</i>	610
<i>Unkana attina</i>	616	----- <i>montana</i>	602, 610
----- <i>harmachis</i>	616	<i>Xantholæma hæmacephala indica</i>	567, 819
----- <i>staudingeri</i>	616	<i>Xenographia adustata</i>	792
<i>Upupa epops epops</i>	568, 879, 1005	<i>Xenopeltis unicolor</i>	559
----- <i>loudoni</i>	879	<i>Xenorhynchus asiaticus asiaticus</i>	576, 820
----- <i>orientalis</i>	568, 1005	<i>Zamenis korros</i>	562
----- <i>subsp</i>	818	<i>Zea mythea</i>	616
<i>Urapteryx ebuleata</i>	783	<i>Zela zenon</i>	615
----- <i>kantalaria</i>	783	----- <i>zeus optimus</i>	615
----- <i>multistrigaria</i>	783	----- <i>zeus</i>	615
<i>Urocissa flavirostris cucullata</i>	817, 977	<i>Zoothera monticola</i>	817
<i>Uroloncha malabarica</i>	862	<i>Zygonyx ida</i>	769
<i>Vanellus vanellus</i>	823	----- <i>ilia</i>	769
<i>Vespertilio leisleri</i>	600	----- <i>iris</i>	741, 763
----- <i>moupinensis</i>	612	----- <i>aenea</i>	766
<i>Vesperugo leisleri</i>	600	----- <i>ceylanica</i>	766
----- <i>oleisleri</i>	600	----- <i>davina</i>	768
<i>Viola cineria</i>	908	----- <i>insignis</i>	766
----- <i>kathiawarensis</i>	908	----- <i>isa</i>	767
----- <i>odorata</i>	908	----- <i>malabarica</i>	763, 768
----- <i>stocksii</i>	908	----- <i>mildredæ</i>	766
----- <i>tricolor</i>	908	----- <i>torridaisis</i>	769
<i>Virachola dohertyi</i>	588				

11752

THE
JOURNAL
OF THE
BOMBAY NATURAL HISTORY SOCIETY.

EDITED BY
SIR REGINALD SPENCE, KT., F.Z.S., and S. H. PRATER, C.M.Z.S.

VOL. XXXI, No. 2.

Date of Publication, 15th August, 1926.

Price to Non-Members ... *Rs. 15-0-0*
or £1-2-6.

For terms of membership see inside front cover.

Honorary Secretary's Address:
BOMBAY NATURAL HISTORY SOCIETY
6, Apollo Street, Bombay.

LONDON AGENTS:
DULAU & Co., Ltd.,
34-36, Margaret Street, Cavendish Square, W.

PRINTED AT THE DIOCESAN PRESS, MADRAS

THE SOCIETY'S PUBLICATIONS

Mesopotamia.

A Survey of the Fauna of Iraq. Mammals, Birds, Reptiles, etc., made by Members of the Mesopotamia Expeditionary Force 'D', 1915-1919. Rs. 7-8

Birds.

A Hand List of Genera and Species of Birds of the Indian Empire by E. C. STUART BAKER, O.B.E., M.B.O.U., F.Z.S., C.F.A.O.U. Rs. 6

(Price to Members Rs. 4-12).

Game Birds of India, Vol. I. (Indian Ducks and their allies) 2nd Edition. Rs. 63

(Price to Members Rs. 50).

Game Birds of India, Vol. II. (Woodcock, Snipe, Bustards, and Sandgrouse) 1st Edition. Rs. 55

(Price to Members Rs. 42).

Set of Game Bird Plates (30), Vol. I. Rs. 20

(Price to Members Rs. 15).

Set of Game Bird Plates (19), Vol. II. Rs. 13-8

(Price to Members Rs. 10).

Indian Pigeons and Doves with coloured illustrations by E. C. STUART BAKER, O.B.E., M.B.O.U., F.Z.S., etc. Rs. 37-12

(Price to Members Rs. 30).

Snakes.

Poisonous Terrestrial Snakes of our British Indian Dominions, 3rd Edition, by COL. F. WALL, C.M.G., I.M.S. Out of print. Orders being registered for new Edition. Rs. 3

Identification of Poisonous Snakes—

Hanging Wall Chart, 30" x 40".

Rs. 5

Folding Pocket Chart.

Rs. 1-12

Snakes of Bombay Island and Sa'sette, by S. H. PRATER, C.M.Z.S. Re. 1

Miscellaneous.

Wood-Destroying White Ants of the Bombay Presidency by Rev. J. ASSMUTH, S.J. Re. 1

Catalogue of the Society's Library. Re. 1

List of Indian Butterflies by LT.-COL. W. H. EVANS, R.E. Rs. 2

Wall Chart to distinguish a Wild Dog from a Domestic Dog or Jackal. Rs. 2-8

TERMS OF MEMBERSHIP.

Life Members pay an entrance fee of Rs. 20 and a Life Membership fee of Rs. 350.

Ordinary Members pay an entrance fee of Rs. 20 and an annual subscription of Rs. 25.

Members elected in November and December are not liable for the subscription for the year in which they are elected.

MEMBERS RESIDING OUTSIDE INDIA.

The terms are the same for members living outside India. Such members should pay their subscriptions by means of orders on their Bankers to pay the amount of the subscription, plus postage—in all Rs. 26-8-0 to the Society in Bombay on the 1st January in each year. If this cannot be done then the sum of £1-15-4 should be paid annually to the Society's London Bankers—The National Bank of India, Bishopsgate Street, London, E.C.

CONTENTS OF Vol. XXXI, No. 2

	PAGE
THE GAME BIRDS OF THE INDIAN EMPIRE. (<i>With a coloured plate</i>). By E. C. Stuart Baker, F.Z.S., F.L.S., M.B.O.U., C.F.A.O.U.	233
SCIENTIFIC RESULTS FROM THE MAMMAL SURVEY No. XLVI. By Helen M. Lindsay, M.A., B.Sc.	239
THE MASON WASP (<i>Eumenes conica</i>). Part I. (<i>With a plate and text figures.</i>) By Major R. W. G. Hingston, I.M.S.	241
NOTES ON SOME NEW AND INTERESTING BUTTERFLIES FROM INDIA AND BURMA. Part I. (<i>With one coloured and one black and white plate.</i>). By Major-General H. C. Tytler, C.B., C.M.G., C.I.E., D.S.O., F.E.S.	248
A CONTRIBUTION TO THE ORNITHOLOGY OF DELHI. Part I. (<i>With two plates.</i>) By the late Mr. S. Basil-Edwardes, M.B.O.U., R.A.O.U.	261
TAKIN SHOOTING IN THE SPRING. (<i>With a plate.</i>) By Capt. E. Maxwell West	274
BIRDS NESTING WITH A CAMERA IN INDIA. Part VI. (<i>With fourteen plates.</i>) By Capt. R. S. P. Bates	277
NOTES ON A COLLECTION OF MOTHS OF THE FAMILY GEOMETRIDÆ MADE BY CAPT. A. E. SWANN. Part II. (<i>With a plate</i>). By Louis B. Prout, F.E.S.	308
THE COMMON BUTTERFLIES OF THE PLAINS OF INDIA. Part XXXVI. By T. R. Bell, C.I.E., I.F.S. (Retd.)	323
THREE MONTHS UP THE VALLEY OF THE SUTLEJ RIVER. Part II. (<i>With two plates.</i>) By Lt.-Col. R. W. BURTON, I.A. (Retd.)	352
ON THE DOWN PLUMAGES OF SOME INDIAN BIRDS. (<i>With a plate</i>). By C. B. Ticehurst, M.A., M.B.O.U. (Late Capt., R.A.M.C.)	368
B.N.H.S. MAMMAL SURVEY OF INDIA, BURMA AND CEYLON. Report Nos. 40, 41 and 42. By Mrs. Helen M. Lindsay, M.A., B.Sc.	379
INDIAN DRAGONFLIES. Part XXIV. (<i>With two plates and five text figures.</i>) By Major F. C. Fraser, I.M.S.	408
THE IDENTIFICATION OF INDIAN BUTTERFLIES. Part X. (<i>With a plate</i>). By Col. W. H. Evans, D.S.O., F.Z.S., F.E.S.	427
ON SOME INTERESTING FEATURES OF THE FAUNA OF THE WESTERN GHATS. By S. L. Hora, D.Sc.	447
RECENT ADDITION TO THE INDO-CEYLONESE COCCID FAUNA, WITH NOTES ON KNOWN AND NEW FORMS. By T. V. Ramakrishna Aiyar	450
A NOTE ON THE BIRDS OF KULU. By H. Whistler, F.L.S., F.Z.S.	458
LADY (ELIZABETH) GWILLIM. By C. A. Wood, M.D., M.B.O.U.	486
SOME NOTES ON THE SECOND EDITION OF THE FAUNA OF BRITISH INDIA —BIRDS. VOLS. I AND II. By C. B. Ticehurst, M.A., M.R.C.S., M.B.O.U., F.R.C.S.	490
REVIEWS:—INDIAN BIRD LIFE. By Douglas Dewar	500
MANUAL OF THE BIRDS OF CEYLON. By W. E. Wait	500
A HANDBOOK OF THE BIRDS OF EASTERN CHINA. By J. D. D. L. A. Touche, Pt. II	501
THE AQUARIUM BOOK. By E. G. Boulenger	502
ACCOUNT OF A PHOTOGRAPHIC EXPEDITION TO THE SOUTHERN GLACIERS OF KANCHENJUNGA IN THE SIKKIM HIMALAYA. By N. A. Tombazi	503
LIFE OF PLANTS. By Sir Frederick Keeble	504
EDITORIAL	506
OBITUARY:—MR. S. BASIL-EDWARDES	508
H. H. THE MAHARAJA OF DHAR	508

	PAGE
MISCELLANEOUS NOTES :—	
I.—Measurements of Game Animals. By Major C. H. Stockley, O.C.F., M.C., D.S.O.	509
II.—Comparative Shape and Measurement of the Fore Feet in Tuskers and Tuskless Elephants. By I. L. Cameron.	512
III.—A Wonderful Sight. By R. C. Morris.	513
IV.—Tigers and Elephants. By R. K. Anderson.	514
V.—The ' Sambhur ' Call of Tigers. By W. R. C. Beadon.	515
VI.—Wild Dogs in Mysore. By Capt. G. E. Windle.	516
VII.—Solitaries. By R. C. Morris.	517
VIII.—A Solitary Cow Gaur. (<i>With a photo.</i>) By Salim A. Ali. ...	518
IX.—A Cross between an Ibex and a Tame Goat. (<i>With a photo.</i>) By Col. A. E. Ward.	519
X.—The Nilgiri Thar (<i>Hemitragus hylocrius</i>). (<i>With a plate</i>). By A. P. Kinloch, F.Z.S., M.B.O.U.	520
XI.—The ' Clicking ' of the Muntjac (<i>M. reevesii</i>). By Max Fre.	521
XII.—Measurements and Weight of a Malay Tapir. By S. H. Prater, C.M.Z.S.	522
XIII.—The Occurrence of the Pink-footed Goose (<i>Anas brachyrhynchos</i>) and the Mallard (<i>Anas boschas</i>) in Jaipur, Rajputana. By Major W. M. Logan Home.	522
XIV.—On the Occurrence of the Pallas' Sandgrouse (<i>Syrhaptes</i> <i>paradoxurus</i>) within Indian limits. (<i>With a plate</i>). By S. H. Prater, C.M.Z.S.	522
XV.—The Occurrence of the Giant Heron (<i>Ardea goliath</i>) in the Khulna District, Bengal. By S. H. Prater, C.M.Z.S.	523
XVI.—The Rock Horned Owl in Kashmir. (<i>With a photo</i>). By B. B. Osmaston, C.I.E., I.F.S. (Retd.)	523
XVII.—Mating Habits of the Common Kite (<i>Milvus migrans gerinda</i>). By Salim A. Ali.	524
XVIII.—Brahminy Kite (<i>Haliastur indus indus</i>) Swimming. By S. H. Prater, C.M.Z.S.	526
XIX.—An Albino Bustard (<i>Eupoditis edwardsi</i>). By Maharaja Kumar Shree Vijayarajji.	526
XX.—The Transport of Birds' Eggs. (<i>With a text figure.</i>) By General R. M. Betham, M.B.O.U.	527
XXI.—Earth Snake (<i>Silybura sp.</i>) and Chicken. By A. P. Kinloch, F.Z.S., M.B.O.U.	528
XXII.—An Interesting Encounter with a Cobra. By W. H. Theobald.	528
XXIII.—The Occurrence of the Butterfly <i>Appias indra</i> var <i>aristoxenus</i> in the Nilgiris. By J. Florence.	529
XXIV.—On <i>Psara phaeopteralis</i> as a Pest on Grasses in South India. (<i>With a plate</i>). By M. C. Cherian, B.sc. and C. J. George, M.A.	529
Proceedings	531



THE INDIAN WATER RAIL

Rallus aquaticus indicus

♂ life size

JOURNAL OF THE Bombay Natural History Society

AUGUST, 1926

VOL. XXXI

No. 2

THE GAME BIRDS OF THE INDIAN EMPIRE

BY

E. C. STUART BAKER, F.Z.S., F.L.S., M.B.O.U., C.F.A.O.U.

VOL. V

THE WADERS AND OTHER SEMI-SPORTING BIRDS

PART I

(*With a coloured Plate*)

I have been asked by the editors of the Bombay Natural History Society to bring out in their journal a series of articles on the waders and other semi-sporting birds of India which will eventually take a place as the final volume of our work on *The Game Birds of the Indian Empire*. Hume and Marshall, in their magnificent work on the game birds, fully realized that every man, when he went out shooting small game, was constantly killing birds which to him seemed to be, to all intents and purposes, game birds, although not coming within the usual acceptance of this term. They accordingly arranged a part volume to include such birds and we propose to do the same.

Possibly there is no country in the world in which the sportsman meets such an extraordinary wealth of birds as he does in India and China coming so near the border line of game birds. Many of these birds are excellent for food and are amongst the wariest and hardest of birds to approach or to bring to bag. Thus anyone who has spent the day after the Demoiselle Crane, whether successful or not, will never grudge this magnificent bird a high place amongst our Indian game birds. Amongst the vast number of birds generally known as Snippets or 'Kuch-nahin' there are

many, such as the Godwit and others, which are almost equally deserving of such a place, for even if not so difficult to approach, they afford good shots on the wing and are not to be despised when on the table. Again, every one who goes Snipe shooting must have once or more shot birds which as they rose, he imagined for the moment to have been Quail but which when picked up proved to be merely one of the small Rails.

Our editors therefore consider that it would be advisable to produce a volume by which the sportsman may be able to identify the hundred and one birds which he may by chance or design or even by mistake, bring to bag in the course of a day's shooting.

Roughly speaking these birds embrace the Cranes and Storks and the Rails and Waders, the latter belonging to the big family *Charadriidæ* which also embrace the really game Plovers, such as the Golden and the Grey.

The number of birds to be dealt with will of course be very large and it is accepted that such a volume will be one required more by the sportsman pure and simple, than by the field naturalist or scientist. It is therefore proposed to eliminate from this volume all synonymy beyond the first reference to the scientific names quoted. Again, in dealing with each bird, each species will be treated as a whole and only a brief reference made to those subspecies or geographical races into which the species may be divided. As regards field notes and, whenever possible, sporting yarns, these will be given in full as also will be breeding notes in so far as these are available.

For illustrations an endeavour will be made to select those species of birds for delineation which most often come in the way of the shooter. For the others, the reader will have to rely on the descriptions given.

RALLUS AQUATICUS

The Water Rail

Three races of the Water Rail have been recorded as occurring in India. These three are

1. *Rallus aquaticus aquaticus*. The Water Rail.

Rallus aquaticus Linn., Syst. Nat., Ed. 10, p. 153, 1758 (Great Britain).

2. *Rallus aquaticus indicus*. The Indian Water Rail.

Rallus indicus Blyth, J. A. S. B., xviii, p. 820, 1840 (Bengal).

3. *Rallus aquaticus korejewi*. The Turkestan Water Rail.

Rallus aquaticus korejewi Sarudny, Orn. Monatsb., p. 209, 1905 (E. Turkestan).

RALLUS AQUATICUS INDICUS

The Indian Water Rail

Description: Forehead, crown and nape black with rufescent olive-brown margins; supercilium from the forehead to above the earcoverts white above the lores, ashy posteriorly; lores and a

line through the eye dark ashy-brown; upper plumage, scapulars, inner secondaries and tail black with broad olive-brown edges; lesser and median wingcoverts olive-brown, with traces of white bars and tips; greater and primary coverts olive-brown, sometimes with faint traces of white bars, sometimes with none at all; quills brown; cheeks and sides of head ashy-grey; chin and throat nearly white; foreneck, breast and abdomen ashy marked with brown; flanks, vent and lower abdomen black or blackish-brown, barred with white; under tail-coverts black edged with rufescent white.

Colours of soft parts: Iris red-brown to brick-red; bill, upper mandible dark brown with a stripe on the base bright orange-red to vermillion; base of lower mandible also red but rather paler, the terminal third grey to dark horny-brown; legs and feet, fleshy brown to pink or brownish-pink.

Measurements: Wing ♂ 120 to 136 mm.; ♀ 110 to 126 mm.; tail 52 to 65 mm.; culmen ♂ about 40 to 43 mm.; ♀ about 36 to 40 mm.; tarsus about 40 to 45 mm. The Turkestan Water Rail is the largest of the three forms, the wing being nearly always over 126 mm., whilst the English Water Rail seldom has it over 130 mm.

Chick in down. Velvety black all over.

Young birds have broader pale rufescent edges to the lower plumage and are much more definitely barred with white on the wing-coverts.

Distribution: Burma, Assam and Eastern Bengal. Nowhere else in India.

RALLUS AQUATICUS KOREJEWI is close to *R. a. aquaticus* but is lighter above and also a paler grey below.

Distribution: Extends to N. W. India, South Sind and to Sehore, Central Provinces. A Sind bird in the British Museum labelled *indicus* is certainly *korejewi*.

RALLUS AQUATICUS AQUATICUS differs from our Indian bird in having the whole of the breast, etc., a much darker, purer grey; the sides of the head are practically all dark-grey and there are only faint indications, if any, of the pale supercilium; the chin is almost as dark a grey as the throat and breast.

Young birds freshly mounted have pale rufescent edges to the feathers of the breast and lower parts but they are never sufficiently conspicuous to make these parts like those of the Indian Water Rail. No specimens have been obtained in India.

In India the true English Water Rail does not occur but we have two geographical representatives which only differ from it as shown above in a very small degree. Of these, *R. a. korejewi* breeds in the North-West Himalayas from the Afghan territory throughout Kashmir to the lakes in West and South-West Tibet. The other is the race *R. a. indicus* which breeds in North-East Siberia and Japan but which finds its way in winter to Burma, Assam and Eastern Bengal but, so far as is known at present, to no other portion of the Indian Empire.

The inclusion of the Common Water Rail within the limits of the Indian Empire rests upon two specimens obtained by Dr. King and Dr. Adams in the Dun, and a single specimen obtained near

Abbotabad, all of which, however, belong to the slightly paler form found in Turkestan. Those birds also which have been found breeding in Kashmir are of the same race and the British Museum does not possess a single specimen which can really be allocated to the European race.

Nidification: The Water Rails breed either in, or very close to, marshes, occasionally, however, in damp meadows and in ditches or similar places at a considerable distance from water. The nests are almost invariably very well concealed, being placed well in amongst herbage of some kind, such as long rank grass, beds of weeds, in amongst long reeds growing in water, or in amongst, or indeed often under, masses of debris which have collected on the sides of ditches or other places. The nest itself is very loosely put together, a bed of dry rushes, aquatic weeds and grass without any true lining, though the inner portion may be made with somewhat softer and more pliant material than the outer. Sometimes nests are a considerable size, measuring as much as a foot across by about six inches deep. The majority, however, are little more than half these dimensions with an internal rough cup of about 5 in. x 3 in. deep. The hen bird, which apparently does most of the sitting, is very hard to catch on the nest, as she sneaks off very quietly on the approach of any person or animal, creeping through the grass and weeds and not rising until some distance from her nest, if at all. In Europe the Water Rail breeds principally in May and early June, and the Japanese bird also breeds principally in those months. The Kashmir bird, however, is rather later, not starting to breed until the end of May whilst a few do not do so until July. The number of eggs laid varies considerably. In Europe seven or eight eggs forms the usual clutch, though I have known as many as twelve and have heard of fourteen. In Eastern Siberia and Japan, seven or eight seems to be the greatest number ever laid, and Allan Owston informed me that he had known as few as three or four eggs incubated. Of Indian records there are but few. Ward obtained it in Kashmir and also in Ladakh. Osmaston has succeeded in taking a certain number of nests in Kashmir and, according to these ornithologists, five to seven seems to be the normal full clutch.

I do not think that the eggs of the various races can be distinguished from one another, but undoubtedly as a whole, eggs laid by the Japanese race are very much richer in colour than are those laid by the two other forms, although individual eggs of any one of these three can be matched with others of the other two.

The ground colour varies from a pale buff to a pale grey-green; a very few eggs have a distinct reddish tinge whilst in a few others it might almost be called a blue-grey. The marking consists primarily of spots and blotches of dark reddish-brown, a few almost purple-black in colour and a few rather washed out and paler reddish in tint. These are sparsely scattered over the whole egg and are sometimes more numerous at the larger end where they may form a very indefinite ring or cap. The secondary, or underlying spots, are rather more numerous and in colour are a pale neutral tint of orange-grey.

Witherby gives the average of 90 eggs of the European race as 35.44×25.75 mm.; maxima, 39×25.5 mm. and 35×27.3 mm.; minima, 31.9×25 mm. and 36.8×24.1 mm.

Habits: The habits of the Water Rails, to whichever race they belong, are much the same. They are all inveterate skulkers and for this reason appear to be a great deal more rare than probably they really are, and it is possible that if one knew where to look for them and how to make them show themselves, they might be found comparatively common in Kashmir, Ladakh and Eastern Tibet. Without dogs, however, it is extremely difficult to flush them. They are certainly most often to be found within a very short distance of some swamp or lake but occasionally they wander away into grass fields and crops at a considerable distance from any piece of water. When disturbed they run at a very great pace, their tail depressed but the head held even lower than the tail. When, however, they think that they are unobserved they strut about very much as a Moorhen does, their head held erect and their tail in the same position. They have a curious little habit when thus moving about, of constantly flicking their tails up and down in little single jerks repeated for two or three minutes. Considering the size of their wings they are quite good fliers as one would indeed suppose them to be considering the distance some of the forms migrate in winter. Thus individuals of the sub-species *R. a. indicus* must in many instances make a return journey from their Northern breeding homes to India and back, of many thousands of miles, crossing great ranges of mountains and wide rivers *en route*. Our Kashmir form *R. a. korejewi* does not seem to be nearly so migratory in its habits as the last bird or as our European bird *R. a. aquaticus*. It is probable that none of the migrating birds travel very great distances in consecutive hours but it has been noticed that both Burmese and Chinese birds sometimes arrive at their destinations in a state of very great exhaustion. Indeed, one of the few birds which I have personally seen in India was so weak that it allowed me to take it up without making any attempt to escape.

The ordinary call note of this little Rail is a softly repeated 'Chip-chip-chip,' but in the breeding season it has a very loud shrill call which Witherby describes as 'a loud, piercing and explosive scream'. It also utters during the breeding season a low grunting note, whilst both birds, whilst at the nest, emit a sound which Mrs. Turner calls a soft purring note. They are excellent swimmers and almost equally good divers but for the most part their food is obtained in the shallow waters where diving is not very necessary. Their food is very varied and consists of any small fish which they can catch, crustacea and mollusca, many insects and larvæ, in addition to all of which they are said to consume a considerable amount of vegetable matter. Seeds of many plants and oak galls have been found in their crops. They feed to a great extent on land as well as in water and on the rare occasion one gets a sight of them, they may be seen feeding indifferently upon either seed or insects as they come across them. A very popular article of diet with these little birds are young

half-grown grasshoppers as well as fully adult ones and I have seen them hopping up into the air to catch these latter as they take to flight when disturbed from the grass.

They are not one of the birds which find their way into the mixed bags so often made in India, especially by those who have recently landed in the country; nor can they claim any status among the Game Birds either from the difficulty they present for a shot or on account of their qualities for the table. At the same time they are birds about which we still have much to learn and concerning which field records are desiderata.

(To be continued)

SCIENTIFIC RESULTS FROM THE MAMMAL SURVEY NO. XLVI

BY

HELEN M. LINDSAY, M.A., B.SC.

FURTHER EXAMINATION OF SINGHALESE SPECIES OF *Funambulus* AND DESCRIPTION OF THE NEW SUB-SPECIES, *F. p. matugamensis*.

From Ceylon the Indian Mammal Survey obtained 152 specimens of *Funambulus palmarum* which were classified by Mr. Wroughton and Miss Ryley. Later, in 1915 Mr. Thomas and Mr. Wroughton published a paper 'On Singhalese species of *Funambulus*' J.B.N.H.S., vol. xxiv, No. 1, in which they give a key to the species and divide *Funambulus palmarum* into five distinct sub-species:—

- | | | | |
|-----|-------------------|-----------------|------------------|
| (1) | <i>Funambulus</i> | <i>palmarum</i> | <i>palmarum</i> |
| (2) | " | " | <i>favonicus</i> |
| (3) | " | " | <i>kelaarti</i> |
| (4) | " | " | <i>brodiei</i> |
| (5) | " | " | <i>olympius</i> |

mainly on considerations of differences in colour and in habitat. They also state that true *Funambulus palmarum* does not occur in Ceylon.

Recently three new sets of material have been received from Ceylon and further investigation has thus been made possible. The Colombo Museum has sent ninety specimens representing all the nine provinces into which Ceylon is divided: Mr. W. W. A. Phillips has presented to the British Museum a set of twenty-one specimens and has also sent another lot of twenty-eight specimens from his own estate, making a total of 139. In the British Museum stock was found an old specimen, presented in 1866 by Viscount Walden and labelled '*S. tristriatus*', which on detailed examination of its skull and skin must now be included in the new collections. In characteristics and habitat these 140 specimens on the whole bear out the division made by Mr. Thomas and Mr. Wroughton, but in addition they also disclose the presence of true *palmarum*, as shown in three specimens obtained on low ground in the east of the North Central Province.

A NEW SUB-SPECIES CALLED *Funambulus palmarum matugamensis*.

These collections show too that 18 specimens from Matugama in the West Province, about 30 miles south-east of Colombo, differ so much in colour and measurement of skin and character of skull that they must form a new sub-species to which I have given the name *Funambulus palmarum matugamensis*. Of these eighteen, Mr. Phillips has sent twelve, while five came from Colombo Museum and the old one from the British Museum collections.

This new sub-species comes closest in size to *F. p. favonicus*. Thus the average measurements of seven adult specimens give

Head and Body: 144'7 (143); Tail: 138 (144); Hind foot: 35 (33); Ear: 17 (16). SKULL:—Greatest length: 38'7 (38); nasals: 11 (12); zygomatic breadth: 21 (22); interorbital breadth: 11 (11). A cleaned skull of this new sub-species shows:—Greatest length 38 (38); condylobasilar length: 32 (31); nasals: 11 (12); zygomatic breadth: 21 (22); basilar length: 30 (30); palatilar length: 16 (16'5); interorbital breadth: 11'5 (11'5); upper tooth row: 7 (7'5).

The figures in brackets are the measurements of the type of *F. p. favonicus*.

But in colour there is much less resemblance. These eighteen specimens are dark as *F. p. olympius*, the saddle being almost black but the three sub-equal dorsal stripes are here all the same colour of tawny or tawny ochraceous, and $\frac{1}{8}$ " to $\frac{1}{4}$ " in breadth. The under parts are ochraceous buff or woodbrown, almost isabelline in some specimens. The tail is somewhat bushy and is orange-rufous below, thus resembling that of *F. p. brodiei*. The neck and shoulders are olive, while the flanks are woodbrown. The face is as a rule coloured like the neck but in winter pelage gets a reddish tinge. The feet are dark as in *F. p. olympius*.

The shape of the skull in this sub-species differs from that of the others in the prominence of the olfactory lobes and the size and shape of the nasals, also in the distance between the point of junction of these lobes and the point of junction of the nasals. In this sub-species the measurement between these two points is 4 mm. while in *F. p. favonicus* it is 5 mm.; in *F. p. olympius* 6 mm. and in *F. p. kelaarti* from 4.5 to 5 mm. Examination of the collection as a whole shows that the shape of the nasals in the different sub-species is fairly constant:—e.g., the posterior edge of the nasals in *F. p. favonicus* is deeply and sharply serrate: that of *F. p. kelaarti* is also pointed but not so deeply serrate and the nasals terminate practically in a line with the frontopremaxillary suture, while that of *F. p. olympius* is of rounded shape almost oval, and ends well in front of the suture.

In classifying these 140 specimens reliance has been placed mainly on the characters of the skull and on the locality whence they were obtained since the variability of colour of coat and size in any one species is considerable, depending on the season and nature of the habitat. This new sub-species comes from a tract of country 100 feet above sea-level in one of the moistest areas of Ceylon with an average rainfall of over 150". Rubber and cocoa palms are grown there and there is much low scrub and jungle around the estates. *F. p. matugamensis* may be found therefore to be a localized form of *F. p. palmarum* just as *F. p. olympius* has been confined to the high hills, and *F. p. favonicus* to the submontane regions.

The classification of these specimens resulted in finding *F. p. olympius* 48; *F. p. favonicus* 28; *F. p. kelaarti* 22; *F. p. brodiei* 10; *F. p. matugamensis* 18. The remaining specimens could not be determined as skull or labels were amissing. There were 3 albino juvenile specimens, one of which had a skull resembling *F. p. favonicus* but as there was no label it cannot be named.

THE MASON WASP (*EUMENES CONICA*)

BY

MAJOR R. W. G. HINGSTON, I.M.S.

PART I

ARCHITECTURE

(*With a plate and text figures*)

This is the narrative of a mason wasp, the common species *Eumenes conica*, which I shall speak of as the cone-shaped *Eumenes*.

This wasp enjoys a wide distribution. It is found in all parts of the Indian Peninsula, crosses over into Ceylon, spreads east to the Burmese plains. From there it extends north to Chinese territory and south into the regions of Malay.

We are concerned with the female only. She is the architect. Her mate in comparison is puny and insignificant. He does no work. His business is to frolic in the tropical sun, to follow the mere dalliance of life. Certainly he performs the bare functions of his sex; but everything concerned with the architectural instinct, the skill, the industry, the intelligent toil, are the attributes of the female alone.

First a note on her appearance. Strong and powerful, a triangular head with incurved jaws, a square-shaped thorax, a thread-like waist, an abdomen swollen into an elongated pear: these are her most striking points. Dark-red in colour. On the front of her head is a patch of yellow, a few black bars cross her abdomen, a delicate purplish gloss tips her golden wings. She is smooth and clean and shining without a trace of any fur.



Eumenes conica



NOCTUID LARVA EXTRACTED
FROM A MUD CELL OF *Eumenes*
conica

In character bold and intrusive, she enters courageously into any apartment likely to supply her needs. Associated with this daring is some suspicion. If aware of any prying interference, she quickly deserts her home. Her habits are solitary in the

strictest sense. She is powerfully armed with sting and poison, not, however, supplied for protection, but in order to paralyze her prey.

I pass to architectural operations.

She constructs a cell composed of mud, building it in the shape of a dome. At the summit she leaves a circular aperture. This serves the purpose of a door. When the cell is built and the door still open she introduces an egg. Then she goes off to search for caterpillars. These she paralyzes, then carries to the edifice, then pushes through the door. This is the provender necessary for her larva. She brings sufficient to fill the compartment, then closes the aperture with clay. The first cell is now complete; but no sooner has she finished one than she commences to build the next. Each cell is placed so as to touch an adjoining cell; consequently there results a cluster of neat little oval domes. She then covers the whole with mud until it looks like a shapeless cake. There are thus five architectural stages. First the raising of the dome, second the laying of the egg, third the introduction of the provender, fourth the closing of the gate. These four successive steps enter the construction of each separate cell. The fifth stage concerns the cell combination. It is the final cover which the wasp applies over the whole.

This is the briefest outline of her operations. I come now to some detailed points.

Work begins in the middle of April, at least in this district of Fyzabad. A preliminary investigation marks its commencement. Exploring verandahs, entering rooms, the wasp examines each shady corner in her efforts to find an acceptable site. Particular attention is given to woodwork. Every object receives minute inspection. Only after prolonged and careful investigation will she content herself with some special place.

She shows definite preference for a wooden foundation, and thus frequently selects the furniture of a room. I have seen her build on a flat table, on the vertical sides of wooden boxes, on the leg of an easy chair. Other materials are not neglected. One wasp chose an iron hat-box, a second the surface of a glass pane, a third a leather trunk. She seems quite indifferent to direction in space. The nest on the table lay in the horizontal, that on the glass in the perpendicular direction, the one on the hat-box had curved foundations to suit the convex base. More attention is given to conditions of light. If possible she avoids bright surroundings, usually choosing some secluded corner liberally supplied with shade. For this reason, I think, she avoids white plaster. Her dislike is for the glaring colour of the lime-wash, not because she thinks the foundations insecure.

The site, being chosen, is minutely examined. It must be prepared for the reception of mud. If an even surface, no attention is necessary; but if of a rough or fibrous texture, we observe that she tears away the shreds, diligently employing every effort to make the area smooth. This is but a preliminary act. Nevertheless it has an end in view. This surface will later become the floor of her cell. Within the chamber her grub will develop. Its body

will come in contact with the floor. Hence there must be no roughness or irregularity, nothing must be left that might injure the larva which is the object of all her toil.

Attend to her workmanship.

Here she comes. A vibrating hum denotes her arrival. In her jaws is a pellet of mud. First there is an aimless circle round the room; then, hesitating as though suspicious, she alights at the chosen spot. An antennary exploration is her next act. Apparently everything is satisfactory for she immediately proceeds to build. Carefully she lays her pellet on the site, then shapes it with her jaws into a low ridge about half an inch in length. On one side the ridge is vertical, on the other it shelves into a gentle slope towards what will later be the interior of the cell. This is the first stone in her edifice, a semi-liquid mud which very quickly dries into a solid brick. It takes her a minute to construct this ridge. When finished, she goes off. There is no delay. A second pellet is brought, applied to the first, spread out in the same manner so as to both lengthen and raise the ridge. More pellets follow. The ridge is extended, shaped into a curve, then bent round into an oval wall which becomes the substructure of the cell.

Architecture becomes now a systematic business. We watch her methodically coming and going. There is no haste, no hesitation. She works in an almost mechanical way. Often she flies direct to her edifice. At other times she alights a little distance away and from there goes on foot to her work. Watch her method. A position is taken on the wall. Her antennae are thrust into the oval where they investigate each step in the work. With hind and middle legs she holds the masonry, then begins to build. Her head is within, her abdomen curled round outside; she looks as if hanging to the edge of the wall. Her whole body is acutely bent; her wings, folded, erect and motionless, are kept well out of harm's way.

Now see the application of mud. With her fore tarsi she supports the pellet while her other legs grip the wall. Her mandibles knead out the mortar. They both spread it sideways into a layer and rake it up into an edge. Her antennae at the same time open and close, sensitive fingers measuring the ground and feeling the pellet being moulded into place. What a display of neatness and expedition! One or two minutes fixes the brick. A wait follows for perhaps a second, just a swift momentary survey. Then satisfied, she takes to her wings, disappearing with a noisy buzz.

She stays away a variable time. Her destination is the clayey patch where brick after brick is made. If the material is close at hand she is back in a minute; otherwise she may be absent more than twice that time. Occasionally she makes a prolonged interruption; I do not see her resume architecture until an hour has elapsed. The clay which she brings is of the finest material. It has been carefully selected and mixed with her saliva. Every particle of grit has been removed. Just sufficient moisture has been added to keep it in the spherical shape. Some species of


Eumenes insert pebbles in their masonry. The object of course is to strengthen the wall. But not so with the cone-shaped mason. She will have nothing but clay. Must she not have trouble in carrying this burden, a spherical globule of sticky mud about the size of an ordinary pea? Certainly if her jaws had not been specially adapted. But observe their suitability. They are long and curved and fit neatly round the pellet so as to enclose it in a ring.

I never caught her gathering clay. But her method must be the same as that of *E. dimidiatipennis*, an allied species that makes similar domes. From this latter I learn the necessary facts. These masons do not use ordinary mud; it is from the hardest and stoniest of available material that they make their architectural bricks. They seek it from the well-rolled thoroughfare, the solid road composed of limestone crushed and firmly driven in. There she fashions the viscid pellet, no doubt preferring powdery material because it can be made into more rigid clay. I see her alight on the open roadway. She digs into it, breaks it into dust, excavates a shallow depression into which she thrusts her head. In this hollow the globule is shaped. Her jaws gather up the finest dust; they thrust aside every granule except those of almost impalpable size. Her salivary glands pour out secretion. With this the dust is thoroughly impregnated until brought to the consistency of paste. Her jaws at the same time make it spherical; her antennae applied on either side examine and test its shape. A minute is sufficient for making the pellet. Then bringing forward her fore tarsi to support it, away she goes for the nest. Often she makes a circle or two around the area. This is a common habit of solitary wasps, a topographical performance, a visualization, a fixing of the spot in their memories with the intention of recognizing it again.

I return to the masonry. Pellet after pellet continues to arrive. For a quarter of an inch the wall is almost vertical. After this it begins to incline inward. The dome is coming into shape. She now builds so that each addition diminishes the oval. The wall curves in on every side. At length it appears to be complete except that it has a central hole. It is in fact a perforated dome.

Now follows an interesting performance. The central perforation is a true circle, in diameter one-quarter of an inch. Its edge is even. It is just a plain perforation in a cell.

This however will not suit the mason. She is not content with an ordinary hole. The time will come when she must plug this aperture. She must therefore have some special device in order to hold the stopper in place. For this purpose she constructs an everted rim all round the edge of the hole. Her last visit is devoted to this purpose. Again we observe her usual skill. Her tarsi hold her tightly to the turret as she lays this last pellet around the hole. She rotates upon the cupola, all the while drawing out her clay and smearing it along the edge. At the same time she neatly bends it outward so as to form an everted lip. What is the result? The opening is no longer a plain hole; rather it is shaped in the form of a funnel owing to the everted edge. What



1



2



3



4

THE MASON WASP

1. Completed nest of *Eumenes conica* with apertures of exit made by young wasps.
2. Cells of *Eumenes conica*.
3. *Eumenes* engaged at building her wall.
4. *Eumenes* introducing her eggs.

suitable preparation for a later stage! It will be an easy matter to close the cell by just thrusting a pellet into the funnel, certainly a simpler mechanical operation than securing a plain hole.

The cell is finished. An inspection follows, the wasp thrusting her antennae into the pot. She soon withdraws them, sits for a little while on the summit, but does not attempt to leave. Perhaps she is about to enjoy a rest. She deserves a respite after all this work.

But no. The next step comes quickly, a fascinating incident. Suddenly a deep absorption overcomes her. She seems as if struck into a trance. Slowly she creeps forward to the summit of her dome, advances so as to sit across it with legs spread out on either side. Gently her abdomen goes into the aperture; her legs clutch the everted rim. Her head bows, her antennae droop, her graceful wings fall down along her back; there she clings, rigid as a statue, half without and half within her cell. Why is she thus mesmerized? What absorbs her? There is only one meaning for such behaviour. She is fulfilling the purpose of her labour, introducing an egg into the dome. Not a motion, not a sound accompanies the act. Even the antennae, so seldom motionless, now fall into a passive state. If we look closely to the base of the abdomen we may note a faint convulsive thrill. This is the effort at egg-expulsion, her only indication of life. For one or two minutes she remains thus fixed, immersed in the glow of that supreme ecstasy which accompanies the fulfilment of these vital acts.

The egg is laid. We can tell from outside that the act is over. For restlessness returns after its expulsion. The antennae begin gently to quiver before she withdraws her abdomen from the cell. Sometimes her head now enters the aperture. Perhaps she is anxious to see her egg. It may not be properly attached. At other times, however, she shows no curiosity. Her abdomen is withdrawn. Away she goes without a trace of any desire to inspect this vital object of her toil.

She does not waste a moment. Provisions must be obtained quickly. Is not her egg exposed to danger owing to the open gate? Away she goes amongst the gardens and trees. Untiringly she pursues her deadly quest. Her motion is a swinging and hovering in the air with the utterance of a gentle hum. Now she sways from point to point, now she poises for a moment to pry into some likely nook, then she sweeps off to some more distant field or alights to pursue her search upon the ground. The period of absence will depend on her success. At length she returns. Suspended beneath her is a large caterpillar. It is almost as much as she can manage since it has to be carried through the air. How well she supports it! Her jaws fix it round the neck; her legs are wrapped about its body, they also gather up its tail so that nothing projects behind.

Her next business is to stuff it in the cell. She thrusts its head first through the gate, then, seizing it further back, pushes it in a little more. Eight or ten similar pushes follow until the whole caterpillar is inside. When the caterpillar's tail disappears, the wasp presses it down with her jaws. Then another inspection;

again the antennae enter the aperture. How often have we had to describe this action; how clearly do we realize the antennary function of determining if all is well? The wasp, if dissatisfied with the manner of storage, again pushes in her head. Another round of adjustment follows, more pushing, more pressing, more testing with the tips of the antennae until the mason is finally satisfied that her provender is safely and economically housed.

More caterpillars are then brought. If of large size, four or five are sufficient. If small, she may require nine or more in order to fill the cell. She is independent of any special number. Capture after capture is pushed in until the edifice is crammed to the mouth.

The next business is to close the door. No difficulty is experienced here. She has already made it conical by constructing a lip. All that she need do now is to push a plug into the cone. She, therefore, brings a pellet, thrusts it in. As a rule one is sufficient. If not, she reinforces it by the addition of one or two more.

Eumenes amedei, as told by M. Fabre, makes use of a pebble for this purpose. She employs a stronger, a more specialized plug. Such is not the method of the cone-shaped mason. It is clearly impossible in this alluvial district. There is not a pebble for hundreds of miles in the plain of hardened mud. Animals must adapt themselves to the environment that offers. A pebble may be a much better kind of stopper, but this mason must build with what the country gives her. She can use but a plug of clay.

The work is finished. Let us take a glance at the completed pot. What a graceful tabernacle, oval in outline, built in the design of a dome! Scarcely an inch in the longest diameter; its width is about five-eighths and its height three-eighths of an inch. The interior is even and smooth. This is the surface of chief importance as against it the body of the larva will rest. Hence the mason has paid it minute attention. Her antennae have kept it under examination at all stages in the work. The external surface is comparatively unimportant; there the ridges and furrows are apparent which mark the superposition of the layers. The whole edifice is neatly formed, a charming example of earthen architecture, a perfect model of the potter's art.

The completion of one cell does not terminate her work. She must provide for many ova. Hence as soon as one cell is finished she immediately commences the next. The second is constructed alongside the first. Others are subsequently added, until about seven or eight are linked together in a group. The number of cells is very variable. Sometimes there may be ten in a cluster; at other times we find only two or three. Occasionally they may be placed in line. More often they seem to have no definite arrangement, being piled into an irregular mass.

I come to the last stage in construction, the cover of mud, the shield with which she envelops the whole. She does not wait till all chambers are built before commencing the manufacture of this shield. Frequently a few smears of mud are applied to it

during the intervals between building cells. It is not made on an elaborate plan. Thick where needed to fill depressions, thin where the underlying architecture is raised, its purpose is just to even off irregularities and cover the surface in a uniform layer. Furthermore it adds strength to the edifice, also opposes an additional barrier against the invasion of parasitic foes. Unfortunately it destroys geometrical symmetry. The oval walls, the dome-shaped roofs, the varied points in geometrical architecture are all permanently concealed from view beneath a formless lump.

What a change! The nest no longer possesses attraction! It is now a mere flattened cake of mud that might have been just thoughtlessly thrown against the wall.

(To be continued.)

NOTES ON SOME NEW AND INTERESTING BUTTERFLIES FROM INDIA AND BURMA

BY

MAJOR-GENERAL H. C. TYTLER, C.B., C.M.G., C.I.E., D.S.O., F.E.S.

PART I

(*With one coloured and one black and white plate*)

INTRODUCTION

The butterflies from the extreme North-West of India are but little known ; Chitral has been worked by several collectors fairly thoroughly, and results have been published by Capt. (now Colonel) W. H. Evans, but the adjoining territories of Gilgit, Hunza-Nagar, Astor and Chilas, which are included in the Gilgit agency, and which are a mass of high and inaccessible mountains, have been practically untouched and very little is known of the butterflies that inhabit them.

In 1920 I proceeded on a shooting trip to Astor and availed myself of the opportunity to collect what butterflies I could in the short time at my disposal.

Through the kindness of Lieut.-Col. D. L. Lorimer, C.I.E., late Political Agent of Gilgit, I have received numerous specimens of butterflies collected by himself and by the local inhabitants in out of the way and inaccessible places of the Gilgit Agency, and but for his kindness and the trouble he has taken on my behalf, the material which has made it possible for me to write much of these notes would never have been obtained.

In 1924 I was transferred to Burma and did a certain amount of collecting at Maymyo and the Southern Shan States during the year I was there. Burma is most interesting entomologically and much work still remains to be done. Certain rich localities such as Thandaung are well known and have been worked thoroughly, but speaking generally most places have been but partially worked and some places on the Chinese and Siam borders are practically untouched, and other places, such as Putao on the extreme N.-E. of Burma, are quite unknown entomologically and a rich harvest awaits the first collector who is able to collect in Putao, the only part of Burma where snow-clad mountains occur to any extent.

At Maymyo I was fortunate to meet Mr. G. Cooper of the Survey of India who has done such good pioneer work in collecting in out of the way places and whose fine collection I was privileged to see. I am also indebted to him for the gift of many rare and interesting specimens.

In the spring of 1925 I proceeded to England on leave and was able to work out at the British Museum, South Kensington, the material I had accumulated and write my conclusions in these notes.

I take this opportunity to thank Capt. Riley of the British Museum for his invariable courtesy and ever ready help afforded me in working out my material but for which my task would have been rendered much more difficult.

1. *Troides ferrari*, n. sp.

The female differs from all other females of the yellow forms of *Troides* in the British Museum in the *different shape of the yellow patch* in the cell of the hind wing which is *oval* and extends from the apex to about $\frac{2}{3}$ the length of the cell but does not touch and is well clear of the upper-discocellular. The yellow spot in interspace 6 is very small and narrow and adjoins v. 6 but not v. 7 ; the yellow spot in interspace 1 is as long as spot in cell ; the yellow spots in inter-

spaces 2, 3, 4 and 5 are less than half the distance from the discocellulars to the outer margin; these are followed by large black spots entirely filling the interspaces and by small streaks on either side of the veins. Forewing black with veins very distinctly edged with grey.

Expanse: ♀ 174 mm.

Three females were caught by Lieut.-Col. Ferrar, C.I.E., Chief Commissioner of the Andamans, in the Great Nicobar Islands. It appears to be very close to *T. darsius*, ♀, from Ceylon but until the male is discovered its affinities cannot be determined.

The type is in my collection.

2. *Papilio noblei haynei*, sub-sp. n.

Bingham in his description of *P. noblei* says, 'Forewing with or without a quadrate white spot in the middle of interspaces 1 (a) and 1"', and Frukstorfer in Seitz's work only mentions a form of *noblei* which has a white patch on the hind margin of the forewing and gives the distribution of *noblei* from Upper Burma to Tonkin. I do not know whether de Nicéville's type has a white spot on the forewing as I have not seen his original description and figure but had his type not had a white spot on the forewing I think Frukstorfer would have noticed it and given another name for the form with a white spot. I am unaware whether specimens with or without the white spot on the forewing occur together or not but a solitary specimen in the British Museum taken by Colonel Broughton at Ye' pank-kan, Southern Shan States, West of Pyinmana and a series from Tonkin in the Adam's collection all have the white spot on the forewing upper and underside; one specimen from Central Tonkin in my own collection in addition to the white spot on the forewing also has three yellow lunules on the hind wing. A series of *P. noblei* taken by Dr. Hayne near Myitkyina, North Burma, does not possess the white spot on the forewing either above or below and it would appear that in North Burma a race has developed without a white spot on forewing whilst further south a form with the white spot occurs. I propose calling the form from Myitkyina as above after Dr. Hayne who very kindly gave me two males from his series.

Expanse.—♂ 102 mm.

The type is in my collection.

3. *Paranticopsis xenocles phrontis*, de N.

A very curious melanitic ♂ of this species was taken by Mr. E. Ollenbach's collector in Sikkim. It is much darker than any ♀ I have ever seen.

The specimen is in my collection.

4. *Parnassius dongalaica*, sp. n. Pl. IV, Fig. 5, ♀

A single ♀ which may be a race of *P. mercurius*, Gr-Gr. from Amdo, N.E. Tibet was taken on the Donga-La, on the road to Mount Everest at 16,000'.

It differs from typical ♀♀ of *P. mercurius* in the British Museum collection in being much smaller and paler, and in having a well-marked series of white marginal and sub-marginal spots on both wings; at the base of the hindwing there is a large crimson spot which is wanting in all the ♂ and ♀ specimens in the British Museum.

In appearance it is somewhat like *P. epaphus everesti* Riley, but the keeled pouch at once separates it from that form and places it in the *Discobulus*—*Jacquemontii* *Mercurius* group.

The type is in my collection.

5. *Parnassius discobulus*, Alph :

The *discobulus*—*jacquemontii* forms occurring in Chitral and Gilgit are somewhat confusing and I find it difficult to place some of the forms under their correct names.

In Chitral and Gilgit there are five forms in which the females have keeled pouches; two of these fly together on the Shandur Pass; one of these is a large insect, 78 mm. in expanse with large red discal spots but no red spot at tornus, which I take to be *P. discobulus* Alph: or a race of it; and the other a much smaller and whiter insect, 60-64 mm. in expanse, in which the females have

red tornal spots; I take this to be a race of *P. jacquemontii*. It is not *P. jacquemontii chitralensis* which is a different looking insect and flies much further south—in Shiski Kuh and Jhela Drosh—and has *all* the red spots very bright. I propose naming the Shandun-form of *P. jacquemontii*, *P. j. shandura* sub-sp. n.

Further North and in the extreme N. E. corner of Chitral on the Baroghil Pass leading into Wakhan, Afghanistan, another form of *P. discobulus* occurs which I propose calling *P. d. baroghila* sub-sp. n.

In Hunza the most easterly province of the Gilgit agency yet another *discobulus* form occurs which I propose calling *P. j. hunzaica*, sub-sp. n.

6. *P. jacquemontii shandura*, sub-sp. nov. Pl. IV, Fig. 3 ♂, 4 ♀:

This is the smallest and palest form of *P. jacquemontii* in the North-West of India; the males are very white looking and both sexes either entirely lack or have very little dusky powdering on the wings. The females have red tornal spots on both sides of the hind wing.

Expanse: ♂, ♀, 60-64 mm.

Described from three males in the British Museum and two males and three females in my collection all from the Shandur Pass, Chitral.

The types are in my collection and para types are in the British Museum.

7. *P. discobulus baroghila*, sub-sp. nov. Pl. IV, Fig. 1 ♀

This form is smaller than *P. discobulus* from the Shandun Pass and not so dark. The females have no red tornal spots on the hind wing and agree with *P. discobulus* from the Shandun Pass in this respect.

Expanse: ♂ 66-70; ♀ 64-70 mm.

A very fine series of this insect was obtained by Lieut.-Col. D. L. Lorimer, late Political Agent, Gilgit, on the Baroghila Pass at the extreme north-east corner of Chitral.

The types are in my collection and para types are in the British Museum.

8. *Parnassius discobulus hunzaica*, sub-sp. nov. Pl. IV, Fig. 2, ♀

This form is very like *P. d. baroghila* but averages smaller and is much darker especially in the females which are as dark as the females of *P. discobulus* from the Shandun Pass. It is very near if not identical to an unnamed form in the British Museum from the Hindu Kush and which is the *rhodius* of other authors, not *rhodius* of Honrath. The type of *P. rhodius* Honr, a male is in the Adam's collection in the British Museum and is according to Mr. Riley an *epaphus* form and therefore the *jacquemontii* forms which have been known by the name of *rhodius* require a new name and I propose the above for this form.

Expanse: ♂ 64, ♀ 64-70.

Numerous specimens of both sexes were received from Misgar in Hunza, taken in August; it appears to be very common.

9. *Parnassius delphiuss stoliczkana*, Fd.

A female taken south of Leh in Ladak agrees well with the figure of the type specimen and the figure given by Moore.

There are no specimens of true *P. stoliczkana* in the British Museum but there were two specimens over the label of this name which have nothing to do with this form, one being a specimen of my *P. delphiuss rileyi* from the Rupal Valley in Astor and the other of doubtful locality.

10. *Parnassius delphiuss chitralica*, Verity

P. delphiuss chitralica from Chitral is very similar to *P. delphiuss hunza* Gr-Gr., the type of which is from the Hindu Kush probably north of Hunza in Eastern Gilgit but rather smaller and darker.

Two specimens were received from Yasin, Western Gilgit, on the borders of Chitral which are similar to typical *chitralica*. Three specimens from Misgar, Hunza are like *chitralica* but larger and are darker than *hunza* but approach the latter in size.

11. *Parnassius delphioides rileyi*, sub-sp. nov. Pl. IV, Fig. 6 ♂, 7 ♀

Male and female very near to *P. nicevillei* from the Burzil Pass, Gurais, which again is very close to *P. atkinsoni* from the Pir Panjal and from Haramosh Mountain in Gilgit; it differs from the latter in the smaller marginal spots of the hind wing; these spots are about the same size in *rileyi* as in *nicevillei*, the former, i.e., *P. rileyi*, however differs from *P. atkinsoni* in not having the red spot in interspace 2 of the hind wing nearly so well developed; in both *P. nicevillei* and in *P. atkinsoni* this spot is large and bright; in some specimens of *P. rileyi* this spot is very small and inconspicuous and in one it is completely absent and is replaced by a very large black spot.

The types are in my collection and came from Rupal Valley, Astor and the Farsat Pass leading into Chilas. Para types are in the British Museum. The Farsat Pass is on a continuation of the Range of Mountains on the south side of the Rupal Valley.

The insect is not rare and a good series was taken in July and August. I did not meet with *P. nicevillei* when crossing the Burzil Pass in July.

12. *Parnassius delphioides zogilaica*, sub-sp. nov. Pl. IV, Fig. 5, ♀

A single ♀ from near the Zogila Pass, N. Kashmir, differs from its allies *P. nicevillei* and *P. rileyi* in being much paler on both wings and the marginal spots are rather smaller and largely centred with blue; these spots do not touch one another and are placed on a white ground with no dusky scales between them; in *nicevillei*, *rileyi* and *atkinsoni* the ground colour between these spots is more or less covered with dusky scales.

The specimen obtained agrees well with the description by Avinoff, in the Transactions of the Entomological Society of London, 1915, p. 354, of a male sent to him from Zogila.

♀ Type is in my own collection.

13. *Parnassius delphioides atkinsoni*, M.

Type from Pir Panjal, Kashmir.

P. delphioides nicevillei Avinoff, the type of which is from the Burzil Pass between Gurais and Astor in Kashmir, only differs from *P. d. atkinsoni* in having the marginal spots on the hindwing rather smaller.

Major Stockley took four specimens of a form of *P. delphioides* on the slopes of Haramosh Mountain in Eastern Gilgit which appear to be *P. d. atkinsoni* and are indistinguishable from specimens of that form from the Pir Panjal. This is curious as these localities are widely separated and two forms at least—e.g. *P. d. zogilaica* and *P. d. nicevillei* occur between them.

14. *Parnassius delphioides florenciæ*, sub-sp. nov., Pl. IV, Fig. 9, ♂

Many years ago I sent a *delphioides* form to the late Col. Bingham for identification and he informed me that it was new and named it *P. florenciæ* at my request. I have been unable to trace the publication of the description of this specimen and it is possible he never published it at all, so it may be advisable to describe it again.

Male: Upperside. Rather similar to *P. delphioides stoliczkanæ* Felder but with the wings still whiter and with no dusky scaling in interspaces 1, 2 and 3 of the forewing except a very small and clear dark spot in interspace 1 and with no dusky scaling on the hind wing; marginal area of forewing very dark and broad bearing a row of small and clear white spots not so disjointed in interspace 4 as in *P. d. stoliczkanæ*. Hind wing marginal area completely dusky with three complete dark spots bearing bluish centres and two small dark spots without blue scales; spots in 2, 3 and 4 are in a line and not as in *P. d. stoliczkanæ* in which 3 is inwardly thrown out of line; a clear small red spot on the disc and a very small one at the tornal angle.

Underside: Similar to upperside.

A single male was taken in June 1906 near Phup, Hundes, North of Tehri Garhwal, at 15,000 feet.

15. *Parnassius simo lorimeri*, sub-sp. n., Pl. IV, Fig. 10, ♂

Male and female very like *P. boedromus*, *f. hohlbecki* Avinoff from the Alexandra Mts., Central Asia which may be a form or race of *P. simo* but which Avinoff has treated as a distinct species. The sexes are exactly alike and differ from the form of *P. boedromus* above mentioned in being somewhat paler, the dark spots in cell of forewing not so well developed and in the greater development of the black spot in interspace 5 of the hindwing which is as well developed as the spot in interspace 7.

I have much pleasure in naming this butterfly after Lt.-Col. Lorimer, C.I.E., late Political Agent, Gilgit, who captured a pair on the Kine-Chish Pass, south-west of the town of Gilgit, at about 14,000'. It appears to be very rare and the two specimens captured were the only ones seen; they were flying about at the extreme summit of the Pass which leads into Tribal territory.

16. *Aporia nabellica ferrari*, sub-sp. nov.

The male differs from typical *nabellica* from Kashmir in being much paler than even the females of that form.

The females are very white with black markings and grey powdering on the disc.

Expanse: ♂ 52–60 mm., ♀ 60–62 mm.

Habitat.—Dalhousie, Dugi Pass and Valossa. Several specimens were taken by Colonel Ferrar at Dalhousie in the Western Himalayas and there are four males and four females in the British Museum from the Dugi Pass and Valossa and one ♂ without locality which belong to this race.

17. *Gonepteryx rhamni gilgitica*, sub-sp. n.

Male: Similar to *G. rhamni nepalensis*.

Female. Differs from the ♀ of *nepalensis* in having the extreme base of the costa and the apex of the forewing distinctly tinged with ochreous yellow and the dorsal area and terminal margin of the hindwing as far as v. 4 also tinged with the same colour.

Habitat: Gilgit; Astor; Chilas. A large series of both sexes were received from the above mentioned places.

The types are in my collection and para types are in the British Museum.

18. *Gonepteryx rhamni burmensis*, sub-sp. n.

Male: *Upperside* very similar to *G. rhamni major*, Oberthur, from western China but smaller and orange spots much smaller. The yellow on the forewing is distinctly darker than the hindwing.

Female: *Upperside* both wings of the same colour as the hindwing of the male and similar to the ♀ of *major* named by Verity ♀ *ab. mascula*. In *major* the typical ♀ is cream colour, a yellow form *mascula* Verity occurring occasionally; in *burmensis* the typical ♀ form is yellow, I have not seen any cream coloured females.

Expanse: ♂ 62 mm., ♀ 66 mm.

Habitat: Southern Shan States.

The types are in my collection and para types are in the British Museum.

19. *Colias wiskotti chrysoptera*, Gr.-Gr.

A male and three females were obtained from the Baroghil Pass Chitral, Yarkhun, Ghizer and Gilgit. The females vary greatly *inter se* :—

One is orange-yellow.


One lemon-yellow.

One white approaching ♀ *ab. leachi* Gr.-Gr., but with the dorsal margin upper hind wing distinctly tinged with yellow.

20. *Colias marcopolo*, Grm.-Grsh.

A male taken in the Thui Nala, Yasin Gilgit, in September agrees well with specimens in the British Museum from Turkestan and the Hindu Kush but the inner margin to the black border of the forewing is *conspicuously* indented and forms a continuous row of luneles.

This species has not previously been recorded from within Indian limits.



SOME NEW AND INTERESTING BUTTERFLIES FROM INDIA AND BURMA.

SOME NEW AND INTERESTING BUTTERFLIES FROM INDIA AND BURMA.

EXPLANATION OF PLATE IV.

- Fig. 1. *Parnassius discobolus baroghila*, ssp. n., ♂
 „ 2. „ „ *hunzaica*, ssp. n., ♀.
 „ 3. „ *jacquemontii shandura*, ssp. n., ♂.
 „ 4. „ „ „ „ ♀.
 „ 5. „ *dongalaica*, ssp. n., ♀.
 „ 6. „ *delphius rileyi*, ssp. n., ♂.
 „ 7. „ „ „ „ ♀.
 „ 8. „ „ *zogilaica*, ssp. n., ♀.
 „ 9. „ „ *florenciæ*, ssp. n., ♂.
 „ 10. „ *simo lorimeri*, ssp. n., ♀.

21. *Colias cocandica hindnencica*, Verity

Four males and three females of a form hardly distinguishable from specimens of *C. hinducucica*, Verity from the Hindu Kush Mountains in the British Museum were received from Misgar, Hunza Gilgit.

This species has not previously been recorded from within Indian limits.

22. *Oeneis garhwalica*, sp. nov. Pl. IV, Fig. 4, ♀

Male: *Upperside* pale yellowish brown, very much the same colour as *Paroeneis pumilus* Felder.

Forewing: basal area and cell brownish and a broad brown marginal area; a row of dark spots in interspaces 2, 3 and 5; *hindwing*: a broad brown marginal area as in the forewing; dark markings on underside showing through on both wings.

Underside: *Forewing* a brown streak across the upper portion of the cell, another closing it and an irregular brown line just beyond it from costa to the dorsum; a broad brown marginal band; *hindwing* pale lilac speckled with brown; a broad dark band starting at the costa bent at right angles at its middle and reaching the middle of the dorsum; a broad marginal band; veins pale lilac.

Female: as in male but paler.

Three males and a female were taken at Shiruans, Garhwal, at 13,500' in July.

The types are in my collection and a para type of the male is in the British Museum.

23. *Pararge menava moeroides*, Fd.

A male and two females from Ladak appear to belong to this form; the male has a distinct small yellow patch on the forewing as in *moerula* and *schakra* but much smaller and not so bright. The females have a large yellow patch as in *moerula* and *schakra* but brighter.

Moeroides is usually sunk as a synonym of *menava* but it is certainly a very different looking insect and appears to be a distinct race of that species.

24. *Anlocera swaha gilgitica*, sub-sp. nov.

Male and female differ from typical *swaha* on the upperside in having the bands pure white instead of cream colour on the forewing.

On the underside both wings are very pale grey with the markings on the termen of the hindwing very obscure; bands pure white.

Col. Evans described the form of *swaha* from the Safed Kot at the head of the Kurram Valley as *Kurrama*; this name should also apply to the form from Chitral from which it is indistinguishable.

Kurrama Ev. appears to be intermediate between typical *swaha* and *gilgitica* *nihl*.

The types are in my collection and para types in the British Museum.

The correct nomenclature and distribution of *Anlocera swaha* appears to be:—

A. swaha swaha Koll. Outer Himalayas. Murree to Darjeeling.

A. swaha garuna, Finhst. Inner Himalayas, Gurais to hills north of Simla.

A. swaha gilgitica, Tyt. Astor, Chilas, Gilgit.

A. swaha kurrama, Ev. N. W. Frontier of India, Kurram Valley to Chitral.

25. *Anlocera padma grandis*, sub-sp. nov.

A pair taken in Yasin, Western Gilgit, differs from the typical form in having the white discal band on both wings under and the band on the hindwing curved and not straight.

Underside: the ground colour is lighter and more greyish and the white spots in interspaces 3 and 4 of the forewing coalesce and are not separate.

Expanse: ♂ 82 mm.; ♀ 90—98 mm.

Major Stockley obtained several specimens in Eastern Gilgit which agree with this form in all respects except that the white band on hindwing is straight and not curved.

The types are in my collection. The specimens obtained by Major Stockley are in the British Museum.

26. *Karanasa regeli chitralica*, sub-sp. nov.

There were several specimens of a form of *Karanasa* from Chitral above the type and label of *K. moorei* Evans in the British Museum which are not that species as they do not agree with the type but appears to be a race of the *regeli*; they are rather larger than *K. regeli boloricus* but are otherwise the same. I propose the above name for the Chitral form.

Expanse: ♂ 42 mm.; ♀ 49 mm.

The types are in the British Museum and a ♂ para type in my collection.

27. *Karanasa regeli boloricus* Gr-Gr.

A large number of both sexes were received from Misgar in Hunza, Gilgit, which agree with the male type and with a series of both sexes from the Hindu Kush in the British Museum.

28. *Karanasa moorei gilgitica*, sub-sp. nov.

Very similar to *K. moorei moorei* Ev., but on the *upper* and *undersides* of both sexes the yellow markings are rather washed out and have lighter coloured patches; on the underside of the hindwing the light markings are paler.

Expanse: ♂ 52 mm.; ♀ 56 mm.

Five males and six females were received from Yasin and Ghizer in Gilgit.

Colonel Evans places his *moorei* as a race of *regeli* but it is quite a different looking insect and it is better I consider to keep it separate.

29. *Karanasa pamirus safeda*, sub-sp. nov.

The form of *Karanasa* from the Safed Koh, Kurram Valley, sent me by Colonel Evans agrees closely with specimens of *K. pamirus josephi* Stdgr, from Bokhara. It is better placed for the present as a race of *pamirus* than as a race of *huebneri*.

I made no note as to how this form differs from *josephi* Stgr. as I was under the impression that Colonel Evans had described it; but on my return to India he informs me that he did not describe it and although he had appointed types he then thought it was not good enough and asked me to describe it myself. I am unable to do this as I have not the specimens with me. There is however a specimen in the British Museum presented by Colonel Evans labelled *Safeda* Evans and placed as a race of *huebneri* and I think it better to leave this form under the name given it by Colonel Evans than to change it.

The male type is in my collection and a male para type in the British Museum.

30. *Karanasa pamirus pupilata*, sub-sp. nov.

There was a series of a *Karanasa* form from Jhila Drosh, Chitral, in the British Museum and there are also a pair from Chitral in my collection which are different to other geographical forms of *Karanasa huebneri* and its allies and appears to be for the present best placed as a race of *K. pamirus*. In appearance the butterfly is somewhat like *K. pamirus safeda* mihi, but the white patch on the inner edge of the upper ocellus on the forewing is not so well developed and there are no white patches on the inner edge of the ochreous band on the upperside of the hindwing as in that form; on the underside the cell in the male is not marked with white as in *safeda*.

Expanse: ♂ 45 mm.; ♀ 48 mm.

The types are in my collection and para types are in the British Museum.

31. *Karanasa huebneri astorica*, sub-sp. nov.

The form of *huebneri* occurring in Astoris very close to *K. h. modesta* M., which occurs further east in the Deosai Plateau, but in the *males* the orange of both wings extends more into the brown basal area and in the *females* the orange colour replaces the brown basal area of the forewing as in the form *cadesia* M., which however is not a racial form but a casual variety and which

occurs in both the races of *modesta* M. and *astorica* mihi. The female of *modesta* M. has not the orange of the forewing so extended as in the female of *astorica* mihi.

Expanse: ♂ 46 mm.; ♀ 48 mm.

The types are in my collection and ♂ and ♀ para types are in the British Museum.

32. *Karanasa huebneri balti*, sub-sp. n. = *K. leechii*. M. n. Gr-Gr.

Males darker and rather larger than *K. h. pallida* mihi; some males are very close to *K. h. astorica* mihi but the females are very different and are much paler.

Expanse: ♂ 48 mm.; ♀ 50 mm.

There is a good series of this form from Skoro La, Baltistan and from Kardhong in the British Museum.

This insect was given the name of *leechii* by Moore but this name must fall as *Karanasa leechii* Gr-Gr. has priority and was given to a very different form of *Karanasa* from the Hindu Kush Mountains.

33. *Karanasa huebneri pallida*, sub-sp. n.

The race of *K. huebneri* from the Chongking Valley can readily be distinguished from its allies by the extremely pale colouring of both wings in both sexes.

It is very close to the typical form from Lahoul but the yellow colour on both wings enters the brown basal area which it does not do in Felder's form.

There is a good series of this form from the Chongking Valley in the British Museum.

Expanse:—♂ 42 mm.; ♀ 41 mm.

The types are in the British Museum and para types are in my collection.

34. *Karanasa huebneri huebneri*, Felder

The types of Felder's *K. huebneri* came from Lahoul and the female only was figured. There are specimens of *K. huebneri* from Lahoul in the British Museum and these may be considered typical in the absence of the type.

There has been a good deal of confusion and doubt regarding the various forms of *Karanasa huebneri* and allied species inhabiting the Indian Region. Seitz in his *Macrolepidoptera of the World*, vol. i, p. 127, in treating *K. huebneri* Fd. states:—Now follow a whole series of forms of which some intergrade completely, and then includes the following forms under this species—*caesia* M (= *Wilkinsi* Ersch., *josephi* Stdgr.), *leechi* Moore, *modesta* Moore, *dissoluta* Stgr., *intermedia* Gr-Grsh., *pamira* Stdgr., and again states 'all these forms often completely intergrade'.

Colonel Evans in his List of Butterflies from the Indian Region lumps all the Indian forms except one—e.g. *moorei* Evans, which he places as a race of *K. regeli* Alph—under the name of *K. huebneri* Fd. and gives no geographical races.

Without sufficient material it is very difficult to appreciate the differences amongst the various closely allied forms; but with the large series I was fortunate in obtaining from Astor and Gilgit together with other forms in my collection and with the fine series in the British Museum it was found possible to appreciate the collective differences of the various species or races when the individuals were placed in their geographical groups. The forms of *huebneri* and allied species appear to be very susceptible to environment and each distinct locality appears to develop a slightly different race.

The following appears to be a fairly satisfactory nomenclature and distribution of the various forms:—

	Type	Other localities
<i>K. regeli chitralica</i> , Tyt.	... Chitral	
<i>K. regeli boloricus</i> , Gr-Gr.	... Hindu Kush.	Hunza Gilgit
<i>K. moorei moorei</i> Ev.	... Shandur Pass, Chitral.	
<i>K. moorei gilgitica</i> , Tyt.	... Yasin, Gilgit	
<i>K. pamirus safeda</i> Ev.	... Safed Koh, Kurram Valley.	

	Type	Other localities
<i>K. pamirus pupilata</i> , Tyt.	... Jhila Drosh, Chitral.	
<i>K. huebneri modesta</i> M.	... Deosai Plateau.	
<i>ab: cadesia</i> M.	... Deosai Plateau.	
<i>K. huebneri astorica</i> , Tyt.	... Rupal Valley, Astor.	
<i>K. huebneri balti</i> , Tyt.	... Skro La Pass, Bal-	
	tistan.	
= <i>leechii</i> M. nee Gr-Gr.		
<i>K. huebneri pallida</i> , Tyt.	... Chongking Valley.	
<i>K. huebneri huebneri</i> Fd.	... Lahoul.	

35. *Eumenis mniszeckii baldira* M.

The forms of *E. mniszeckii* can at once be separated from those of *E. of lehana* occurring within Indian limits by the presence of two small white centred dark spots or dark spots without the white centres placed between the two large black ocelli on the forewing.

Numerous females were received from Yasin, Gilgit, and a male undoubtedly of this form from Baluchistan was sent to me by Mr. E. Ollenbach. There are numerous specimens of both sexes in the British Museum and all have the dark spots with or without a white centre on the forewing as in typical *E. mniszeckii* but which are absent in *E. lehana*.

36. *Eumenis lehana droshica*, sub-sp. n.

A form of *Eumenis* occurring in Drosh, South Chitral and in the Shandur Pass, North-East Chitral appears to be a race of *E. lehana* but differs from typical *lehana* from Kardong, Ladak in the yellow bands and the basal brown areas of both wings being darker and in the yellow band of the hind wing in both sexes being broader and extending to the tornal angle.

The types are in my collection and para types are in the British Museum

37. *Eumenis lehana gilgitica*, sub-sp. n.

The form of *lehana* occurring in Ghizer, Gilgit, differs from the typical form from Ladak in having the yellow bands on the forewing narrower and darker as in *E. lehana droshica*, and the yellow on the hindwing is also darker as in *droshica*, but the extent of the yellow is as in typical form and does not reach the tornal angle as in *droshica*. A good series of both sexes were taken by Lieut.-Col. Lorimer, Political Agent, Gilgit at Ghizer, Gilgit, and I possess specimens from the Shandur Pass where this race apparently meets with *droshica* mihi.

38. *Eumenis lehana clarissima* Seitz.

There were specimens in the British Museum of a form of *E. lehana* from Skardo Dras, the Chongking Valley and Hunza over the label *Eumenis lehana clarissima* Seitz and I have received specimens from Misgar Hunza and from Upshi which agree with these specimens but I am not at all sure that the name *clarissima* should apply to them as the type of *clarissima* is said to come from West China but this may possibly be a mistake. Those specimens agree with Seitz's figures of *clarissima* and the yellow bands anteriorly are paler.

The form from Skardo and Misgar are very close to typical *E. lehana* and are hardly worth separating.

39. *Erebia mani kamriana*, sub-sp. nov. Pl. I, Fig. 12, ♂

Male and Female: Forewing, apex more pointed than in the typical form and similar to *E. mani chitralica* Ev. but not so pointed as in *E. yasina* mihi and *E. lorimeri*, mihi. Upperside, a large patch on the forewing as in typical *E. mani* de N., but of a deep ferruginous colour somewhat as in *E. kalinda chitralica* Ev. but brighter; basal half of forewing brown; yellow rings to ocellus small. Underside: Forewing the red ferruginous patch large reaching base of v. 1; yellow ring to ocellus as on upperside.

Expanse; ♂ 50 mm.; ♀ 52 mm.

Several males and females were taken on the Kamri Pass, Gurais and a male on the Babusar Pass, Chilas, in August.

The types are in my collection and a male para type is in the British Museum collection.



SOME NEW AND INTERESTING BUTTERFLIES FROM INDIA AND BURMA.

EXPLANATION OF PLATE I.

- Fig. 1. *Lycæna baroghila*, sp. n. ♀.
 „ 2. „ *christophi lesliei*, ssp. n., ♂.
 „ 3. „ *astorica*, sp. n., ♂.
 „ 4. „ *florenciæ*, sp. n., ♂.
 „ 5. „ *galathea depreei*, ssp. n., ♂.
 „ 6. „ *metallica chitralensis*, ssp. n., ♂.
 „ 7. „ „ *gilgitica*, ssp. n., ♂.
 „ 8. „ *orbitulus astorica*, ssp. n., ♀.
 „ 9. *Erebia mani shandura*, ssp. n., ♂.
 „ 10. „ „ *lorimeri*, ssp. n., ♂.
 „ 11. „ „ *yasina*, ssp. n., ♂.
 „ 12. „ „ *kamriana*, ssp. n., ♂.
 „ 13. *Amblypodia suffusa atarana*, ssp. n., ♀.
 „ 14. *Polyommatus sarta rupala*, ssp. n., ♂.
 „ 15. „ „ „ „ ♀.
 „ 16. „ *iolas gilgitica*, ssp. n., ♂.
 „ 17. *Virachola dohertyi*, sp. n., ♂.
 „ 18. *Rapala rubida*, Tytler, ♂ (n.n. pro *rosacea*, Tytler preocc.)
 „ 19. *Amblypodia maymyoica*, sp. n., ♂.
 „ 20. *Orthomiella putaoica*, sp. n., ♂.

40. *Erebia mani shandura*, sub-sp. nov. Pl. I, Fig. 9, ♂

The types, a pair from the Shandur Lake, Chitral, in the British Museum collection are different to any other form of *E. mani* that I have seen and appear to be a distinct race.

Male and Female : Forewing rounded as in the typical form.

Male. Upperside : Forewing, the reddish fulvous patch as in the typical form but darker and also darker than in *E. mani lorimeri* mihi ; but not so dark as in *E. mani kamriana* mihi. Underside, the reddish fulvous patch fills the whole of the forewing almost up to the base ; the pale yellow ring to ocellus broad and distinct.

Female. Upperside : Forewing, fulvous patches as in typical *E. mani* female and much yellower than the male. Hindwing, a small dark ferruginous discal patch present. Underside as in the typical form.

Expanse : ♂ 46 mm. ; ♀ 53 mm.

41. *Erebia mani yasina*, sub-sp. n. Pl. I, Fig. ♂, 11

Forewing : apex pointed and not rounded as in typical form.

Male. Upperside : differs from *E. mani shandura* mihi, in having the apical fulvous patch on forewing much reduced in size ; underside very similar to *E. mani shandura*.

Female. Upperside : very similar to the male but the basal two-thirds of the forewing reddish yellow and the apical third pale yellow.

Expanse : ♂ 50 mm.

A pair were received from Yasin, Gilgit.

The types are in my collection.

42. *Erebia mani lorimeri*, sub-sp. nov. Pl. I, Fig. 10, ♂

Male and Female : Apex of forewing pointed as in *E. mani yasina* mihi.

Male. Upperside : on forewing, a large fulvous patch as large as in the typical form with its inner edge ferruginous ; ocellus without a white centre as in *E. yasina* mihi. Underside : the coloured patch on forewing extending into cell up to V. 1, the outer half being yellowish fulvous and the inner half ferruginous.

Female very similar to the male.

Expanse : ♂ 50 mm. ; ♀ 54 mm.

The types were received from the Haitu Nullah, Ghizer, Gilgit, and are in my collection.

43. *Maniola pulchra chitralica*, sub-sp. nov.

This form is nearest to and between *M. pulchra neoza* Lang from the Kashmir Valley and a form of *pulchra* which occurs in Gilgit up to 9000'. It is smaller than the former but larger than the latter and the female is rather brighter than the female of *M. pulchra neoza* Lang.

Expanse : ♂ 41 mm. ; ♀ 44 mm.

Habitat : Chitral, up to 9000'.

The form from Gilgit besides being smaller is rather paler and approaches the next form *M. pulchra baroghila* mihi which occurs at high elevations.

The types are in my collection and para types are in the British Museum.

44. *Maniola pulchra baroghila*, sub-sp. nov.

The race of *pulchra* from the extreme north-east corner of Chitral and the watershed between Chitral and Wakhan, Afghanistan and the adjoining country on the extreme north-west of Yasin in Gilgit at elevations over 9000' differs from the typical form in being much smaller, being the smallest of all the races, and on the upperside rather paler. On the underside of the hindwing there are small patches of pale yellow in the cell and on the outer edge of the dark discal irregular line.

Expanse : ♂ 37 mm. ; ♀ 38 mm.

A fair series was received from the above mentioned localities.

The types are in my collection and para types are in the British Museum collection.

45. *Maniola pulchra astorica*, sub-sp. nov. Pl. III, Fig. 5, ♀

This race agrees in size with typical *pulchra* from Lahoul but smaller than *M. pulchra neoza* Lang. It differs from the typical and other forms of *pulchra* in the female having the disc of the upper forewing entirely yellow as in *M. pulchella* Fd. The female of *astorica* can however be separated from the same sex of *pulchella* by having a pale yellow iris encircling the black apical ocellus on the forewing which is entirely wanting in *pulchella*.

Expanse: ♂ 42 mm.; ♀ 44 mm.

Numerous males and females were taken in the Rupal Nullah and at Rama in Astor and also a few from Ladak in July and August at 11,000'. The form from the hills surrounding the Kashmir Valley is *M. pulchra neoza* Lang and is the largest of all the races.

46. *Maniola pulchra pulchra*, Felder

The type is probably from Lahoul and a male specimen in the British Museum from that locality agrees with the description of the type and is very yellow on the forewing, approaching *M. pulchella* to which Felder likens it; the hindwing upperside has yellowish discal patches. This specimen and probably Felder's type are aberrant as all the other male specimens from Lahoul, Spiti, Panji, etc., are like the ordinary males from other localities and have the forewing covered with dense brown scales and no traces of yellow on the hindwing.

I possess specimens from Garhwal which probably belong to this race.

47. *Maniola cheena chitralica*, sub-sp. nov.

The Chitral race of *M. cheena* has the yellow iris to the ocellus on forewing as well developed as in the typical form, but the insect is much smaller and of the same size as *M. cheena kashmirica*. The types of *M. cheena* M. are from Kunuwar and are in the British Museum. The male has yellow patches below the apical ocellus on the upperside of the forewing and the female has also the yellow on the forewing well developed and conspicuous. The male type appears to be aberrant as other specimens from the same locality have only traces of the yellow patches below the apical ocellus on the forewing.

The Kashmir race *M. cheena kashmirica* is smaller than the typical form from Kunuwar and the yellow iris to the ocellus on forewing is smaller and not so conspicuous.

The types are in my collection and para types are in the British Museum.

The following appears to be the nomenclature and distribution of the various forms of *Maniola* within Indian limits:—

	Type locality	Other localities
<i>M. pulchella</i> , Fd. Ladak.	Astor, Gilgit, Deosai Plateau, Dras.
<i>M. pulchra chitralica</i> , Tyt.	... Chitral, below 9,000'	Gilgit below 9000'.
<i>M. pulchra baroghila</i> , Tyt.	... Baroghil Pass, N. E. Chitral, above 9,000'	Yasin, N. W. Gilgit, above 9,000'.
<i>M. pulchra astorica</i> , Tyt.	... Astor.	G u r a i s, D r a s, Ladak.
<i>M. pulchra neoza</i> , Lang.	... Kashmir.	Tragbal, Pir Panjal, Shamshiri, Dugi Pass.
<i>M. pulchra pulchra</i> , Fd.	... Lahoul?	Lahoul, Dalhousie, Kokser P a s s; Garhwal.
<i>M. cheena chitralica</i> , Tyt.	... Tarban, Nala and Utzen, Nala Chitral.	
<i>M. cheena kashmirica</i> , M.	... Kashmir.	
<i>M. cheena cheena</i> Kunuwar.	

48. *Lethe armandii khasiana*, M.

There has been a great deal of confusion about this form and in the Journal of the Bombay Natural History Society, vol. xxiii p. 223 I attempted to clear up the matter and showed this *khasiana* was not a form of *bhadra* but that it was an *armandii* form and I thought it was true *armandii* but erroneously so. I have

lately compared the type of *hasiana*, which is a dry season form, with true *armandii* from China and find that there is a marked difference in the two forms and that the former will now have to stand as a race of *armandii*. In the wet season males of *hasiana* from the Naga Hills all the yellow markings are brighter and larger than in the wet season form of *armandii* from China and the cell of the forewing has at its apex a yellowish bar; the females are very similar to the females of typical *armandii* with the exception of the presence of the apical yellow bar to the cell of the forewing, absent in typical *armandii*.

In the dry season male type of *hasiana* the markings on the upperside of the apical area of forewing are white and not yellow as in the dry season form of typical *armandii* and the yellow on the hindwing is darker and tinged with orange and enters the cell.

A single male of the dry season form from Bernardmyo, Upper Burma, in the British Museum is intermediate between *hasiana* from the Khasi and Naga Hills and *armandii* from West China; the forewing agreeing with *hasiana* and the hindwing with *armandii* in the shade of yellow which however enters the cell as in *hasiana*.

The following appears to be the correct nomenclature and distribution of *L. armandii* :—

<i>L. armandii armandii oberthur</i>	...	d. s. f;	West China.
= <i>fusca</i>	...	w. s. f;	
<i>L. armandii hasiana</i> M	...	d. s. f;	Khasi Hills, Naga Hills.
= <i>unnamed form</i>	...	w. s. f.	Naga Hills.
<i>L. armandii</i> sub sp.	...	d. s. f.	Bernardmyo, North Burma.

<i>L. armandii laticolora</i> , Fruhst	...	w. s. f.	
= <i>saggitata</i> Wilen	...	d. s. f. ?	Formosa."

The form from Bernardmyo may form a local race but more material is required to determine whether the differences above noted are constant or not.

49. *Mycalesis suaveolens sebonga*, sub-sp. n.

A male and female from the North Chin Hills in the British Museum and a large series taken by my collectors at Sebong, Manipur, and in the Manipur Valley and Naga Hills are quite distinct from the typical form of *Suaveolens* from Sikkim and Bhutan and form a well marked race.

Underside: wet season form very similar to the typical form but all the ocelli very much smaller; the dry season form in addition to having the ocelli much smaller has the outer area beyond the discal band on both wings very conspicuously paler.

The types from Sebong are in my collection and para types from the Chin Hills are in the British Museum.

50. *Mycalesis suaveolens duguidi*, sub-sp. n.

Male and female: *Upperside* very similar to *M. suaveolens sebonga* mihi. *Underside* differs in being paler brown and the white bands on both wings tinged with brown; cilia brown and not white as in the typical form and in *sebonga*.

Several males and females were received from Loimwe in the Southern Shan States caught in May and August at about 5,000'.

The types are in my collection and a male para type is in the British Museum.

The occurrence of this form in the Southern Shan States considerably extends the range of the species to the south and east.

I have much pleasure in naming this form after Capt. Duguid who obtained the specimens and kindly presented them to me.

51. *Faunis arcesilaus* F.

A male and female received from Loimwe, Southern Shan States, differs from the typical form in having the apical third of the forewing on the underside conspicuously paler than the rest of the wing.

Loimwe is close to the Chinese border in the Southern Shan States and the butterflies from this locality are but little known. It is quite possible that these specimens represent a race of this species.

52. *Faunis faunuloides* de N.

Colonel Evans in his 'Identification of Indian Butterflies' states that the genus *Faunis*, Hubner has a prominent tuft on the upperside hindwing from v. 1 below base of cell and that there may be a brand on 1a of the hindwing near the tornus covered by a tuft; and in his key to the genus *Faunis* states that the male *faunuloides* on the upper hindwing has a brand on v. 1 (a). He gives a figure of *F. faunula* from the Malay Peninsula which clearly shows the brand on v. 1 (a). He also places *F. faunuloides* de N. as a race of *faunula*.

Fruhstorfer in Seitz's work also mentions that *F. faunula* has a second tuft on the hindwing and also places *Faunuloides* as a race of *faunula*.

Fruhstorfer states that only five females have so far been secured from the Chin Hills and presumably he has never seen a male and it is possible that Colonel Evans also has not seen the male and that the male has hitherto been unknown.

I have lately received two males and three females from the Chin Hills and they are exactly the same as the female but for the sex mark.

The males however do not agree with Evan's key in the following respects :—

(a) On the hindwing the tuft at base of cell is from below the median nervure and not from v. 1.

(b) There is no brand on v. 1 (a) near the dorsum hindwing.

As regards (a) Evans probably intended to say *median vein* instead of v. 1; but as regards (b) he is incorrect in stating that *faunuloides* has a brand on v. 1 (a) of the hindwing. If *F. faunula* from the Malay States has the brand on v. 1 (a) near the dorsum of hindwing as Evans and Fruhstorfer state then I consider that *F. faunuloides* de N. which has not got this brand cannot be conspecific and must retain its specific rank.

The specimens received by me were taken by my collector in September; de Nicéville and Fruhstorfer record it as having been taken in June. It is therefore probably double brooded.

53. *Aemona lena haynei*, sub-sp. nov. Pl. III, Fig. 9, ♂

Male. *Upperside*: forewing; yellow with patches of paler colour in the interspaces beyond the discal brown band; apical area dusted with smoky brown scales. *Hindwing*: yellow the area between the double marginal lines in interspaces 4, 5 and 6 sometimes but not always paler than the ground colour. *Underside* as in the typical form.

Female as in the typical form.

In *Aemona lena* the sexes are alike; in *haynei* they are very different the male being a yellowish insect lacking nearly all the pale markings of the female.

Habitat: Maymyo, N. Shan States.

A fair number of both sexes were obtained in April and May and again in September, in dense shade and near water. The insects as a rule keep to the low undergrowth a foot to three feet above the ground and on being disturbed fly a short distance and settle down again. On one occasion I saw about half a dozen males sitting on the stem of a tree 20' up. It is not rare but appears to be very local.

The types are in my collection and para types are in the British Museum.

I have much pleasure in naming this form after Dr. Hayne who first discovered it and told me where to look for it.

(To be continued)

A CONTRIBUTION TO THE ORNITHOLOGY OF DELHI

BY

THE LATE S. BASIL-EDWARDES, M.B.O.U., R.A.O.U.

PART I

(With 2 plates)

During the winter of 1924-25 while I was in Delhi I made a small collection of birds and accumulated a number of notes. This was done with the dual object of adding specimens new to my collection and of enabling me to prepare a list of the birds of Delhi at some future date. As, however, I cannot readily trace any literature dealing specifically with the birds of this locality, and as I was fortunate in obtaining some interesting specimens, some useful purpose may perhaps be served by placing these notes on record now, instead of waiting indefinitely. Five months is too short a time in which to obtain anything approaching a complete knowledge of the birds of the district; but two or three years' continuous observation and collecting in all types of country ought to produce interesting results. I did not obtain, nor even did I see, any specimens of certain families of birds, e.g., Buntings or Rails, not because there were none to be had, but because I could not find the time to work suitable localities. I have included the names of all species observed in order to make this paper more complete; but the following notes can only be regarded as a nucleus for a more comprehensive paper on the ornithology of the Delhi Province. If and when a suitable opportunity arises, I hope to add further notes on the birds of Delhi, and by this means place on record a fuller account of the avifauna of the Province.

My activities were confined to the immediate neighbourhood of the City itself, chiefly at Raisina, with occasional excursions to places nearby, such as Okla, Kingsway, etc. Bird-life is abundant in the Delhi Province, and quite a number of common species visit one's garden. I found the babool jungles in and around Raisina very convenient and profitable hunting-grounds, as also the Ridge, and the canal near Kingsway. Unless otherwise stated, my notes refer to the neighbourhood of Raisina. I was not on leave and most of my leisure hours were occupied in collecting birds only, so that I could pay little or no attention to birds' nesting. I have, however, included a few stray notes on nests found during my stay in Delhi in the winter of 1924-25 and on some eggs collected by me a few years ago. I have also included particulars of eggs taken by Mr. F. H. Cole, an ardent young oologist, who was in Delhi during the months of November 1924 to March 1925. Scattered throughout *Nests and Eggs of Indian Birds* are a number of notes by Major C. H. T. Bingham on the eggs collected by him in Delhi, and these notes I have culled and incorporated briefly in my paper, as I think they form a useful and valuable complement. Reference to Bingham's oological notes on Delhi birds will be found in square brackets, thus: '[Eggs taken by Bingham April-July]' I have endeavoured wherever possible to indicate the status of each species recorded below; it should be explained, however, that I have in many cases regarded certain species as resident, basing my conclusion on my own personal observation of such species throughout the winter, and on Bingham's oological notes which refer to the summer months. I think that this is justifiable, and will, in the majority of instances, be found to be correct. I arrived in Delhi on November 2, 1924, and left for Simla on April 3, 1925, and during this period of five months I collected almost daily. The total number of specimens obtained was about 230—a fair result, as I had to work unassisted at collecting, skinning, and writing up my notes. In the case of certain species a small series of each was obtained.

As most of my collecting was done in and around Raisina and the outskirts of Delhi City, with occasional excursions to places near by, such as Kingsway and Okla, it does not appear necessary to describe the boundaries and extent of the Delhi Province. Similarly, I shall not enter into any detail about the flora or climate of the province as a whole. All this may be left for some future occasion when my ornithological activities in Delhi have been more extended. For the present, therefore, I shall confine myself to a few remarks on the kind of country I have actually seen and worked.

For the most part Raisina, or New Delhi as it is also called, is a gently undulating, open tract—a city and its environs in embryo, which is developing rapidly. The country round about consists of open spaces, covered either with short grass or low, scrubby bushes, with the babool (*Acacia arabica*) growing freely everywhere. On the outskirts of Raisina, scattered about in patches of greater or lesser extent, are babool jungles. Fields and cultivated tracts occupy certain portions. Deserted towns and tombs lie in every direction; some cared for under Government supervision, some dilapidated and crumbling-falling to pieces amid desolate and dreary surroundings. Where there are pools, canals, or borrow-pits, reeds and long grasses fringe the water's edge. The banks of the River Jumna are, in winter, bare stretches of sand on which the tamarix or jhow (*Tamarix dioica*) grows in great profusion. At Okla is situated a weir and sluice-gates where the waters of the Jumna are controlled and diverted by canals. Some sort of a park and garden exists here—a pleasing contrast to the bare and rather monotonous aspect of the banks which stretch out on either side beyond the grounds at Okla. The historic Ridge of Delhi, which is an extension of the Mewat branch of the Aravalli Hills, forms a conspicuous and characteristic physical feature. The vegetation on the Ridge consists, as may be expected, largely of low, thorny bushes and cactus. At Kingsway there are low-lying fields, subject to inundation during the rains. The Nuzzargarh Cut Drain also passes through Kingsway. Off the Karnal road lies the comparatively large 'Horse-shoe Jheel', so-called on account of its shape.

Leaving out of consideration the babool (*Acacia arabica*), which is excessively common, some of the chief trees are *Salvadora persica* and *Butea frondosa* (which predominate in the uncultivated tracts), *Tamarindus indica*, *Melia azadirachta* and *azadarach*, various kinds of *Ficus*, etc. The only wild palm is *Phoenix sylvestris*. *Trappa bispinosa*, *Nelumbium speciosum* and *Typha latifolia* are common water-plants; while *Opuntia dillenii*, *Agave americana* and *Ala vera* are the commonest succulents. The following wild plants may be mentioned: *Zizyphus jujuba* and *nummularia*, *Prosopis spicigera*, *Cryptostegia grandiflora*, *Argemone mexicana* and *Adhotada vasica*. *Saccharum arundinaceum* grows in profusion on embankments, etc.; *Berberis aristata*, a dense, thorny shrub, grows along the borders of cultivated fields and in dry, rocky situations; *Capparis aphylla* and *horrida* are common in waste lands and among ruins; and *Tribulus terrestris* is a common weed.

I am only able to write of the winter in Delhi, as I have had no personal experience of the summer months or the rains. It is rather pleasant during the months of October and November, but the weather thereafter becomes cold. By the middle of February it begins to warm up again, and the latter half of March is unquestionably hot, though the nights are fairly pleasant if there is a breeze. In March high winds and dust-storms are of rather frequent occurrence.

A few photographs have kindly been taken for me by Mr. E. S. Keymer. These illustrate various types of country and serve to indicate the haunts of certain species of birds, while they also possess a topical interest.

The scientific nomenclature used is that employed in Mr. E. C. Stuart Baker's *Hand-List of the Birds of the Indian Empire*. Modifications have been made where necessary in the light of the first two volumes of the second edition on Birds in the *Fauna of British India* series and of other publications.

In conclusion, I wish to express my thanks and gratitude for the kindly assistance in many ways which has been rendered me by Mr. H. Whistler, F.Z.S., M.B.O.U., and by Mr. A. E. Jones, M.B.O.U. My thanks are also due to Dr. C. B. Ticehurst, M.D., M.B.O.U., for the willing manner in which he has identified certain skins and for some useful suggestions in regard to the preparation of this short paper.



BABOOL TREES ALONG THE MUTTRA ROAD
Haunts of *Argya malcolmi*, *Phoenicurus o. phoenicuroides*,
Pericrocotus p. peregrinus, *Lanius vittatus*, etc.



BABOOL JUNGLE AT RAISINA, DELHI
The haunts of a variety of small birds.



THE CANAL AT OKLA, DELHI

Haunts of *Ceryle e. leucomelanura*, *Ardeola grayii*, etc.
Clump of trees in background—haunts of *Cuncuma leocorypha*.



A PORTION OF THE RIDGE AT RAISINA, DELHI

Haunts of *Argya c. caudata*, *Franklinia buchanani*, *Lanius e. lahtora*, *Uroloucha malabarica*, *Cyrtostomus a. asiaticus*, etc.

Corvus corax laurencei (Hume). The Punjab Raven.

Hume writes: 'Even in the Delhi Division of the Punjab they breed sparingly . . .' I did not meet with any Ravens.

Corvus splendens splendens (Vieill.). The Indian House-Crow.

Four winters ago I remember seeing a specimen which had some white in the wings. This bird would frequent the neighbourhood of Kingsway Camp. The bones or flesh of this bird, in the eyes of the Hindus of Delhi, appear to possess some medicinal or magical properties. When out collecting near a village one day I was asked by a Hindu villager to shoot a Crow which he said he required for a certain purpose. I was not able to ascertain the exact purpose, however. Subsequently, I saw a man hawking round many dozens of live Crows for sale. These were being carried in flat baskets covered with string meshing—one basket at either end of a bamboo slung across the shoulders. The baskets were similar to those in which poultry and wild duck are carried. I have no information as to the method of capture of the Crows.

Corvus coronoides subsp. Jungle-Crow.

Bingham writes: 'This Crow (*macrorhynchus*, Wag.), so common at Allahabad, is very scarce here at Delhi. In fact, I have only seen one pair.'

I can endorse Bingham's remarks about the scarcity of Jungle-Crows in Delhi: they are certainly not common. A pair were seen on December 14, 1924, haunting an old ruin known as the Hauz Khas. The birds were not seen again when I visited the place on December 28, 1924, and during the rest of my stay I saw no *coronoides* Crows though I was always on the watch for them.

Dendrocitta rufa vagabunda (Blyth.). The Bengal Tree-pie.

Quite common. A specimen obtained belongs to this race. Wing 165 mm. Tail 285 mm.

Argya caudata caudata. (Dum.). The Common Babbler.

Generally found among long grass, or on thorny bushes on the Ridge. I neglected to collect a sufficient number of this common bird until it was too late. Only two specimens were procured. A ♂ on February 28, 1925 and a juv. ♀ on December 5, 1924. These appear to be of the typical race. Wing of ad. ♂ 76 mm.

[Eggs taken by Bingham. March-June.]

Argya malcolmi (Sykes). The Large Grey Babbler.

This is by far the commonest Babbler in Delhi. Abundant everywhere in exceedingly noisy parties. Generally nests in babool trees.

[Eggs taken by Bingham. March-August.]

Turdoides terricolor terricolor (Hodg.). The Bengal Babbler.

Common. Appears to prefer shady gardens and groves to babool trees, which is the typical haunt of *Argya malcolmi*. This Babbler seems to be rather a wary bird at times.

[Eggs taken by Bingham. April-July.]

Ægithina nigrolutea (Marsh). Marshall's Iora.

I did not find this pretty little bird at all common. Two obtained on December 11, 1924, were haunting babool jungle. One of them, apparently the ♂, was observed to fly up into the air and come spinning down to the tree with his feathers fluffed out. This habit has been described in the *Fauna of British India*. Birds, vol. i, page 342, second edition. An examination of the genital organs of the birds obtained showed them to be normal, so that this curious habit is not confined to the breeding season. Another pair of Ioras were seen on the Muttra Road, about 8 miles from Delhi, on February 8, 1925, and both were obtained.

At the time I obtained these Ioras I thought they were *Ægithina tiphia*, *tiphia* but on examination I found them to be undoubted examples of *nigrolutea*. Hume states (*Stray Feathers*, vol. vii, page 454) that all the Ioras collected by Cleveland in the Gurgaon District were, without exception, *nigrolutea*. In this connection his 'Remarks on the Genus *Iora*' in *Stray Feathers*, vol. v, pp. 420-452, are worth perusal. Of thirty specimens obtained by Hume, five were from Delhi (Gurgaon?). According to him *nigrolutea* is a species with a defined area. Mr. Stuart Baker (*Fauna of British India* Birds,

Ægithina nigrolutea.—(contd.)

vol. i, p. 344) states that this species 'takes the place of *Æ. t. tiphia* and *Æ. t. humei* to the north-west of India', overlapping the range of both in certain parts of India. Wing of four specimens 62–66 mm.

Molpastes hæmorrhous pallidus (Stuart Baker). The Central Indian Red-vented Bulbul.

Common. Seen along with *M. l. leucotis*. Writing from the neighbourhood of Delhi, Mr. F. R. Blewitt says: 'This Bulbul breeds from the middle of May to about the middle of August.'

[Eggs taken by Bingham. April–July.]

The notes in the new edition of the *Fauna of British India* regarding the *hæmorrhous* Buleuls are not quite clear. Two specimens obtained belong to this pale race. My identification has kindly been confirmed by Mr. Whistler.

Molpastes leucotis leucotis (Gould). The White-eared Bulbul.

Also very common. This species and the last-mentioned are often seen together. Mr. Jones tells me that this species does not occur in the Amballa District.

Sitta castaneiventris castaneiventris (Frank.). The Chestnut-bellied Nuthatch.

Collecting as I did chiefly in babool jungle and wasteland, I did not meet with this Nuthatch whose haunts are shady groves and gardens. Cole tells me that he saw it on several occasions in the Kudsia Gardens. He obtained a ♀ on December 11, 1924, and this specimen I have seen.

Dicurus macrocercus macrocercus (Vieill.). The Black Drongo.

Very common everywhere, and, I think, a resident species.

[Eggs taken by Bingham. May to (?) August.]

This species breeds much earlier, I think, in Delhi, for I remember seeing a nest towards the end of March (1922).

Salpornis spilonota (Frank.). The Spotted Grey Creeper.

There is an old record of a nest found by Mr. Cleveland at Hattin, in the Gurgaon District, on April 16.

Acrocephalus dumetorum (Blyth.). Blyth's Reed-Warbler.

A single specimen of what may have been this species was seen among babool trees on the Ridge on March 25, 1925. I much regret having missed the bird, which disappeared among the trees and bushes. As Dr. Ticehurst remarks, this species is more of a Tree-Warbler than a Reed-Warbler in its habits (*Ibis*, July 1922, p. 550). In Simla, when this Warbler passes through in May, it is found among bushes and trees. As I did not get the bird its exact identity is doubtful. The bird seen may have been *Hippolais rama*: in fact, Mr. Jones thinks this was probably the species. Both look much alike in the field, and I may have been mistaken.

Orthotomus sutorius sutorius (Forst.). The Indian Tailor-Bird.

Abundant. A very common garden bird.

[Eggs taken by Bingham. April–June.]

Cisticola juncidis cursitans (Frank.). The Rufous Fantail-Warbler.

Bingham writes: 'I have not yet observed this bird at Delhi.' He seems to imply that it is likely to be found there. I met with this peculiarly marked Warbler on only one occasion on February 21, 1925, when I was out after Sand-Grouse in the outskirts of Raisina. Marking down a spot where some Grouse had settled, I happened to pass through a small patch of grass about two feet long, when I disturbed a Warbler which was new to me. It flew some little height above the grass, immediately being lost to view when it dropped down among the yellowish brown grass. In trying to find this bird I flushed another which behaved in the same manner. I decided to watch the exact spot where a bird settled, and when one rose again I marked it sitting quietly at the base of some grass. This specimen I shot; but I was not able to find the second bird. The way in which the streaked plumage of this species blends with the grass among which it lives is remarkable.

Franklinia buchanani (Blyth.). The Rufous-fronted Warbler.

Another common species, especially among jherberry bushes on the Ridge. It is also commonly seen among babool trees. This Warbler has rather a distinctive song, which consists of a trill, followed by a triple note like *sirriget-sirriget-sirriget*. A ♂ shot on December 12, 1924, was breeding.

[Eggs taken by Bingham. October.]

Sylvia curruca affinis (Blyth.). The Indian Lesser Whitethroat.

Abundant. Always to be seen among babool trees, though it has also haunts low bushes. Its harsh note is always to be heard in babool jungle.

Phylloscopus collybita tristis (Blyth.). The Siberian Chiff-Chaff.

When out collecting along the Bela Road at the back of the Fort on March 8, 1925, a specimen (sex?) was procured. Several were seen among the jhow bushes on alluvial land. Another specimen (a ♂) was shot on a Neem tree near the District Jail on the same date. Both specimens were in fat condition preparatory to migration.

Phylloscopus indicus (Jerd.). The Olivaceous Willow-Warbler.

A specimen (ad. ♀) was obtained in babool jungle on March 21, 1925. This species was also observed passing through on migration. Towards the end of March : I have no details of dates.

Phylloscopus humii humii (Brooks.) Hume's Willow-Warbler.

The note of this Willow-Warbler was heard throughout my stay. Unfortunately no specimens were obtained, but a specimen shot for me at Okla proved to be this species.

Phylloscopus nitidus nitidus (Blyth.). The Green Willow-Warbler.

Three or four were seen on March 21, 1925, in babool jungle on the outskirts of Raisina, and an ad. ♂ obtained. Noticed during March and April passing through on migration.

Prinia gracilis lepida (Blyth.). The Streaked Wren-Warbler.

The haunts, *par excellence*, of this Warbler appear to be long grass. I did not meet with it anywhere else. On February 7, 1925, numbers were seen, flitting from one clump of grass to another, at the back of the Old Fort.

There is an interesting account of the nidification of this species in Delhi by Colonel C. H. T. Marshall in *Nests and Eggs*, vol. i, p. 288.

[Eggs taken by Bingham. March.]

Prinia socialis stewarti (Blyth.). Stewart's Ashy Wren-Warbler.

Bingham says : ' At Delhi I have not yet found its nest. I once found in July three nests all attached together in a sort of triangle, but whether built by separate pairs of birds I cannot say. Only one nest contained eggs.' This is a very common species, affecting the same type of country as *Prinia i. inornata*. Like that species it loves to investigate clumps of long grass, bushes, etc.

Prinia inornata inornata (Sykes.). The Indian Wren-Warbler.

Abundant. A common garden bird. Flits from bush to bush making a curious snapping noise. In the hills *Suya crinigera crinigera* indulges in the same habit.

Lanius excubitor lahtora (Sykes.). The Indian Grey Shrike.

Common all over, but its chief haunts appearing to be the rocky Ridge. In *Nests and Eggs*, vol. i, pp. 307-308, there is an interesting note by Mr. F. R. Blewitt on his experiences of the nidification of this Shrike in Delhi. Resident.

[Eggs taken by Bingham. March--August.]

Lanius vittatus. (Valenc). The Bay-backed Shrike.

Undoubtedly the commonest Shrike in Delhi. Seen among babool trees along the roads. This Shrike is a good mimic. Several specimens collected, including a nicely marked juv. ♂ obtained on December 3, 1924. Blewitt has a note on this species in *Nests and Eggs*, vol. i, pp. 312-313.

[Eggs taken by Bingham. May-July.]

Lanius schach erythronotus (Vig.). The Rufous-backed Shrike.

Not near as common as either of the two Shrikes already mentioned. If it has any particular haunts, these appear to be thin babool jungle interspersed with long grass, and also large gardens. I have seen specimens in the grounds of Metcalfe House and the Secretariat. As we require more information about this Shrike, I may mention that I met with it on several occasions during my stay in Delhi. The first occasion on which I saw it was on November 25, 1924, on which date a single bird was seen in a clump of bamboo and flowering shrubs in the grounds of the Secretariat in Old Delhi. On December 4, 1924, I obtained a fine ♂ in an open patch of ground opposite my residence in New Delhi, and on the following day I procured a ♀ in the same place. These two birds must have been together. On December 19, 1924, another specimen was seen, and this bird was also found in the same locality. On January 16, 1925, a ♀ was shot for me. A month later, on February 17, 1925, I saw two of these Shrikes in thin babool jungle at Raisina, and I saw a specimen in the grounds of Metcalfe House on March 30, 1925, when I was collecting examples of *Pastor roseus*, but I have no record of this in my diary. Bingham found it breeding in June.

[Eggs taken by Bingham. One nest containing two eggs on June 3, in the Nicholson Gardens.]

Lanius cristatus isabellinus (H. and E.) The Pale Brown Shrike.

Uncommon. I met with this species as follows :—

- (a) A single bird seen on November 10, 1924, was shot at but not obtained.
- (b) Juv. ♀ obtained on November 26, 1924.
- (c) Ad. ♀ obtained on February 1, 1925.
- (d) A single bird seen on February 17, 1925.
- (e) A single bird seen in open fallow land off the Bela Road near the Jumna on March 8, 1925. A friend and I had several long shots at it, but in the end it eluded us!

Tephrodornis pondicerianus (Gmel.) Wood-Shrike.

Rather common among babool trees, its curious plaintive crescendo notes attracting attention. The north-western Indian bird has been separated by Ticehurst and called *pallidus* (vide B. O. C., xii., p. 56). Without a good series it would be better to treat the Delhi bird binomially.

[Eggs taken by Bingham. Date not mentioned.]

Pericrocotus brevirostris brevirostris (Vig.). The Indian Short-billed Minivet.

About half a dozen birds seen on January 15, 1925, in a thick clump of babool trees at Raisina. No ad. ♂♂ in the black and red plumage were observed however. I was unable to obtain any specimens as my supply of twenty-two cartridges had run out. Two days later two more ♀♀ (or immature ♂♂) were seen in the same place, but the birds managed to elude me by disappearing among the trees. On March 14, 1925, three more were observed. Two specimens were shot, one of which could not be found. It is strange that I saw no red ♂♂.

Pericrocotus peregrinus peregrinus (L.). The Small Minivet.

Rather common. Two ♂♂ in first plumage obtained on the road to Okla on November 22, 1924. Two ad. ♂♂ on January 1925.

[Eggs taken by Bingham. Earliest nest on March 21.]

Pericrocotus erythropygius (Jerd.). The White-bellied Minivet.

This elegant little Minivet is not uncommon at Delhi. It is usually seen in small parties of about half a dozen birds. Seen pecking at weeds a foot or two high. A party of four or five birds seen on November 25, 1924, and three ♂♂ obtained, one of which, however, was too badly damaged for preservation. Another ♂ obtained on January 25, 1925, and another of the same sex on February 4, 1925. A solitary ♂ seen near the Jantar Mantar on January 24, 1925. I did not get a ♀ specimen.

Oriolus oriolus kundoo (Sykes.). The Indian Oriole.

I saw no Orioles in Delhi during my stay there.

[Eggs taken by Bingham. April-July.]

***Sturnus vulgaris poltaratzskii* (Finsch).** Finsch's Starling.

This, I think, is the common race found in Delhi. At the commencement of winter this Starling is to be found in small parties, in twos and threes or even singly, but by the middle of February enormous flocks are to be seen settling in fields and open spaces. A decent series of Starling would probably reveal the occurrence of one or two other races besides *poltaratzskii*.

***Pastor roseus* (L.).** The Rosy Pastor.

This beautiful Starling was not seen at all till the end of my stay in Delhi. They must have arrived during the third week of March, as the first birds I saw (a couple) were on March 3, 1925, and a friend told me of some he saw two or three days before. On March 25, 1925, I saw flocks of varying size flying about, and others were seen on peepul trees on the same date. On March 26, 1925, I went out after them securing three specimens. On March 30, 1925, I obtained three more, in the grounds of Metcalfe House near the Secretariat. Up to the time I left Delhi flocks were seen daily, flying north in the mornings and evenings. This species is generally supposed to regulate its migratory flight with the flowering of the cotton tree, and it is therefore worth mentioning that I did not once see a single bird on a cotton tree, though this tree was in bloom at the time that the birds were found in Delhi. Four specimens were shot on trees bearing small green berries (I do not know the name of the tree). This tree appears to be the favourite resort of Rosy Pastors, the peepul also being frequented for its fruit, but not to such an extent. The song of this species consists of a jumble of harsh grating noises, intermingled with some really tuneful notes. In the field the presence of the birds may be known at once from the clamour they make. The long black feathers on the head and throat of a bird in full plumage stand out prominently when the bird is calling.

The colours of the soft parts are as follows :—

Fully adult birds in the pink and black plumage.

Irides—very dark brown, almost black.

Bill—fleshy, with the tips and culmen brownish, the base of the lower mandible blackish-plumbeous. Inside of mouth plumbeous posteriorly, pink anteriorly.

Legs and feet—brownish-yellow, tinged fleshy.

Claws—blackish-brown, with pale bases.

Birds in immature plumage.

Irides—As above.

Bill—pale flesh, with tips and culmen dark horny-brown, and base of lower mandible dull yellow. Inside of mouth dull yellow tinged plumbeous, with anterior portion of inside of both mandibles fleshy.

Legs and feet—dark-brown.

Claws—blackish-brown, without pale bases.

***Temenuchus pagodarum* (Gmel.)** The Black-headed Myna.

This elegant little Myna was seen in pairs or in small parties in thin babool jungle and in other suitable localities.

[Eggs taken by Bingham. May-July.]

***Acridotheres tristis tristis* (L.)** The Common House-Myna.

Abundant. Resident.

***Acridotheres ginginianus* (Lath.)** The Bank Myna.

Common. Often seen attending grazing cattle. Numbers of these Mynas are to be seen at the Delhi railway station and at other stations on the line towards Ambala. They seem, in fact, to take the place of *A. t. tristis* at the railway stations.

***Sturnopastor contra contra* (L.)** The Pied Myna.

This species is found bathing in the evenings along with *Acridotheres t. tristis*. It seems by preference to haunt country in the vicinity of water, and although a common species, is not usually found elsewhere.

[Eggs taken by Bingham. May-July.]

Siphia parva parva (Bechst.). The European Red-breasted Flycatcher.

A ♂ seen on January 11, 1925, in a large peepul tree in the grounds at Okla. Quite a number seen in babool jungle on March 22, 1925, on which date a ♂ and two ♀♀ were obtained. These were hawking insects among the trees. The note of this species is a *chit-trr* and is much like that of *Cyornis superciliaris*, but decidedly softer. ♀♀ appeared to be far in excess of ♂♂. A pair seen on March 26, 1925 and the ♀ obtained.

Stoparola melanops melanops (Vig.) The Verditer Flycatcher.

Miss Holmer in her book entitled *Indian Bird-Life* states that she has seen this Flycatcher in 'kikar jungle in New Delhi.' I did not meet with it.

Culicicapa ceylonensis ceylonensis (Swains.) The Grey-headed Flycatcher.

The note of this little Flycatcher was heard emanating from a clump of neem trees on the Okla road. No specimens obtained.

Terpsephone paradisi paradisi (L.) The Indian Paradise Flycatcher.

I did not see this species, and it had not arrived before I left Delhi.

[Eggs taken by Bingham. Seven nests on May 27, and June 12.]

Rhipidura aureola aureola (Less.) The White-browed Fantail Flycatcher.

Commonly seen singly or in pairs in babool jungle along roadsides.

[Eggs taken by Bingham. July 8.]

Saxicola caprata bicolor. (Sykes). The Northern Indian Pied Bush-Chat.

Common among bushes and shrubs in waste-land.

I obtained a nest at Kingsway towards the end of March 1922 containing four eggs.

[Eggs taken by Bingham. May-June.]

Saxicola torquata indica. (Blyth.). The Indian Bush-Chat.

Not quite as common as the species last mentioned, but, of course, haunting the same kind of ground.

Oenanthe picata (Blyth.). The Pied Chat.

Fairly common. Not infrequently seen on the roofs of houses and on walls. Far more ♂♂ than ♀♀ seen. Dr. C. B. Ticehurst noticed this same phenomenon in Lower Sind, but is unable to offer any reason for it. (*Ibis*, 1922, p. 633). I did not obtain, or even see, any specimens of the *capistrata* form of this species, though I kept a very sharp look-out for it. This is probably a distinct species or subspecies, rather than a dimorphic form of *Oe. picata*.

Oenanthe isabellina (Cretzsch). The Isabelline Chat.

Several pairs of what I took to be this species were seen on fallow land near the Hauz Khas on December 28, 1924. A stiff breeze was blowing that day and I tried several shots without success. A single bird seen on February 21, 1925, in bare, open ground, interspersed with a few thorny bushes and deep dry drains. This bird was far too wary and I did not get it. Four winters ago I saw this species in fields near Kingsway, but on two occasions when I visited this locality in 1924-25 I did not meet with a single Chat of this species.

Cercomela fusca (Blyth.). The Brown Rock-Chat.

Very common among piles of stones, and in rocky ground; also a most domesticated species, coming right into the house, and perching in the early mornings on the tops of windows, etc. This bird is a good mimic. I have heard it imitate the plaintive notes of *Tephrodornis pondicerianus*, *Tringa ochropus* and one or two other species.

[Eggs taken by Bingham. March-August.]

Phoenicurus ochrurus phoenicuroides. (Moore). The Western-Indian Redstart.

Very common, especially among babool trees on which it perches freely. Seen throughout my stay in Delhi.

Cyanosylvia suecica pallidogularis (Zarud.) The Eastern Redspotted Bluethroat.

Haunts bushes and scrub-jungle close to water, and clumps of plants in gardens. Two ♂♂ and a ♀ belong to this race.

Saxicoloides fulicata cambaiensis (Lath). The Brown-backed Indian Robin.

Abundant everywhere and a very common garden bird. A nest with three eggs found in a hole in a mound at the foot on a bush on March 22, 1925. Another nest built (no eggs) in a crevice in an old tomb March 29, 1925. Several pairs were seen towards the end of March carrying food to the young.

[Eggs taken by Bingham. March-June.]

Copsychus saularis saularis. The Indian Magpie-Robin.

A common bird on the Ridge and in shady groves and gardens; but not very prominent on account of its somewhat retiring habits. A pair used to haunt the grounds at Okla.

[Eggs taken by Bingham. May-June.]

Turdus atrogularis. (Temm.). The Black-throated Thrush.

I remember seeing two or three specimens of this species, which I know well from my experience of Simla birds, two or three winters ago in Delhi. Unfortunately no specimens were shot then; but a fine ♂ was obtained on February 14, 1925, in the thick babool clump where I met with *Pericrocotus b. brevirostris* a month earlier. Not common.

Monticola solitaria pando (Sykes.). The Indian Blue Rock-Thrush.

Not very common. A single bird or a pair may be seen among piles of stones and bricks. Only one ♂ obtained on November 13, 1924. A pair used to haunt the new Secretariat buildings at Raisina.

Ploceus phillipinus phillipinus (L.). The Baya.

Old nests seen on babool trees. A flock of about twenty birds seen on December 6, 1924, and two specimens obtained, an ad ♂ and unsexed bird. Three or four seen among long grass at the back of the Old Fort on February 7, 1925.

Ploceus bengalensis. (L.). The Black-throated Weaver-Bird.

The British Museum catalogue mentions a ♀, dated March 25, in the Hume collection.

Ploceus manyar flaviceps. (Less). The Indian Striated Weaver-Bird.

I did not get any specimens of this Weaver.

[Eggs taken by Bingham. July-September.]

Uroloncha malabarica (L.). The White-throated Munia.

Observed in small parties among long grass and thin babool jungle. This Munia is also very common among the thorny bushes on the Ridge, and seems to have a partiality for the kikar in which it builds its large globular nest. Cole found a nest in a kikar on November 14, 1924, with five slightly hard-set eggs. A note by Blewitt is quoted in *Nests and Eggs*. vol. 11, p. 139.

[Eggs taken by Bingham. February-September.]

Amandava amandava amandava (L.). The Indian Red Munia.

Seen in small parties among long grass along the canal banks at Kingsway and in other suitable localities where there is long grass. Three specimens obtained are all ♂♂.

Erythrospiza githaginea crassirostris (Blyth.). The Eastern Desert-Finch.

It is worth noting that this species has been obtained in the Gurgaon District near Delhi. Hume records the following note in *Stray Feathers*, vol. vii, p. 454:

'Mr. E. W. Cleveland sends me a beautiful specimen of *Bucanetes githagineus*, which he shot near Hattin in the Gurgaon District (Punjab) on December 16, 1877.

'No doubt the bird occurred here, as Mr. Cleveland remarks, as a mere straggler, but still its occurrence so far east is most remarkable. When I discovered it years ago in Sindh, this was an enormous extension eastwards of its range (which westwards stretches to the Canaries).

'Again, last year, when I shot it at Jodhpoor, we had a further easterly extension, and now this new locality extends the range to the 78° E. Long.'

Carpodacus erythrinus roseatus. (Hodgs). Hodgson's Rose-Finch.

A pair obtained on January 24, 1925, in thin babool jungle near the Jantar Mantar. The ♀ was first noticed perched on a low branch and was shot. I then made a search for the ♂ which I suspected must be nearby. I soon discovered him and shot him also. On January 31, 1925, I saw a party of about eight or ten birds feeding on the ground in a patch of babool trees, and secured another fine ♂. The only previous record of the occurrence of this species in Delhi appears to be ♀ dated April 2, 1872, in the Hume Collection—*vide* Mr. H. Whistler's 'Note on the Weavers and Finches of the Punjab.' (J.B.N.H.S., vol. xxx, p. 186).

Gymnorhis flavicollis (Frank). Yellow-throated Sparrow.

Observed first in a small party from which a bird was obtained. This was on November 13, 1924. Later on I got a solitary ♂ on February 5, 1925, and on March 7, 1925. I obtained two birds separately. On March 23, 1925, a pair was seen perched on railing on the roadside. Others were seen from time to time. This species is not at all uncommon in Delhi. The presence of this Sparrow may be detected in the field by the prolonged twittering notes of the bird which remains perched in one place for quite a long time while it keeps on twittering. The notes are like those of *Passer d. indicus* but softer and sweeter in tone. The colour of the bill in winter is variable: it may be either dark brown or black. Specimens collected appear to be intermediate between *transfuga*, and *flavicollis* and agree with examples obtained by Mr. A. E. Jones at Amballa. Please see in this connection Mr. Whistler's notes on this species (J.B.N.H.S., vol. xxx, p. 411).

[Eggs taken by Bingham, June.]

Passer domesticus indicus. (Jard and Selby). The Indian House-Sparrow.

Abundant. Resident.

Emberizinae. Buntings.

A small party of birds seen in the evening among the reeds at Okla on January 11, 1925, and apparently about to roost there, may have been Buntings of some sort. It was impossible to obtain specimens, however, and the species must remain in doubt. Although I watched for Buntings and visited all likely spots, I was unsuccessful in getting any specimens.

Emberiza cia per. (Hart.) The Transcasian Meadow-Bunting.

Cole obtained a ♂ on February 24, 1925, on the Ridge at Kingsway. This was the only one seen by him.

Riparia brevicaudata (McClell.) The Indian Sand-Martin.

Found breeding in colonies towards the end of November and in December, January and February. A pair obtained on November 16, 1924. Several holes were opened up on December 13, 1924, but held no eggs. Cole obtained eggs as follows:—

(a) February 2, 1925. Colony of eight nests. Three nests with four eggs each.

(b) February 6, 1925. Colony of four nests. Two nests held three and four eggs each.

(c) February 26, 1925. Four nests, all with young.

[Eggs taken by Bingham. January-May and October-December.]

Ptynoprogne concolor (Sykel.) The Dusky Crag-Martin.

Some Crag-Martins, which, I think, must have been of this species, were seen flying about near the Old Fort on February 7, 1925.

Hirundo rustica gutturalis. (Scop.). The Eastern Swallow.

Seen flying over grassy plots along with *laurica* Swallows and *Riparia brevicaudata* towards the beginning of November. Later on these Swallows were not seen so prominently. An ad. ♂ and an unsexed juv. were obtained on November 10 and 11, 1924, respectively. In the field this species is easily recognized by its pinkish underparts and the dark patch on the throat. The specimens obtained seem to be intermediate between typical *rustica* and *gutturalis*.

Hirundo daurica erythropygia. (Sykes.). Sykes's Striated Swallow.

I found a nest with young on March 14, 1925. in a small ruined tomb in babool jungle at Raisina. In the dome of the same tomb several pairs of *Micropus a. affinis* were also nesting. A single egg was once brought to me by a small boy.

[Eggs taken by Bingham. March-June and September.]

Hirundo daurica nepalensis. (Hodg.). Hodgson's Striated Swallow.

A specimen (unsexed) obtained on March 8, 1925. On which date three or four were seen perched on telegraph wires along the Bela Road. It is not easy to differentiate between this sub-species and *erythropygia* without shooting specimens.

Hirundo daurica striolata (Temm. and Sch.). The Japanese Striated Swallow

On the morning of November 16, 1924, I saw numbers of Swallows of the *daurica* type perched on telegraph wires at the back of the Secretariat in Old Delhi. An ad. ♂ obtained has the striations on the lower parts very coarsely marked. It is also a large bird. It appears to be a specimen of this race, though the distribution does not normally extend so far west. Mr. A. E. Jones has compared this bird with a fair series of the commoner *daurica* Swallows in his collection, but cannot match it.

This specimen has very kindly been sent by Mr. Whistler to Dr. Ticehurst for identification.

Measurements. Wing 122 mm. Tail 120 mm. Depth of fork 68 mm.

Hirundo smithii filifera (Steph.). The Wire-tailed Swallow.

This species builds under culverts and bridges above water. Some winters ago I saw nests at Kingsway. I was told of some Wire-tailed Swallows seen on the telegraph wires along the Bela Road, but I visited the place two or three times without success. No specimens were collected. Some birds were seen on April 3, 1925.

[Eggs taken by Bingham. April-May.]

Motacillidæ. Wagtails.

Wagtails were exceedingly common when I first arrived in Delhi, and continued to be so till the middle of February. I made the mistake which seems to be difficult to avoid, viz., omitting to collect a sufficient number of specimens of a common species till one realizes it is too late to make up for lost ground. The result was that I got hardly any Wagtails. The only species I am able to record with certainty are enumerated below. From about the middle of March large numbers of Wagtails are to be seen flying over on migration in the evenings. Hundreds of birds may be watched passing over in a continuous stream. The direction appeared to be N. W. or W. N. W.

Motacilla alba dukhunesis (Sykes.). The Indian White Wagtail.

This was the commonest species during the winter. It remained behind till after *M. feldegg melanogrisea* had left. Stray birds were seen in grassy plots and open ground till the very end of my stay in Delhi, though no other species were observed except in flight on migration. A specimen obtained on March 21, 1925, is in very heavy moult.

Motacilla maderaspatensis (Gmel.). The Large Pied Wagtail.

The classification of this Wagtail in the *alba* group seems incorrect. I am inclined to agree with Dr. Ticehurst that, for the reasons given by him (J.B.N.H.S., vol. xxviii, p. 1090) this species should be kept apart from the White Wagtails. I did not see this large Wagtail anywhere else except at Okla and at Kingsway. In the latter place, I found a nest under a low bridge near Shah Alam's Tomb, on February 27, 1922. The nest held three fresh eggs which I took.

Motacilla feldegg melanogrisea (Hom.). The Black-headed Wagtail.

This species was common enough up to the middle of February. Birds were generally to be seen in open spaces, in fields, near patches of water and in gardens. In winter most birds are a dingy, brownish colour; in March some green begins to show on the upper parts and the lower plumage shows indications of turning yellow, and by the end of March numbers of birds in full

Motacilla feldegg melanogrisea—(contd)

breeding plumage are to be seen. This Wagtail perches freely on trees and telegraph wires. The note is a penetrating *queegik, queegik*. A ♂ specimen shot on February 6, 1925, is beginning to assume the summer plumage, and a fine ♂ shot on March 8, 1925, is in full breeding dress.

Anthus trivialis trivialis (L.) The Tree-Pipit.

Common throughout my stay. One greenish specimen appears to be of this race, not being near as green as typical *maculatus* with specimens of which it was compared.

Anthus campestris griseus. (Nicoll). The Eastern Tawny Pipit.

The first specimen was obtained by my brother who shot a ♂ on November 10, 1924. The next day he shot two more. This species is to be found in open country covered with short grass, on large open *maidans*, lawns, etc. On February 7, 1925, I saw some Pipits which were probably of this species on the lawns in the Old Fort. The specimen shot on November 11, 1924, is the palest obtained, while the ♂ shot on the previous day is inclined to be rufous on the upper parts and tail. The colours of the soft parts are as follows :

Irides—brown.

Bill—Upper mandible brown to brownish-black, lower mandible fleshy.

Legs and feet—fleshy.

Claws—pale horny or fleshy brown.

Anthus richardi rufulus (Vieill). The Indian Pipit.

On March 7, 1925. I obtained a Pipit (which could not be satisfactorily sexed) which has a whitish throat and is of smallish size. This bird I could not identify. Mr. Whistler has been good enough to examine it carefully and is satisfied that the bird belongs to this species. In this specimen the wing measures 85 mm., and the hind-claw 8.5 mm.

Anthus sordidus subsp. (Finsch.) The Brown Rock-Pipit.

This large Pipit first came to my notice on November 29, 1924, on which date a ♀ was shot for me. No more were seen for some little time. On January 6, 1925, I was taken to a place where some more birds had been seen. Although *Anthus hodgsoni* was found in numbers, none of the larger species was met with. On February 14, 1925, I discovered a spot where there were several birds. This was a fairly open grass-covered stretch, interspersed with thin babool jungle and jherberry bushes. About half a dozen Pipits were seen, and a specimen (which could not be sexed) was obtained. On February 17, 1925, I again visited this place and obtained two more specimens—both ♂♂. I saw this Pipit from time to time till the end of my stay. I would not regard this bird as rare in Delhi. The fact of the matter is that it frequents country of the kind described and is therefore not usually seen except in such haunts. It is always found on the ground, rising suddenly from near one's feet if not noticed ; but running quickly along the ground, taking cover behind tufts of grass, clods of earth, etc., if followed up. It frequently perches on large exposed rocks, and also on bushes and trees, I did not, however, observe it wagging its tail in the slow manner of *Anthus t. trivialis*. The note is a soft *plip, plip*. The colours of the soft parts are as follows :—

Irides—dark brown.

Bill—Upper mandible dark horny brown, lower mandible pale yellowish-white with dusky tip.

Legs and feet—pale brownish-yellow.

Claws—brown.

The ♀ shot on November 29, 1924, has more rufous in the lower plumage than the other specimens, and the unsexed bird obtained on February 14, 1922, has the dark-brown streaks on the breast most distinct.

Wing 95–99 mm. Tail 84–86 mm. Hind-claw 9–10 mm.

Calandrella. Short-toed Lark.

On three or four occasions I met with flocks of some small birds in bare, open country, where such species as *Pterocles s. erlangeri*, *Cusorius coromandelicus*, etc., are found. The birds were exceedingly wary, however, and I could never get close enough to shoot any with my .22 bore or collecting gun. Flocks were also met with in ploughed fields. The birds seemed rather

Calandrella—(contd.)

restless, wheeling about and settling on the ground to feed, then suddenly flying off again and after a few wide circles settling on the ground once more. These little birds were probably *Calandrella* larks.

Alaudula raytal raytal (Blyth.). The Ganges Sand-Lark.

Cole obtained a specimen near the riverside on November 26, 1924.

Mirafra erythroptera (Jerd.). The Red-winged Bush-Lark.

This Bush-Lark is found commonly all over Delhi. It is generally seen singly or in pairs, crouching on the ground near a tuft of grass, a bush or clod of earth. When disturbed it often flies on to a tree or bush.

[Eggs taken by Bingham. One nest found on the Ridge on September 21.]

Galerida cristata chendoola (Frankl.). Franklin's Crested Lark.

Common in open country, in fields, etc. Has rather a plaintive note.

[Eggs taken by Bingham. Two nests, on March 31 and April 23.]

Pyrrhuloxia grisea grisea (Scop.). The Ashy-Crowned Finch-Lark.

Common in dry, sandy ground. Seen on the footpaths of the more open roads on the outskirts of Raisina and on bare ground near the new Secretariat. Three ad. ♂♂ obtained on March 14, 1925. One of these is curiously deformed. There is no foot on the left leg, which terminates in a hard, horny process! This bird did not apparently suffer much on account of its malformed foot, for it moved about on the ground just as easily as other birds and it was not until it was shot that the deformity was noticed. Two ad ♀♀ secured on March 18, 1925, in front of the new Secretariat at Raisina. A juv. bird was caught alive on March 17, 1925. It lived for a day and I made it into a skin when it died. Mr. S. C. Law has given a full description of the song-flight of this peculiar Lark (*Ibis*, 1924, pp. 645-647). A low, churring note is also uttered when the bird is seated on the ground.

Zosterops palpebrosa elwesi (Baker). The Western White-eye.

Heard throughout my stay. No specimens obtained, but the birds must have been of this race.

[Eggs taken by Bingham. May-July.]

Cyrtostomus asiaticus asiaticus (Lath.). The Purple Sun-bird.

Abundant. The very first bird collected on the day of my arrival was a ♂ in undress plumage. A series of seven ♂♂ was obtained with the object of ascertaining whether the purple dress is a seasonal change of plumage assumed during the breeding period, or whether it is merely the fully adult garb acquired after some time, and which when once assumed, is not lost again. A careful examination of the skulls of the specimens I shot showed them to be adult birds. I also noted carefully that all ♂♂ seen during the months of November-December and the first few days of January lacked the full purple plumage; whereas, several ♂♂ were in purple plumage by the first week of February, and by the middle of that month hardly a specimen was seen in undress. The first purple ♂ was seen as early as January 4, 1925; but along side it, on the same flowering shrub, was another ♂ in undress. This leads me to think that the purple plumage is seasonal. There is a note on this subject by Dr. Ticehurst in the *Ibis* (1923, p. 24). Breeds commonly in March. Nests usually suspended from the thorns on the fleshy leaves of the cactus. Of six ♂♂ measured, three have the bill 19 mm., and three 20 mm. (Measured from gape).

[Eggs taken by Bingham. March-May.]

(To be continued)

TAKIN SHOOTING IN THE SPRING

BY

E. MAXWELL WEST

(*With a plate*)

It is four o'clock in the morning and bitterly cold as you force yourself half out of the blankets to drink your morning tea. That biting wind, which springs up just before dawn to herald the sun, is whistling through the thin walls of the bamboo hut and you wonder why you ever took up big game shooting and resolve, on your return to civilization, to sell your rifles and go in for poodle-faking or some less exacting pastime which does not involve living in the 'blue', rising by lamp light and being frozen withal.

Camp is at 11,000 feet, and at this hour, as you tramp up the hill, the deodar and rhododendron forest with its undergrowth of bamboo all festooned with moss and with drifts of snow in every hollow presents a dismal picture. Presently, however, as the first rays of the sun strike through the interlacing branches, the scene is transformed, and, incidentally, your former depression vanishes.

From a dark and gloomy vista of greys and blacks you find yourself, as if by magic, in an enchanted forest. The dawn brings out the scent of the rhododendrons and magnolias which carpet the ground with their fallen blossoms in patterns of pink, scarlet and white. A thousand points of light shine from the dew-drenched greenery, and the moss, which previously had such a depressing effect, now appears like hangings of jewelled lace.

The wind has now died away and the breathless stillness which prevails is suddenly broken by the sound of, surely, a flute. As you halt the notes are repeated and amid a shower of sparkling dew, a Tragopan drops from his roost in the trees, a vision of grey, spotted crimson and blue. His upright blue horns and flute like call remind one irresistibly of Pan and his pipes, but this is a Temminck's Tragopan, only to be met with on these far frontier hills, and were it not for the thought of possible Takin higher up, sentiment would be dismissed and he would be added to the bag. The Yawyin guide, however, cannot understand your forbearance and by his excited gestures urging you to shoot, soon removes any chance of your doing so by frightening the bird away.

As you reach the higher levels the forest thins out, the rhododendron, magnolia and bamboo being replaced by larch, juniper and dwarf rhododendron, while blue gentian stars the more exposed spaces from where the snow has melted. Again you are startled by a strange call and catch sight of a smallish bird in a livery of apple-green grey and scarlet, as he runs up the hill. This is the Blood Pheasant, a beautiful bird, but from the sportsman's point of view annoying, as he never rises and has to be shot running.



TYPICAL TAKIN COUNTRY



THE END OF A PERFECT MORNING

At last you reach your objective, the open grassy slopes from which the snow has almost all melted, and where you hope to find the Takin. Here you stop for a well-earned rest and to search the hills around with glasses, but before you sit down pull on the sweater which your guide has carried, for it was hot work climbing that last five hundred feet, and here, at over 12,000, in spite of the sun, you will soon be chilled.

It is a wonderful scene. A wide semi-circle of black cliffs picked out with patches of dazzling snow show up clear cut against the brilliant blue of the sky, sinking towards the centre to a col, which is the pass into China. The grassy slopes you are on are relieved here and there by clumps of deodar and pine in black silhouette on the unmelted snow, while beneath stretches the dark forest from which you have just emerged. Far below again, in the green jungle clearings are a few scattered hamlets of the Yawyins, the mountaineers of this frontier.

As you sit here in the midst of Nature's grandeur drinking in the champagne-like air, you wonder at your heretical thoughts of two hours ago, and pity the poor misguided fools who spend their leave in a round of gaiety, long nights and short drinks, at popular hill stations. This feeling of exaltation of being removed from the world of dull care and monotonous routine, of being in part with Nature, is worth living for. You feel that God is near

A rattle of stones above you breaks into your reverie. You slowly turn your head, and behold, the Devil? . . . No! at long last you have found the Takin, or rather he has found you, for suddenly coming over the ridge above you he sees something strange and stops to stare. He is not unlike the Devil with his large Roman nose, sharp black horns and large cloven hoofs.

Not a move now or he will be off. After what seems an age he turns his head to look for his companions. Now is your chance. Slowly you bring the rifle forward and get a sight on to the point of his shoulder, when he again turns to stare at these strange intruders. His curiosity is his downfall and as the report of your rifle is thrown back from the cliffs above, your first Takin lies kicking on the ground. After firing it is as well to jump to one side behind a rock if possible, as the Takin is reputed to live up to his appearance and charge anyone in sight when wounded, his business like horns, his size and considerable agility making him an animal not to be despised on a steep hillside.

This time however the bullet has done its work cleanly and well, and you scramble up the hill to view your trophy, but first take cover behind that rock and see if there is anything the other side of the ridge. Yes, not far below are three more of these strange animals, one good bull and two small ones. They have never heard a rifle shot before and do not know from which direction the danger comes, but steady now, and get your breath, for you may never see a Takin again. Once more the rifle shatters the stillness and the big bull drops, the youngsters disappearing at an amazing pace into the jungle below.

Forgotten are the weeks of fruitlessly toiling up and down the hills as often as not in snow and rain, with nothing to gladden

your eyes but old tracks. Or if remembered for a moment you dismiss the thought as you look down on your hardly won trophies and around at the glorious panorama, now rendered doubly beautiful by your success.

And so we will leave you with your measuring tape and camera, and to enjoy a well-earned breakfast in the finest surroundings for which man could wish, while your guide drops down to the camp to bring up men for the final obsequies.

BIRDS NESTING WITH A CAMERA IN INDIA

PART VI

BY

CAPT. R. S. P. BATES

(Continued from page 804 of Volume XXX)

(With 14 plates)

In the plains of India the lot of the ornithologist on most occasions is far from being all that can be desired. From April till the end of September, the months of greatest activity amongst the resident feathered population, the discomforts of first the scorching heat of the early summer, next the humid and sultry days of the rains with their attendant and everchanging hordes of noxious insects, and lastly the final burst of heat before the cool weather sets in, are at times well nigh unbearable.

By those who have never braved the inside of a small hiding tent when the shade temperature is in the neighbourhood of 110° , or have never had occasion to spend hours at a time in a small open boat on a shadeless expanse of water, the attendant discomforts cannot be realized. For these reasons and the fact that I have only been able to devote myself to this work during the all-too-short and far-between periods of leave obtained during the late unsettled years, I fear my efforts with regard to the birds of the plains and the hills of continental India are most incomplete, even for the comparatively short time I have spent in the country.

However, though my photographic records are somewhat scant, I have indeed had the fullest value out of my exploits, and on going through the notes jotted down for the subject matter of this article, I find that, rather than impressions of the general habits of the birds dealt with, they are largely records of abnormalities, curious nesting sites, etc., and experiences at times annoying and unpleasant but all the same amusing when one looks back on them. This chapter therefore might more aptly have been headed 'Abnormalities of Indian Bird-Life' or 'The Foolishness of being an Ardent Ornithologist'.

I first conceived the idea of making bird photography my hobby when stationed in a native state in Rajputana, with rather more leisure than one now has in the post-war army. Peafowl were an absolute nuisance and being sacred one was not allowed to impress upon them with a charge of shot that they would be the more welcome if only seen and not heard, with the result that they were so tame that they even used to come into the compound, and, if anything disturbed them during the night, one would be rudely awakened by piercing yells from their roosting places in the large trees around the bungalow.


I discovered one morning that a Peahen had a nest in some long

grass just the other side of a privet hedge and not more than 20 yards from the bungalow, so I sallied forth with all my paraphernalia to photograph it, and having set up the camera stepped behind it to start focussing. Unfortunately I put my foot in the very middle of a bees' nest, nasty little grey fellows which stuck to one like leeches. The camera flew in one direction, and I in a bee-line in the other. It really must have been a bee-line as they too followed it, and incidentally me, in most unpleasant numbers. The remainder of that day I was somewhat unhappy and only ventured forth to collect my apparatus after sundown. A couple of days later I summoned up courage to try again, and approaching the nest from a different direction was disgusted to find nothing but some broken egg-shells.

Curiously enough, one of the first nests I found in Bharatpur, the said native state, was on the roof of the Bungalow and was that of a Brown Rock-chat (*Cercomela fusca*) whose confiding ways and sweet subdued little song soon attracted my attention. The nest was in a curious spot, being built inside an old nest of a Syke's Striated Swallow, the tubular entrance to which had somewhat crumbled away. I succeeded in getting quite a pleasant photograph including the bird standing on the ledge above the nest. Later on the same pair made two attempts to raise a second brood: the first of the two quite close to the old site in a crevice in the wall below a window sill, and the second on a shelf actually inside the lamp room, a small room at the back of the Bungalow with a single barred window without glass. When this latter nest was almost complete, some ass of a servant swept it away. These confiding little birds were very common in Bharatpur, and as they have a very misguided preference for human propinquity, almost every bungalow had its lodgers. The nests, I noticed, were rather untidily put together, and had little in the way of lining. The eggs were in no way unusual, being pale blue green with small reddish spots mostly around the larger end; unspotted eggs I never came across. I wonder why this chat should ape Robin Red-breast? Perhaps it is because it rather resembles it in plumage, much more so, in fact, than do the Indian Robins, and having realized this fact and seeing that the Robins are far from familiar, it may think that it is up to it to do its best to fill the gap in the heart of every new arrival from Home who longs for anything English.

The Rock-chats were not alone in choosing weird places for their nests. Stranger than the above was the situation of a Yellow-throated Sparrow's (*Gymnorhis xanthocollis xanthocollis*) nest. This little bird is just what an aristocratic House Sparrow should be but is not--well mannered, quiet, and not too familiar.


It approximates to the female of that species in appearance, but has a rather long and slender bill, the distinctive pale yellow throat-patch, and for nesting sites largely patronizes holes in trees and, although found round about habitations, is also addicted to scrub jungle and fairly open country. Still, one of the strangest of nesting sites was that chosen by a pair of these birds close to the house. An iron gate, seldom closed, and situated but some



THE BROWN ROCK-CHAT
(*Cercomela fusca*)



THE LARGE GREY BABBLER
(*Argya malcolmi*)



THE CEYLON RED-VENTED BULBUL
(*Molpastes hæmorrhous hæmorrhous*)



THE SOUTHERN INDIAN STONE-CHAT
(*Saxicola caprata atrata*)



THE WHITE-SPOTTED FANTAIL FLY-CATCHER
(*Rhipidura pectoralis*)



THE SOUTHERN RED-WHISKERED BULBUL
(*Otocompsa emeria fuscicaudata*)



THE NILGIRI LAUGHING THRUSH
(*Trochalopteron cachinans cachinans*)

15 yards or so from my bed-room door, led from the side of the compound into a field. The gate posts were of stone and through one of them a hole had been pierced for the bolt. This was certainly not more than $1\frac{1}{2}$ inches in diameter on the outside and I very much doubt if it got any bigger inside, and certainly there did not appear to be room for the birds to turn, as I noticed that they invariably reappeared from the hole on the opposite side to that by which they had entered. This may, of course, have been merely due to habit. Unfortunately I find I have made no entry in my diary as to whether their efforts were crowned with success or not, and I cannot now remember what happened.

Only last year I struck another of these nests in nearly as strange and certainly a more unpleasant situation, namely in the top of a metal telegraph post on one of the rifle ranges near Madras. The diameter of the apex of a similar post lying close by, which also lacked its lid, I judged to be just about 2 inches. There were evidently hefty youngsters in the nest nearly ready to fly. It had been constructed some few inches down from the top, and how both the parents and young stuck it I don't know. The heat of the post at 8 a.m. was infernal, so what it must have been like by the time the sun had been playing on the metal for some hours I cannot possibly imagine.

However to return to Bharatpur, a further peculiar Yellow-throated Sparrow's nest was one in the old nesting hole of an Indian White-breasted Kingfisher (*Halcyon smyrnensis fusca*), an abode some two feet deep. This nest I opened up as it was only about nine inches down from the top of a sandy bank. There were four eggs in a small cup in a large and rather loose untidy collection of grass and feathers, the latter material greatly predominating. After photographing it, I roofed in the hole again with small slabs of stone which I discovered near by, covering these up finally with earth and sand, and had the satisfaction of seeing the birds carrying on quite happily. Some days later, however, the nest was raided and the eggs eaten, either by rats or some other of the many pests with which Indian birds have to contend.


More confiding than the above were the Bulbuls, two species of which lived in the compound, namely the Central Indian Red-vented Bulbul (*Molpastes haemorrhous pallidus*) and the White-eared Bulbul (*Molpastes leucogenys leucotis*). As the sun used to set behind the house, in front of which was a small lawn with a *nim* tree in its centre, I always had my tea out of doors, and it was not long before I had collected one or two acquaintances, the first to become thoroughly tame being a pair of Bulbuls of the former species, which soon took to descending on to the tea-table to peck at anything and everything which looked eatable. A White-eared Bulbul used to hang about the tree, but would not actually join in the repast except when I left the table and went into the verandah. Though I did everything I could to gain their complete confidence, I never succeeded in getting any of them to actually feed out of my hand, and had to console myself by ministering to the wants of a bold little tree-rat, who used to run up my leg to pinch bits of cake from off my knee.

These two species of Bulbuls were excessively common, and one was always coming across their nests in almost every description of bush or sapling, the small prickly bushes at the back of the bungalow being especially favoured by the White-eared variety, which in Bharatpur very much outnumbered the Red-vented species. In the construction of their respective nests there was nothing to choose, both being pretty rottenly made thin-walled cups of bents, in comparison to which the nests of the Southern Red-whiskered Bulbuls in the illustrations here are most solid structures. It really does seem extraordinary that these cheery little birds should be so common when one thinks of the astonishing number of nests which are destroyed, many through the bird's own stupidity I'm sorry to say. I soon discovered that if I found a particularly interesting nest, even to put off photographing it till next day was exceedingly risky, the only safe thing to do being to return for the camera then and there.

Before long my tea parties were joined by a band of noisy seven-sisters—Large Grey Babblers (*Argya malcolmi*), and their advent might be termed the beginning of the end, as quarrelling at once became the order of the day between these people and the tree-rat family, which resulted in the complete discomfiture of the cheerful little Bulbuls, who evidently disagreed strongly with such unseemly behaviour. In the north these Babblers are exceedingly tame, whereas here in the south, and especially in the hills, these same birds particularly shun observation. In fact wherever I have struck them in the peninsula they cannot possibly be described as anything but rather shy skulking birds, which, though given to entering gardens and orchards, certainly resent being watched. I do not consider, however, that they are as shy as Mr. Douglas Dewar leads one to expect in his 'Birds of the Indian Hills'.

Besides this Babbler, the Common Babbler (*Argya caudata caudata*) used also to frequent the compound, and, in a thick tangle of creeper-like growth enveloping a piece of the fence bordering the railway line, I came across a compactly built nest, made chiefly of the thinner strands of the same material in which it was placed, and containing three eggs of the usual very glossy texture and deep spotless blue colouration.

To expound fully on every description of bird which was noted in the compound would fill a volume, but even a bare list of those others not yet mentioned, which I noted as of frequent occurrence during the whole or a large part of the year, is of decided interest, including as it does such widely divergent forms as Quails and Partridges (the Black and the Grey). Ibis, Nuktas, Spotted-billed Ducks, Sandpipers and Stilts, Paddybirds and Night Herons, during the floods, Paroquets, Hornbills, Tree-pies, Bee-eaters and Fly-catchers—the Paradise Fly-catcher and the White-browed Fantail—Rosy-backed Starlings and Mynas of various species, Sun-birds, Barbets and Golden-backed Woodpeckers, Crow-pheasants, Koels, Crows, Shrikes and King-crows, Munias, Weaver-birds and Wren-Warblers, Bush-chats, Finch-Larks, Pipits, Hoopoes, Rollers, Doves and Blue Rock Pigeons, and probably many others which at the moment I do not recall, and, of course, Kites, Scavenger



THE WHITE-EARED BULBUL
(*Molpastes leucogenys leucotis*)



THE CENTRAL INDIAN RED-VENTED BULBUL
(*Molpastes hamorrhous pallidus*)



THE SOUTHERN INDIAN BLACK BULBUL
(*Microscelis psaroides ganeesa*)



THE SOUTHERN RED-WHISKERED BULBUL
(*Olocompsa emeria fuscicaudata*)

Vultures, an occasional White-backed Vulture and, on a couple of occasions at least, a King Vulture.

I don't suppose for an instant that this extraordinary variety of species could have been met with in any other garden in Bharatpur. It was largely due, of course, both to its constitution and to its position, as it was something like a mile from the city and considerably farther from any other bungalow, with the exception of the Moti Jhil, and had scrub jungle in front and to one side, and on the other side and rear cultivation. One face of the compound was bounded by a railway embankment, to construct which deep pits had been dug, and these, besides being full or overhung by masses of vegetation, connected up into a long stretch of water during the floods. Babool trees and scrubby bushes were also scattered about throughout the compound and between its rear boundary and the main broad-gauge railway line which also ran pretty close. In some places, too, long grass obtained. Lastly, the bungalow was situated on the direct line between the Keoladeo Ghana and large Jhils of like nature at Dig and Kama; hence waterfowl of all kinds and descriptions were continually passing over.

The remaining birds whose nests I actually took in the compound were the Indian Ring-Dove, the Brown-backed Indian Robin, which in true eastern fashion puts the cart before the horse in that it carries its red patch under its tail instead of on the breast, the Crimson-breasted Barbet, with whose monotonous call every dweller in the plains is only too familiar, and the energetic little Purple Sunbird.

A pair of Indian Ring-Doves (*Streptopelia decaocta decaocta*) started to build their apology of a nest on a rafter in the verandah. I say started advisedly, as at one point I really despaired of its ever-reaching completion. It did however get to the egg stage, though no further. I noticed a most curious fact about this pair of Doves—perhaps only one of them was responsible for the nest construction; I don't know, but hope so. I hate to think they were both equally brainless—and that was that any material which dropped to the floor was never retrieved, and I am sure I'm correct in saying that those fool birds dropped enough twigs, in trying to balance them in impossible places, to build at least four of the nest which was eventually the outcome of their unnecessarily hard labours. This and the Little Brown Dove (*Streptopelia senegalensis cambayensis*) were the two species common to Bharatpur, and they were excessively common too. I don't think I ever saw the Spotted Dove (*S. chinensis suratensis*) there, though it seemed numerous enough a bit further south in Kotah, and seems to be the commonest Dove of all the places of South India in which I have been stationed, as well as of the Hills.

The Indian Crimson-breasted Barbet (*Xantholæma hæmacephala indica*) bores a hole about the size of a half-crown in the trunk or a stoutish branch of any tree up to about a foot in diameter. Near the extremity of a broken branch or of one that has been sawn off and has commenced to decay is perhaps the most favoured spot. It is generally on the under side too, that is if

the branch slopes at all. On one or two occasions I have noticed holes on the underside of perfectly horizontal boughs.

In Secunderabad I watched one of these birds excavating its abode. The spot chosen was only about 10 feet from my bedroom window, so I was able to lie on my bed and watch it at ease. When I first noticed it, the hole was only about half an inch deep, so I could see that the bird worked with the bill slightly open, and that after every few taps it would jerk the chip or chips thus gouged out over its left shoulder. It worked most intermittently, sometimes only for half a minute or so at a time, and often with intervals of some hours in between. Nevertheless, in three days only the latter half of its body remained in sight and it had commenced to turn the cavity downwards. The branch, incidentally, was perfectly healthy and was that of a *nim* tree. The manner in which this quaint little bird arrives at its full-blown 'song' always amuses me. It apparently needs to work itself up to the necessary pitch of excitement and usually starts with a very subdued 'wuk', which steadily increases in volume, at the same time changing slightly in tone, till the rounder and not unpleasant even 'tonk' is attained, and thereafter monotonously reiterated.

The nesting of the Brown-backed Indian Robins (*Saxicoloides fulicata cambaiensis*) has nothing of particular interest about it, excepting perhaps that they have the same habit of choosing strange sites as has the English Robin. One nest I found was in an old *ghurra* which had been thrown into the centre of a bush, and another in a battered kerosine oil-tin lying in a patch of weed. Those I found in my Bharatpur compound were:—two in cavities in the railway embankment, one in the roots of a low prickly bush and a couple more in holes in the back wall of the stables. At the present moment I have a nest full of youngsters a few feet above my head in a pipe intended to drain rainwater from off the roof, but this, of course, is of the Black-backed variety which replaces the Brown-backed Robin in the southern half of India.

The Indian Purple Sunbird (*Leptocoma asiatica asiatica*) is a highly interesting personage as its nest is such a well thought out structure. It is, of course, known to every dweller in this country on account of its great predilection for flowers and gardens, its unbounded energy, sharp call notes, and habit of ceaselessly flicking the wings, as it turns and twists about the flower stalks in its efforts to extract the honey and insects from the innermost recesses. The male in the breeding season is also possessed of quite a sweet little song. A pair suspended their nest from a rafter in the verandah, but for some reason or other I failed to spot it until the eggs had been laid and the very day following its discovery it was torn down and its remains strewn about the floor, probably by a tree-rat in search of material for its own ball of rubbish stuffed into a corner by a rafter a few feet distant. In Karachi I took a photograph of a nest of this species suspended from the telephone wire close to one of the insulators on the mess wall. Occasionally they are very well concealed, but whether by intent or accident I do not know. Nevertheless it is a most noticeable fact that a large proportion of



THE DUSKY HORNED OWL
(*Bubo coromandus*)



THE SMALLER WHITE SCAVENGER VULTURE
(*Neophron percnopterus ginginianus*)



THE INDIAN PURPLE SUNBIRD
(*Leptocoma asiatica asiatica*)



THE MALABAR SMALL MINIVET
(*Pericrocotus peregrinus malabaricus*)

the nests to be found in and around Secunderabad are constructed within large closely woven untidy webs, which abound in the camel-thorns. The result is that, failing the presence of the bird, most of them are extraordinarily liable to escape detection. The usual impression conveyed by most descriptions I have read of the nesting of this bird is that a grass nest is first constructed and on to this all manner of decorations in the form of spiders' webs, chips, bark, scraps of paper, etc., are then stuck with an idea of either making it pleasing to the eye or more probably to camouflage it. Incidentally I once remember seeing a nest outside a regimental office door plastered with many scraps of white paper. I don't for a moment wish to assert that this is incorrect. To do so on the strength of observations on a single nest would be absurd, and I have only once witnessed the construction of a nest from its very commencement. However, this was so diametrically opposed to the above conception that it is certainly worthy of notice.

On the 24th of July I was sauntering round the compound of my bungalow on the outskirts of Madras when my attention was suddenly attracted to a Purple Sunbird, a female, which flew down to a very low date-palm and seemed to have become greatly interested in a point a few inches from the extremity of a drooping frond. On looking at the spot, but some 3 feet from the ground as the palm grew at the foot of a bank, I at first noticed nothing at all unusual, but on going closer I discovered a minute piece of web had been wound round the stem with a loose end perhaps half an inch in length left hanging down. Five days later the nest had progressed to the extent of being in shape not unlike the upper half of a crinkled paper-bag suspended from the branch, or a small edition of an unfinished Weaver-bird's nest without the cross bar, but of grass there was no sign. It was but a scant collection of the usual materials found on the outside of these nests. On the 4th of August the outer shell was completed and reminded one of nothing so much as of a deflated penny balloon, the entrance hole appearing like a rent in its side. The next step appeared to be the construction of the porch and by the 8th this and the outer shell were completed even to the ragged little bits which can be seen hanging down in the photograph, but of inner grass nest there was still no sign. In the making of the porch, short lengths of grass had been employed and from this time on grass was the main item brought by the builder. During the next week the balloon was quickly inflated, the bottom even on the afternoon of the 9th presenting a more or less rounded appearance, and, on looking into the nest on the 16th, I found two eggs had been laid. I do not know for certain on what days these eggs were deposited as for three or four days previously I unfortunately omitted to look inside. At any rate, the building undoubtedly took a full three weeks. I find in my diary for the 30th a remark to the effect that the Sunbird's eggs appeared to have been hatched about two days, and later, that they left the nest either on the afternoon of the 15th September or early morning of the 16th, as they were there after lunch on the former date, but had left it when I again visited the nest after breakfast the following day. The whole operation

therefore took 54 days, made up approximately as follows :—nest building 22 days, incubation 14, and rearing 18. On the 11th September my bearer brought to me a half-naked young Sunbird of apparently the same species, which he had found lying plumb in the centre of the main road. I suppose it had been dropped by a crow or something, but nevertheless it showed no signs of injury. Taking pity on its sorry plight, I popped it into my Sunbird's nest where it was at once adopted, and, though very much younger than its foster brothers or sisters, it thrived, and was comfortably ensconced in sole possession on the 16th. How long it remained there I do not know, as I did not go near the place again for some days.

In strong contrast to the tedious nest-building of these Sunbirds were the efforts of a pair of Malabar Small Minivets (*Pericrocotus peregrinus malabaricus*), which started building in the same compound on the 19th of July. So quick were they that by the 29th the full clutch of 3 eggs which the nest then contained showed distinct signs of incubation. The nest was certainly completed by the 25th, as the female was on that morning observed to be seated on it for an unbroken period of at least 3 hours, i.e., from 7 a.m. till I left the house at 10 a.m.

Previous to this the first hours of daylight had been spent in energetic nest-building, both birds flying to and from the site at very short intervals. Operations seemed to get gradually slower as they went on and appeared to cease entirely about 4 p.m. each afternoon. From where I slept on the verandah I could look straight across to the *nim* tree, some 40 or 50 yards distant, on one of the lower and smaller branches of which the nest was constructed at about 12 feet from the ground, and I had therefore ample opportunities of watching their early morning antics at the nest. From below and even from slightly to one side the nest was absolutely invisible, being but a tiny shallow cup of a small quantity of fine grass cemented together with webs and lichen and decorated with bits of dead leaves and bark-scrap, the whole barely the thickness of the nearly horizontal branch on which it rested and with the colouration of which it blended perfectly. To be exact its extreme diameter was just under $2\frac{1}{4}$ inches, and I should think the whole affair could have been squashed up into a little ball the size of a small walnut, so it is hardly surprising that the Minivets somewhat outclassed the Sunbirds in the matter of speed of house construction.

From amongst the other birds mentioned in the list a few pages back, I took in close proximity to the bungalow the nests of the following :—The Smaller White Scavenger Vulture (*Neophron percnopterus ginginianus*), Nukhta (*Sarcidiormis melanota*) White-browed Fantail Flycatcher (*Rhipidura aureola aureola*), Northern Golden-backed Woodpecker (*Brachypternus aurantius aurantius*), Bay-backed Shrike (*Lanius vittatus*), and Black Drongo (*Dicrurus macrocercus macrocercus*) White-throated Munia (*Uroloncha malabarica*) Baya (*Ploceus philippinus philippinus*) Indian Wren-Warbler (*Prinia inornata inornata*), the Ashy Crowned Finch-Lark (*Pyrrhuloxia grisea grisea*) and Indian Pipit (*Anthus richardi rufulus*).



THE SOUTHERN JUNGLE-CROW
(*Corvus coronoides culminatus*)



THE NILGIRI BLACKBIRD
(*Turdus merula simillimus*)

Besides Pharaoh's Chicken, the Common Vultures of Bharatpur were the Indian White-backed Vulture (*Pseudogyps bengalensis*) which outnumbered any of the other species by at least thirty to one, the Black Vulture (*Torgos calvus*), and the Indian Griffon Vulture (*Gyps fulvus fulvescens*). Whenever I shot a Black-buck or Chinkara, after having first removed the head and skin and any portions we required for the larder, I used generally to sit down more or less under cover about 40 or 50 yards away from the carcase just to see what birds and beasts of prey would turn up. The speed with which Vultures made their appearance from almost every quarter of the Heavens was most amazing, though occasionally I have watched near a carcase right in the open for the best part of an hour without any sign from bird or beast to show that the carcase had been spotted although distant specks high up proclaimed that Vultures were on the lookout. Generally the sky would appear altogether untenanted, but within a very few minutes there would be a heaving, hissing and gibbering mass of ungainly birds, fighting one another for places at the revolting feast. They have, of course, more or less parcelled off the upper air amongst themselves, and one bird seeing its next door neighbour, probably a mile or two away, dropping to a dead or dying animal in its own area, at once sets off on a colossal glide towards the spot for which its neighbour is obviously heading. Number 3, perhaps another mile or more farther on, spots No. 2 commencing its volplane and of course follows suit, and this sort of thing is repeated until all the Vultures for many miles around the spot to which the first bird dropped have been apprised of the fact that a meal awaits those who can arrive in time to participate in the torn flesh of the departed. Numbers I fear arrive only to envy their gorged brethren flopping heavily and sleepily around a useless mass of scattered bones, only of use to the slinking cowardly Hyæna, who will not dare to put in an appearance till dusk in spite of the fact that he possesses colossally formidable jaws, which enable him to crush up the largest bones as if they were those of a chicken, and with which he could lay out almost any adversary with which he is likely to meet.

The rush of air through the semi-closed wings of half a dozen Vultures dropping at a steep angle at terrific speed, makes a noise so loud and so strongly resembling that produced by a gust of wind in the tree-tops, that I have been completely taken in by it on more than one occasion, and not until the falling birds have actually come within my line of vision have I realised my mistake.

Such greed do these brutes possess, that, in spite of an enormous wing-spread, which enables them to drift literally on motionless pinions from one horizon to the other in practically still air, at the end of their disgusting repast they can hardly raise themselves from the ground, and so lethargic do they then become that one can walk right in amongst them and with a reflex, which I unfortunately did not possess in Bharatpur days, their photography becomes almost too easy to be worth while.

Not alone did Vultures appear on these occasions but crows, kites, an occasional Tawny Eagle and almost always a couple or

so of skulking jackals, which, running hither and thither with much snarling and snapping of jaws and with tails for ever between their legs, would endeavour, not invariably with success, to chase off the birds. On one occasion and one I shall never forget, as my rifle was completely out of action owing to the fired case having separated during extraction, a pair of wolves trotted up to within about 40 yards, and then having spotted me seated by a bush about an equal distance on the other side of the kill, calmly sat down on their haunches and derisively watched me feverishly trying to clear the chamber of my weapon. That, incidentally, was the only occasion on which I have ever had a possible chance of getting one of these animals, as in most parts of India they are now somewhat scarce.

The White-backed Vultures used to breed very freely in the larger pipal trees, one tree frequently having two nests in it and occasionally three. I never once saw a nest that was not difficult of access, so never troubled to investigate any of them. The other true Vultures likewise seemed always to choose situations difficult to reach, and only the filthy Scavenger Vulture was unparticular as to where its abode was placed. The roof of a tumbled down building, the mainfork of a tree but a few feet from the ground or yet a ledge on a church or temple tower, anywhere did for it.

In Secunderabad a short time ago I was watching one of these birds building in the top of a very tall toddy-palm, when an intruder of the same species alighted on the ground near by. The owner of the nest at once took objection to this and set off to chase it from its area, but, as I expected, a fight did not ensue. The two birds merely stood motionless and glared at one another, till at last the intruder thinking discretion to be the better part of valour, made off. All wild things appear to me to be imbued with this instinct which tells them never to fight unless forced to. The reason of course is obvious. A bird or animal injured in any way stands far less chance than its whole brethren in the appalling struggle for existence which takes place in nature, and which is apt to result solely in the survival of the physically fit.

Where a little brute force would, I think, have been of great assistance, was in the absurd farce I saw enacted between a couple of pairs of Common and Black-Headed Mynas (*Acridotheres tristis tristis* and *Temenuchus pagodarum*), which both chose for their nests the same recess between a beam and the verandah roof of my Secunderabad bungalow. I first saw the latter pair at the hole towards the end of April. I was away at the beginning of May and returning on the 11th at once noticed a pair of Common Mynas hanging around. During the next few days they did nothing but pull their weaker cousins' building material out of the hole as fast as it was put in, and after about a week of this appeared to have won the unequal contest and to have started building on their own account. The Black-headed Mynas, however, did not admit defeat, but surreptitiously added material to the Common Mynas' new pile during their absence with the result that the collection of rubbish soon reached colossal proportions. This sort of thing continued till June 4th, when the situation again became critical,



THE BLACK-THROATED WEAVER-BIRD
(*Ploceus bengalensis*)



THE INDIAN WHITE-BACKED VULTURE
(*Pseudogyps bengalensis*)



THE WHITE-SPOTTED FANTAIL FLY-CATCHER
(*Rhipidura bectoralis*)



THE SPOTTED BABBLER
(*Pellorneum ruficeps*)

as both birds appeared to wish to make the egg cavity in somewhat different spots. I don't quite remember what happened after this, and the only other note I find I have made is dated June 18th and is to the effect that 'The Common Mynas win'.

I regret I did not do much in the bird-line during my few months in Secunderabad, but amongst other things I took a number of nests of the Dusky Crag-Martin (*Ptynoprogne concolor*) and a couple of the Singing Bushlark (*Mirafra cantillans cantillans*).

A mystery, worthy of solution by the great Sherlock Holmes himself, surrounds the one and only Nukta's nest I have ever found. It was placed in a hollow in the trunk of a fairly large *nim* tree about ten feet from the ground. When the site was originally chosen the tree would be near water, as floods early in July inundated a great part of the State, the main road, within 50 yards of which this tree stood, being under some four feet of water for many days. However when I found this nest on the 25th October, a rather late date by the way, the water had receded and the nearest wet spot was some marshy ground with a few reed-covered stagnant pools roughly quarter of a mile away. On this date the nest contained five eggs and two newly hatched ducklings. And here is where the mystery comes in—one of the chicks was minus its head, which important piece of its anatomy was nowhere to be seen. I am sure it had not been hatched long, and the duck was on the nest and only flew out on my tapping the tree with a stick. I thought this was rather queer and removed the corpse, but next day my mystification was increased still further by finding but three eggs and two dead ducklings, both treated in exactly the same manner. The heads had completely disappeared, apparently having been quite neatly bitten off. The anguished mother was again in evidence and flew two or three times round the tree while I was scratching my head over this extraordinary occurrence. I did not get an opportunity of visiting the nest again for some days, when I found it obviously deserted, smelling foully, and containing two bad eggs and the decaying body of one headless youngster. A civet cat strikes one as being the most likely murderer, but why take away the heads, and would not a civet cat, mongoose or any other animal have laid out all the ducklings and sucked all the eggs straight away, and do mongooses climb trees, by the way? Perhaps the parents had themselves turned cannibal. Surely rats could not have been responsible? One would expect a Nukta to be able to protect its hearth and home against such lesser vermin.

The Indian Spotbill (*Anas pæcilorhyncha pæcilorhyncha*), another resident Indian Duck, I also saw not far from my bungalow in Bharatpur, but I never had the good fortune to find a nest there. In fact, the only eggs of this species I have ever seen were a clutch of eleven, which were collected by an old shikari from a nest in a patch of reeds at the edge of a tank outside Bellary in the Madras Presidency. I was walking round the tank in the hopes of getting a snipe or two and met the old man as he emerged with his spoils. This pair of birds was also breeding somewhat late as the date was the 2nd of November.

The Ashy-Crowned Finch-Lark's nest shown here was actually taken in December, so I hardly know whether one would call it an exceptionally late effort or a very early one, as the breeding season is usually from about February to August or September. Some of the older books I have, call this bird the Black-bellied Finch-Lark, which strikes me as a most appropriate name, as it helps immensely in its identification. The Finch-Larks are close sitters, and I saw an extraordinary case of devotion to duty a couple of months back, in March to be exact. I was putting a number of recruits through a rapid practice on the rifle range, when I noticed a couple of Ashy-Crowned Finch-Larks about twenty yards in front of the firing point. They were gradually working their way closer and closer up in short little runs. The female eventually took up a position on a small stone, perhaps twelve or fifteen yards away, where she remained for the rest of the practice. The male, however, easily distinguished from his consort as she lacks the deep black underparts, continued his advance to a tiny prickly apology for a bush, where he stood for a few moments close to a nest which was plainly discernable now that its position had been given away. Having inspected the contents and after walking round it once or twice, he settled down on the single egg it contained and ceased to take any further interest in the fiendish noise which was going on within some 20 feet of him; even though I could see the muzzle blast perceptibly shaking the little bush at each discharge of the rifles opposite to him. A few days later, when I was again at the same firing point, the egg had been hatched, and this time both birds brought food to the infant on two or three occasions and the female covered it once for about ten to fifteen minutes. This infant, by the way, was clothed in reddy-buff-coloured down which matched to perfection the sandy surroundings. The nest was a very very scant affair of grass, a mere lining to a small cavity scratched out in the bare ground, but more than half the rim was composed of a strip of dirty grey cloth about four inches long. I have also included here a photograph of a nest of the Desert Finch-Lark (*Ammomanes deserti phœnicuroides*) belonging to another genus which inhabits the more barren tracts of the extreme north of India. I took this in Waziristan at Palosina, the scene of some of the hardest fighting during the Mahsud campaign of 1919 and 1920. The chief interest in the photograph, however, lies in the fact that it shows very well the peculiar habit these birds have of incorporating a ring of pebbles in the nest's perimeter, a habit originally acquired, I should think, to prevent the nest becoming silted up by the flying particles of sand blown by the strong winds which so often occur.


Munias and Common Weaver-birds' nests were numerous everywhere, the latter seeming to have a great predilection for babool trees close to wells or overhanging borrow-pits along the two railway lines. Everyone is familiar with the retort-shaped nests of the Baya and the quarrelsome activity of these little finches so soon as the rains start, but perhaps not with those of the Striated Weaver-Bird (*Ploceus manyar peguensis*). The latter's nest

shown here was just within the edge of a patch of elephant-grass and quite alone. I noticed it while photographing a colony of Common Weaver-Birds in a babool tree nearby. This Weaver-Bird always makes its nest in the manner shown, bending over the tops of the reeds to form a sort of scaffolding from which to suspend it.

Though a distinctly catching tuneful whistle from the corner of the compound by the railway embankment, close to the Common Babblers' nesting site, continually proclaimed the presence of a pair of White-browed Fantail Fly-Catchers, the beautiful little cone-shaped structure created by this species was not to be found within my garden's limits, and it was not until the breeding season was nearly over that I discovered two nests in the scrub jungle some 200 yards across the railway. One was a used one at the extremity of a branch of a prickly babool tree at about 5 feet from the ground; the other was in process of construction across a twig springing from the lowest branch of a *nim* tree, and about ten or twelve feet up. Unfortunately it was torn down a few days after I found it.

I had an amusing experience with a very close relative of this species in the Nilgiris, i.e., the White-Spotted Fantail Fly-Catcher (*Rhipidura pectoralis*), which is the common variety of the hills of the southern half of India, and which, unfortunately, though it does possess quite a pleasing whistle, lacks the tunefulness of the song of the White-browed Fantail. This species seems as a rule to confine its choice of building site to positions very low down, at times nests being scarcely 18 inches from the ground, though the usual elevation is generally between 3 and 5 feet. Around Kotagiri I have taken many nests in apple and pear trees and also in creeper, a favourite spot in the latter being on a more or less horizontal strand crossing the hollow space formed by the creeper drooping over the sides of a bush. These birds seem to suffer setbacks in their domestic life almost as much as do the Bulbuls. In Kotagiri last year I had three nests under observation at the same time and was merely waiting for the completion of nest-building in the one case and the laying of the complete clutch of three eggs in that of the other two. The first was amongst the lowest branches of a coffee-shrub, and close to the ground; another in a bunch of creeper and the third at the extremity of a thin twig of an apple tree, both the latter at an elevation of perhaps $2\frac{1}{2}$ or 3 feet. These last came to grief after the laying of the second egg. The eggs of the one in the creeper were in all probability sucked by a lizard or a mouse, as all I could find was a small perfectly dry piece of shell directly beneath an undamaged nest. The apple tree nest probably had its eggs bounced out of it by a thieving crow, which I am sure learnt of the nest by the absurd action of the fussy little Fly-Catchers who swooped and swore at every crow or other intruder which alighted anywhere near. The weight of a crow, attempting to pitch on the frail branch supporting the nest, would have been fatal to its contents, and when I came along after witnessing the mobbing of the said crow, beneath the nest were two broken eggs, their contents still soaking into the

grass. This pair commenced another edifice a couple of days later in an isolated apple tree some 50 yards from the scene of their recent bereavement, but this had its centre all pulled out just when it looked about complete, incidentally only eight days after their first loss. However, it is the nest in the coffee-shrub which chiefly concerns us here. The coffee patch was but a small one, some 50 yards long and varying from only 10 to about 30 yards wide. So, hearing a pair of these highly energetic Fantails making the unmistakeable fuss, consisting of harsh squeaks interposed with occasional little whistles, which augured the presence of a nest, I went down and hunted through the plantation from end to end, the search resulting in the discovery of three, the first of which was under construction, the other two obviously being earlier efforts. The state of this nest gave one a very fair idea of the method of construction. A rupee placed on top would have more than covered it, nevertheless it was a perfect replica of the finished article and its symmetry was faultless. I imagined it was a finished but abnormally small one, but on both of the two occasions I visited it during the next five days it had grown very perceptibly in size but showed no difference in other respects. On the 10th day I took along with me the half-plate camera and the hide and was relieved to find that two eggs had at last been laid. The place was gloomy, as coffee is always grown in the shade of larger trees, and these bushes were very closely planted with the result that their usual umbrella shape rendered the gloom around the nest doubly intense. On this account it was evident that time exposures would be required. I therefore wished to allay the fears of the nest-owners as much as possible, hence I erected the tent and started to camouflage it with great care. My wife was amusing herself covering the top and sides with her usual success, so I crawled round to the front to tie the flap round the lens of the camera and to complete matters generally. I had to look out what I was doing with my nether appendages, as a kneeling position in front of the tent brought my feet very close to the nest indeed. Having fixed the lens to my satisfaction, I glanced over my shoulder to make certain that the camera was pointing in approximately the correct direction. Imagine my surprise to behold the Fly-Catcher reposing quite calmly on its nest, eyeing without the least concern the soles of my shoes at a range of scarcely a foot. I can hardly see myself reclining at ease directly behind the recumbent form of a dinosaur which is the best analogy I can think of. However, without further delay I got inside the tent, and having waited a little to let her settle down, I loosed off five plates one after the other. I may be quite wrong, incidentally, in saying her, as half-way through the proceedings the two of them changed places, showing that both birds take part in incubation. Unfortunately there were factors I had not taken into account. Time exposures appeared to be out of the question. Firstly, because the bird would not condescend to keep its head still; secondly, because the wind was gently swaying the nest almost continuously. However, by a series of short exposures on each plate I succeeded in attaining my object. But, low be it spoken, I had made the one mistake about which I



THE ASHY WREN-WARBLER
(*Prinia socialis socialis*)



THE CEYLON WREN-WARBLER
(*Prinia inornata jerdoni*)



THE ASHY WREN-WARBLER
(*Psaltriparus ashy*)



have already issued a word of warning. I was too close, with the result that the abnormally large and interesting tail of the sitting bird is out of focus. Of course in this case it was a mistake difficult to avoid, as to get away to a decent distance and yet retain a sufficiently clear view of the nest was almost impossible, especially as I did not wish to damage the coffee in any way.

The only Wren-Warbler I identified in Bharatpur was the Indian Wren-Warbler (*Prinia inornata inornata*) and I took a number of their nests at different times, most of which were of the long purse-shaped variety. In fact, when I come to think of it, I only found one of the small domed nests and this was in a sort of prickly weed on 'Chink Hill' and was only about six inches off the ground. 'Chink Hill', incidentally, being a low elongated feature about one and a half miles behind the bungalow, and so christened as it was a most excellent hunting ground for that beautiful little gazelle, the Chinkara. The photograph here of the Ceylon sub-species (*Prinia inornata jerdoni*) was taken on the summit of the Nilgiris where this bird and the Ashy Wren-Warbler are exceedingly numerous; in fact round the southern and eastern edges of the plateau the latter is, I consider, possibly barring the Southern Indian Stonechat (*Saxicola caprata atrata*) and the Southern Red-Whiskered Bulbul (*Otocompsa emeria fuscicaudata*), perhaps the commonest bird. While on the subject I might make mention of the fact that a bird which a very few years ago was unknown on the top of the Nilgiris can now be reckoned among the common birds, of the edge of the plateau at any rate. This is our old friend the Common Indian House Crow (*Corvus splendens splendens*), which in such places as Coonoor and Kotagiri is now almost as numerous as the Jungle Crow. Another bird, which is not usually counted among the Common Nilgiri birds but which I find very numerous in both the above places, especially so all around Kotagiri and overlooking the Mysore Ditch, is the Ceylon Red-Vented Bulbul (*Molpastes hæmorrhous hæmorrhous*). It is however absent, I believe, from the vicinity of Ootacamund and the Kundahs. Whether the House Crow has penetrated to Ooty I do not know, but if not, I am certain it will not be long before it does so. However I was talking primarily of the Wren-Warblers. The last time I was up in Kotagiri was in April and in this month large numbers of both the above mentioned Warblers were breeding. The majority, however, were in the early stages, and even towards the end of the month, in one patch of tall weeds which I searched pretty thoroughly, out of three nests of the Ashy Wren-Warbler and four of the other, only one of each contained eggs. One of the Ceylon Wren-Warblers' nests was in a wild gooseberry bush and was domed, while those in the straight stemmed weeds were all purse shaped. The Ashy Wren-Warblers' nests were of the type which is constructed by cemented together scraps of grass fibre, vegetable down, etc. The other two types of nest which are described in the new *Fauna* (Second Edition) as being built by this species I also took in the Nilgiris. The loosely-woven grass nests often seem to be constructed in thick weed or grass, and, where the nest is built in a large-leaved plant, the Tailor-bird type of nest is made.

In fact I have always been inclined towards the opinion that there are in reality but two types of nest, these two being the roughly woven affair, somewhat resembling that put together by the Common Wren-Warbler, and what might be termed the scrap nest, but, where these nests are constructed in large leafed plants, the leaves are conveniently incorporated in the structure and at times render a great saving in other materials possible. If the nest happens to be built in a spot where there are even small leaves, those leaves contiguous to the nest will always be found sewn to its sides, making a structure which is, practically speaking, midway between the leaf-nest and the other. In other words they largely depend on what they are built in, and probably mainly differ only on this account. It would be interesting to see if each pair sticks exclusively to one type of nest. Personally I do not think this will be found to be the case. I am of opinion that a bird can and will vary its type of nest to accord with the situation chosen, and will not necessarily stick to one kind of site in order that it may always build the same sort of nest. Unfortunately I have never had a chance of elucidating these points, and all the above is, I fear, mere conjecture.

To return to the plate of the Ceylon Wren-Warbler, here this bird evidently disliked the look of the lens, and, though it visited the nest a goodly number of times during the morning, I was only able to get off three exposures on it. It had not really settled down to incubation, the third egg having been deposited only that morning, so it was content with an occasional glimpse of its possessions just to see that all was well, and these glimpses it usually contrived to obtain by working its way up to the nest from directly behind, sneaking through the weeds till it got right up without ever having come out into the full view of the lens. It would then cling to the back of the nest and peer through the thin wall at the eggs within.

Northern Golden-backed Woodpeckers were fairly numerous in Eastern Rajputana and gave one plenty of opportunity for observing their ways, but I find in my all too scanty Bharatpur notes that I made no mention of a habit which I noticed in the Southern race in Madras, namely that they feed almost as much on the ground as they do on trees. Often have I noticed a pair revelling in the treasures of an ant-heap, and occasionally as many as three at once distributed over the surface of the compound, moving about in ungainly hops in company with Mynas or White-headed Babblers (*Turdoides griseus griseus*). Round about Madras they seemed very common indeed, but of course their loud high pitched trilling call at once draws attention to them and probably causes one to imagine them more numerous than in reality they are. In the hills of South India they are replaced by Malherbe's Golden-backed Woodpecker (*Chrysocolaptes guttacristatus delesserti*) which though a bit larger is practically indistinguishable except on close examination, but may easily be recognized in the field by the very feeble imitation of the former's cry. It is also a higher pitched effort and decidedly tinny.

At Mercara in Coorg we were once making our way

along the side of a pretty narrow ravine when the rapid tapping of a Woodpecker, probably this species, was wafted up from below, and on looking down the nullah I spotted a lonesome dead tree the best part of half a mile away, the only likely-looking thing in the neighbourhood. The jungle was very thick and from down below the tree was difficult to locate, but once found there remained not the least doubt that this was whence the noise had emanated, for every single branch was so riddled with the holes of both Woodpeckers and Barbets, that it was literally but a tottering mass of pulp, and if it has not yet collapsed I shall be very surprised. In one large branch, which was almost in two owing to the wood having decayed round some adjacent borings, I counted on one side of it alone fifteen holes and on walking round discovered that about an equal number had been drilled into it from the far side too.

This same little valley was also the hunting ground of many Black-headed Babblers (*Rhopocichla atriceps atriceps*), happy little creatures true to the family in their way of going about in bands continually talking to each other in subdued cheeps. I feel almost inclined to accuse these birds of nesting in colonies. Of course I have only met with them in Mercara, so cannot really judge, but there I don't think I ever came across less than two nests together and on two occasions I struck seven within a few yards of one another. Both of these batches were in deep wide ditches practically concealed by undergrowth and creeper and running through coffee plantations. Once in the ditch one could walk down its centre quite comfortably. The nests were loose spheres of coarse grass leaves or bamboo leaves, with occasionally a little moss or other material intermixed, and faced into the middle of the ditch, being anchored as a rule in hanging strands of creeper at about six feet from the ground. They were in a row at an almost even interval of about six or seven yards. The breeding season appeared to me to be over, though I see that they are said to breed from December to August, and this was but the end of April. However, two only of the nests were still being used as sleeping apartments by their respective families, one of which I found to consist of two parents and two children, the normal thing I believe. These ditches were rather secluded, but a love of seclusion for breeding purposes is not one of this bird's traits though shy itself, as I also found nests overhanging paths and close to roads. In nearly every case the nest was constructed in hanging creeper or brambles.

In this valley too were many nests suspended over the pretty little stream which tumbled down its centre. It was here also that I made the discovery that the extraordinary notes, which someone unsuccessfully attempted to make me believe were those of the Malabar Whistling Thrush (*Myiophoneus horsfieldii*), were in fact produced by the Travancore Spotted Babbler (*Pellorneum ruficeps granti*). I consider the Spotted Babblers very interesting birds, but, being of a retiring disposition, they are somewhat difficult of observation. Thick cover with a sufficient carpet of fallen leaves is most to their liking. Hence forest with considerable

undergrowth is perhaps their favourite haunt; where they can hop, or perhaps I should say waddle, unobserved, turning over the leaves in their search for insects, occasionally darting in prodigious leaps to catch a bug fleetly than usual. I have seen them in parties of three to five, but more often perhaps in pairs and alone, and, when hunting about the ground, they move with a peculiar somewhat dove-like gait with the body and head close to the ground and the tail bent downwards; a position very much resembling the characteristic poise of the Tree-Creepers. This is their general mode of progression, but, when wishing to move quickly, they are capable of covering a considerable distance in a single hop. They do not confine their activities altogether to the immediate neighbourhood of *terra firma*, but are fairly often to be noticed in low trees and hopping about thick bushes, every now and then leaving off chattering with their neighbours to give vent to their most attractive whistle, a clear far-reaching and very pleasantly toned effort. The song of the Western Spotted Babbler (*Pellorneum ruficeps jonesi*) was lately described in this Journal and is apparently somewhat similar to that which I heard in Coorg. The full lay of the Tranvancore Spotted Babbler, however, struck me as consisting of thirteen notes, and I append here my attempt at recording it in a playable form. Unfortunately it is more than a year since I heard it, and though I memorised it pretty thoroughly at the time, by whistling it after every bird I heard singing, it is quite probable that in the course of the year I have introduced slight variations, which in the aggregate may amount to my version being considerably at fault.



The full song is not always indulged in by the way. The songster often starts off quite merrily and then for no apparent reason suddenly breaks off; it may be towards the end of its effort or almost at its very start. Just like someone who has commenced to whistle the latest popular ditty and in the very centre of a bar has suddenly recalled an imminent appointment with a particularly cruel dentist. Last April I came down the Kotagiri Ghaut from the Nilgiris by rickshaw and when passing through the thick jungle near the bottom, that is, when I was certainly not more than a thousand feet above the plain, I suddenly heard the whistle of a Spotted Babbler quite close. Unfortunately I could not see the minstrel and had not time to stay to listen. Anyhow, the song consisted of the same few notes uttered repeatedly, and, as far as I could make out, was absolutely similar to that recorded for *jonesi*. This I presume would almost certainly be *Pellorneum ruficeps ruficeps* (the Common Spotted Babbler), altogether it produced three bursts for my benefit. In Coorg I was not lucky enough to come across any nests, but at Mahableshwar three years ago I remember finding two nests of the latter species. Though shy birds one of them had very unwisely built its nest in a leaf-filled hollow just within

the fringe of trees which encloses one of the tees on the golf links. It soon became known to the caddies, one of whom pointed it out to me. The bird stuck the publicity till the full clutch of three eggs had been laid, but the frequent disturbing of its attempts to incubate soon caused it to desert. The nest was absolutely unconcealed, its back alone resting against the stem of a small and scraggy prickly stalk, which I suppose called itself a bush. Nevertheless the untidy domed nest, surrounded and half-submerged by the dead leaves, of numbers of which it was itself composed, with the dome a fairly bulky but loosely built affair of fresh grasses, was exceedingly difficult to spot, and only the fact of the bird invariably leaving when one was some fifteen or twenty yards off, gave it away; especially as she always flew straight off the nest into the upper foliage of the surrounding trees.

The second nest was also in forest and close to, but not actually in, a patch of thick undergrowth. Like the other one it was amongst a crackling mass of dead leaves, but its back was let into a little depression in a bank at the foot of a tree. On close examination there turned out to be no real foundation to this nest at all. A hollow had apparently been scraped in the leaves and in this a loose hemispherical roof of thin dry seed grass had been constructed, its diameter being about six inches. The entrance was rather large and somewhat untidy, and the three eggs which it contained rested merely on the fallen debris. I put up the hiding tent and settled down within to wait for the birds to appear, but after spending the whole morning in a most cramped position without result, I discovered to my extreme annoyance, on making a closer inspection of the eggs, that they were all bad and the nest obviously a deserted one. The yolks of all three were thickly clotted. In fact it took me some days to thoroughly clean one of them, so it must have been given up some days before. My misspent morning was not altogether wasted, however, as I had spotted through a spy-hole a Northern Indian Stonechat (*Saxicola caprata bicolor*) visiting its mate on a nest in a bank a few yards higher up the slope.

On this occasion I spent the first ten days of May in Mahableshwar, and besides the above my discoveries included many nests of the Southern Red-whiskered Bulbul, a couple of Southern Black Buleuls' (*Microscelis psaroides ganeesa*), a couple of Indian Pipits', a Ceylon Red-vented Bulbul's, and a Bombay Green Barbet's, a White-Spotted Fantail Fly-Catcher's, and an amazing series of three nests of the Malabar Whistling Thrush. The latter were empty but one and all surprisingly fresh looking and were all clustered together round the face of a small boulder in an almost dry stream, the interval between each nest being at most two feet. I cannot now think why on earth I never photographed them, as, though empty, the photograph would have been of considerable interest. They were, of course, in all probability successive nests built by one pair of birds.

Of the Black Buleuls' nests, one was in a decidedly unusual position being barely five feet from the ground in a thickish bush close to a forest ride. The other was in a more usual situation,

resting, as it was, in a fork at the end of a slender horizontal branch of a small forest tree in a clump at the edge of the race course. Even this was considerably lower than the normal as it was certainly not more than 15 feet up. As the tree did not look as if it could bear much weight, I sent up a small and light chokra to see what was in the nest, and before I could stop him, the young imp had stepped on to the very branch holding the nest. Down it bent to a fearful angle, and fearing it was going to break, he hurriedly relieved it of his weight with the natural result that he gave to us below a very creditable demonstration of the working of the old Roman ballista. Up shot two eggs about four feet into the air, one of them to flatten itself against a branch higher up the other spreading its greasy remains over my hand, as I made a valiant but unsuccessful attempt to retrieve it intact.

The owner of the jerry-built Southern Red-Whiskered Bulbul's nest in the accompanying illustration provided me with considerable entertainment. The nest, as one can see, was built with the usual disregard of durability, concealment and want of forethought displayed by many species of Bulbuls. All the supports being on one side only, the increasing weight of the fast growing youngsters was causing it to heal over to an increasingly perilous angle each day. I was not there to witness the rude awakening of the infants to the realization of the lack of common sense of their revered parents, but I hope the structure withstood the forces acting on it until they were old enough to look after themselves. The bravery of the bird I took to be the female was really most amazing. On my discovery of her treasures she retreated but to a twig within a few inches of the nest and danced there in a perfect fury with beak agape and wings vibrating half-open, swearing and spitting like an infuriated lynx, and when I presumed to put my fingers into the nest and to snap off one or two twigs which were in the way, it really looked as if she were about to attack my hand. Seeing her attitude, I at once rigged up the camera, but the moment the fearsome lens was attached her extraordinary courage evaporated, and not until I had brought the hiding tent into use would she venture to the nest again.

In my wanderings in Mahableshwar through a number of patches of rather stunted and fairly open forest growth I came across a great many old nests of what I suppose must have been the White-Throated Ground Thrush (*Geocichla citrina cyanotis*). They were placed on an average about 5 or 6 feet from the ground in the fork of a tree or fairly open bush, and moss and mud figured largely in their construction. I saw a pretty fair number of these birds about and besides possessing an energetic and very pleasant song, which they proclaim most vigorously in the evenings, they have a peculiar note, if note it can be termed, extremely like the noise of a screeching slate-pencil.

However, let us hark back yet once more to Bharatpur, as I cannot finish this chapter without some reference to the bird noises of the night. Around the bungalow these were many and varied, but the most frequently heard and easily distinguished were the chuckling of the nightjars and the grunting hoots of the

Owls. The Common Indian Nightjar (*Caprimulgus asiaticus*) was excessively common and once, when making my way across a little hollow close to the railway embankment, I found myself on the point of stepping right amongst a bunch of five. One does not often spot a motionless Nightjar, when on the ground at any rate, but it was the black shining eyes of one of the five which just caught my attention. The call of this bird has indeed been very rightly likened to the noise which is produced by a stone skimming over ice.

Chief among the owls were the Dusky Horned Owl (*Bubo coramandus*) and secondly his near relative the Indian Great Horned Owl (*Bubo bubo bengalensis*), at least one pair of which lived at Chink Hill. These birds used to stray as far as my compound and consequently I often heard their grunting calls. The distinctive voice of the Dusky Horned Owl was to be heard from all sides, but chiefly from the grounds of the Moti Jhil, a bungalow just across the railway with a superabundance of large pipal and other leafy trees around it. This bird is very common in Bharatpur and every single grove and garden appeared to have its tenants, as well as the large patches of jungle such as the Chota Ghana and those around the Keoladeo Ghana. During the rains and the cold weather this owl is most noisy and not only at night. In the monsoon months, especially, their voices are often to be heard in the very middle of the day. True, dusk is the time when they are most vociferous and are more generally met with, but in habits they are very diurnal. Nor do they shun mankind, but on the other hand, like the noisy little Spotted Owlets, are commonly to be discovered in groves and gardens in and around villages. The most curiously placed nest of this species, which I have ever come across, was placed in a dead and leafless pipal on a low horizontal branch which barely cleared the roof of a kutch building in the very centre of one of the largest villages in the State. The nest was large, probably the appropriated nest of a Kite, and was a conspicuous object from all over the village, as was also our friend the owl sitting on it. A number of naked children playing around the base of the tree in no way disturbed her Majesty, nor did she appear in the least put out by the unwonted sight of shining topis and white faces. Another rather curious situation was in the main fork of a very lonesome tree standing in some flooded fields, also close to a village. It was certainly not more than 7 feet from the ground.

This fine bird breeds from the end of the rains onwards to about December, and, besides appropriating the disused nests of other birds, is said to build for itself a fairly large stick affair. The nests, which I suspect to be of the bird's own manufacture, are rather shallow but nevertheless pretty compact and about eighteen inches or so across and are generally placed fairly high up amongst the lesser and leafy branches of large trees, and consequently are fairly well concealed. The appropriated nest on the other hand may apparently be anywhere, so long as it takes the fancy of the bird and appears to be a large and strong enough structure for its nursery.

In 1918 I noticed a White-backed Vulture on its nest half way up, and close to the trunk of, a perfectly tremendous but climbable pipal tree. Wondering if by any chance they ever make use of a nest twice over, I revisited the spot again in December 1919. Over the rim of the nest there peered at me, not the ugly head of a Vulture, but the catlike visage of a Dusky-horned Owl. This is the nest which appears in the accompanying plate, and over the getting of it I thought my last moments had come. Unfortunately I am small of stature but I possessed an orderly of considerable length, and with his aid I managed to triumph over the long stretches between the branches and eventually reached my destination, let down a cord and drew up the camera and its appendages and obtained a number of photographs. But as is so often the case, the descent was very much more difficult than the ascent had been. When I had been unable to reach a branch the orderly had gone up first and then pulled me up, and we ought to have adopted a like method of going down, instead of which he went right down first and I followed. Having lowered myself over a particularly thick branch, at what felt to me some 40 feet from *terra firma*, I suddenly found that my feet were dangling in mid-air, and that I was unable to reach the branch below. That from which I had come was so fat that my hands began to slip, nor was the main trunk any use to me as it was about 4 feet thick. Well to cut matters short I continued to slip inch by inch expecting every moment to feel my toes in contact with the longed-for foothold, but I had no luck and perforce had to drop. No, I am sorry to have to disappoint you and end up so tamely, but my flattened remains were not peeled from off mother earth and carried home on a stretcher. Instead I hit the branch plumb in its centre and did the only decent balancing feat I've ever done in my life. Nevertheless, as below me there was an unbroken drop to the ground, I must admit that on the way home my nerves felt somewhat shattered.

In a previous article I have made mention of the conclusions I have been forced to draw from numerous observations of the attitude of birds towards suspicious sounds where the evidence of sight is lacking, and lately I had occasion to test my theories on the subject somewhat severely. Firstly in the case of a Fly-Catcher which is common to the Nilgiris, and inhabits practically every sholah in the place, no matter how small it may be. This is the Nilgiri Blue Fly-Catcher (*Stoparola albicaudata*). The second case was that of a Ceylon Hoopoe (*Upupa epops ceylonensis*), whose mate I had observed incubating a clutch of five eggs in the base of a stone wall. Both these birds, one must admit, are well used to human beings, the latter especially so, and are likely in consequence to recognize the human voice.

At Kotagiri the Longwood Sholah especially swarms with the former, and immediately on entering the wood their sweet and penetrating song attracts one's attention. Normally they appear to hold cavities in the sides of banks in great esteem as positions for nesting sites. Thus the banks along the inner side of the paths intersecting the sholah are most favourable spots to search and

yielded three nests containing eggs, all in convenient positions for photography. April, by the way, is an excellent month in which to look for fresh eggs. The first of these nests had close to it some small roots projecting from the bank-side, and I counted on the bird alighting on one of these before hopping into the nest, which was almost completely out of sight in a pretty deep and hence dark cavity well protected by a number of thin hanging roots. To photograph successfully the bird on the nest would have been impossible without completely removing and altering the whole surroundings, which would also probably have caused desertion. I therefore put up the tent on the opposite side of the path, and focussed the camera on the exposed roots. Fortunately my wife had come out with me and after shutting me up had retired round an adjacent corner to write letters. Within five minutes I heard the Flycatchers singing lustily within a few feet of the rear of the tent and next moment a whirl of wings over the top proclaimed the fact that one of them was coming to the nest, and to my extreme annoyance, come to the nest it did, my roots being given a complete go-by, as she—it turned out to be the female—flew without stopping outside straight into the cavity and settled down on the eggs. The male after singing for a little longer close by, flew off, evidently quite satisfied that all was in order. Now what was to be done? If I shooed her off it might make her too frightened to come back again, and even if she did reappear she would in all probability go straight in again. I therefore started talking; first in a subdued voice, and then loudly, until I felt that a sudden shout would not startle her too much. She sat on quite contentedly under this treatment and did not even depart when I yelled as hard as I could to my wife to come to the rescue. I had with me a pretty small watch of the folding travelling variety about $2\frac{1}{2}$ inches square and this my wife laid over the egg cavity and again took her departure. Within a very few minutes the female was back, flying straight in as before. For a moment she stood still on the edge of the nest, then she began to peck at the watch all round, and finally started trying to scratch it off and to force her way underneath it. With such frenzy did she persevere that I became alarmed for the safety of the eggs. After a bit, however, she quietened down and I expected her to fly out and at last make use of the roots as a resting place while she thought matters out. But not a bit of it; instead, she sat down on the top of the watch and remained there until my patience gave out. A fact worthy of record, by the way, and one which provides much food for thought, is that the whole time she was frenziedly scratching away she was singing as hard as she could. Again I had to appeal to my wife and this time the whole cavity was stuffed with bracken, but this too turned out no good, as the mother hovered about outside it like a humming bird, eventually worming her way into a bit of a hole in the bracken and sitting only half in sight of the lens. A third time I was obliged to call to my long-suffering memsaheb, and success was at last achieved by pushing in a little stick just by the side of the bracken, and on this the female had the decency to alight repeatedly with the result that

a number of plates were soon exposed. After the fifth exposure, however, I inadvertantly bumped the camera while she was on the perch, causing the lens to shift a trifle. Immediately away she flew and it was some twenty minutes before she dared return. Noise she could put up with, but the moment something moved as well, suspicion of the hide invaded her. Unfortunately Longwood Sholah is a pretty gloomy place at any time, and the negatives obtained on this occasion were all under-exposed. A few days later I tried my luck on one of the other nests and, profiting by the experience gained in the above episode, I blocked up the entrance to the nest before ensconcing myself within the hide. To provide a perch was unnecessary, I am glad to say, as I do not like introducing anything extraneous into the picture, but always endeavour to depict both birds and their nests as they are naturally. This time I had no difficulty except with the light, as, although I gave as long exposures as the bird would permit, these negatives too show distinct signs of underexposure, mainly traceable to the fact that my ultra-rapid plates had come to an end and I had had to revert to ones of slower speed. I also tried to photograph a Rufous-bellied Short-wing (*Brachypteryx major major*) in this same wood, but once more the gloom proved too much for me. Hence illustrations of the nest and eggs of this bird and of those of the Black and Orange Fly-Catcher (*Ochromela nigrorufa*) and the Indian White-Eye (*Zosterops palpebrosa palpebrosa*), two more denizens of these Nilgiri Sholahs, are all I can run to.

The Hoopoe episode was as follows:—The strong-smelling scattered collection of rubbish which they presumed to call a nest was, as I have said, in a stone wall some fifty yards above a road. After I had used up four plates on the male bird who was engaged in bringing food to its mate, whom I discovered was not only sitting on eggs but also on two newly hatched babies, my wife called up to me that she was going home as she was unable to discover a shady spot in which to sit. Although at that moment the female was within and the male actually on the ground but a few feet from the tent, I shouted down to my wife telling her what my intentions were so soon as I had finished with the Hoopoes, and the illustration here was actually taken in the midst of the conversation, as our friend took not the slightest notice of my voice but paddled up as usual to the nest-hole, inviting his spouse the while to come and take the fat grub he held for her in the tip of his bill. The male Hoopoe is a good husband, and certainly works hard for his living, as his lady seldom leaves the nest once she has started laying, until the young have been hatched some time. As this one came flying to the nest he always uttered a series of not unsnipe-like 'penches', and when at the hole invariably talked to his wife in a subdued rapidly-repeated sort of coo.

The Hoopoe is, of course, far from being a man-fearing creature, but even so I am perfectly certain that no Hoopoe would enter its nest with a man sitting but 5 or 6 feet away, without, at any rate, an enormous amount of tuition first. I am all the more convinced, therefore, that as a general rule birds cannot and will

not believe that real danger is to be apprehended when they can only hear, but cannot see, anything of which to be afraid. They seem, in fact, to lack all powers of deduction. A few of the smaller birds, and with a decrease in size a corresponding increase in foolhardiness appears to be the rule, can be got with but little expenditure of patience used to the idea of coming to the nest with one close by. In my school-days I remember being busy photographing the abode of a Spotted Fly-Catcher. The pair concerned had nested in the ivy on the wall near my study window for some years and were quite at home with the genus school-boy. While I was focussing with my head under the cloth, one of the birds landed on the top of the camera and flew thence straight on to the nest, where it sat quite unperturbed during the remaining operations.

The photograph of the Southern Grey Tit (*Parus major mahrat-tarum*) here depicted was taken about 30 yards from a garage in front of which a car was being washed, but here too I made use of the hiding tent though without troubling greatly about its concealment, as even a cheeky little Tit is apt to differentiate between 5 or 6 and 100 feet, though, why—goodness only knows, as it is just as easy for it to be spotted entering its nest at the longer distance as it is at the shorter.

Laughing Thrushes are as a rule put down as shy customers. Yet one of the only two cases I know of, where a bird has really not seemed to mind a tinker's curse about my presence was in that of a Nilgiri Laughing Thrush (*Trochalopteron cachinans cachinans*). This bird was only building, yet it came to the nest while I was erecting the tent late in the evening with the intention of photographing it the following morning, my idea being that it would be advisable to give the birds a chance of getting used to the hide during the night as otherwise they might fear the structure and refuse to come near for hours of valuable daylight, as birds often do not appear greatly attached to their homes until after the eggs have been deposited. As, however, it was then too late for photography, I went on with the erection of the hide, while the bird remained on its half-finished collection of moss, although I was working in full view and within 8 or 9 feet of it most of the time.

The next morning there were no signs of either of the birds until I had been in the tent for upwards of half an hour. Then, however, one of them arrived, incidentally without any fresh material, sat in the still untidy nest and commenced to shape the egg cavity with its breast, by turning round and round as it sat. It was in the midst of this process when I took the photograph opposite. After a time it suddenly gave forth a few piercing yells in answer to its mate, and then settled down for a bit of a snooze. Perhaps to talk of this bird's call as a 'Crow' rather than a 'Yell' would be the more accurate, as, in order to produce the noise it raised its head high in the air, puffed out its chest and opened wide its bill, assuming thereby a very cock-like stance. Unfortunately I was unable to photograph it in this attitude, as it was in such a position that one of the supporting branches of the nest

was directly between it and the lens. After I had taken a number of photos, I pulled aside the front flap of the tent, and lo and behold the bird still sat on, and it was only when I emerged completely that it took it into its head to quit. It is, of course, altogether fallacious to conclude that because one bird is fearless, any other of the same species will be equally so. This is emphatically not the case, and I had sat in front of another Nilgiri Laughing Thrush's nest but a few days previously for a couple of hours before the female, who had been sitting for some days on two eggs, picked up enough courage to return, and I am convinced that she would not have faced the camera at all, had it not been for the advent of a heavy rain-storm. She obviously objected to her treasures getting wet. The resultant photos of this bird are in consequence somewhat peculiar, as the light is strongly reflected from the many rain-drops which cover her back and tail.

The hiding tent, to which I have at times been alluding, is not the one which I mentioned as ordered from Cawnpore, as the makers thereof evidently considered that they knew more about my requirements than I did myself, and sent a thing altogether unsuited to the work in hand. I therefore drew up the plans for another in Secunderabad, whither I was transferred early in 1922, and having enlisted the services of a local *durzi* and later of a carpenter, we evolved between us the contraption which has since proved so useful and quite up to everything so far required of it. It is made of good quality khaki drill, double along all seams and edges, and consists of a centre piece 3 ft. by $2\frac{1}{2}$ ft., which forms the roof, on to which are stitched end pieces $3\frac{1}{2}$ ft. deep which widen to 4 ft. along their bottom edge, and two side pieces 3 ft. long at the top to fit the roof and $3\frac{1}{2}$ ft. deep to correspond with the ends. These widen to 5 ft. along the bottom edge. When spread out flat the tent therefore looks something like a Maltese cross. The reason why I made the walls in the above manner is that it allows one a certain amount of latitude in the ultimate shape of the erected tent. Along each of the sides of these four flaps or walls are four eyelets so that all or any pair of the edges may be laced up as required. Horizontally on the outer surface of each flap are stretched three khaki ropes caught at intervals to the drill so that loops are formed, and across the top ropes are likewise sewn. Into these loops, grass, branches, ferns, etc., can very quickly be rammed; thus the tent takes but a short time to camouflage. The frame to hold this up is of wood, and consists of but four separate pieces as follows; two plain end-pieces, in length the width of the roof, and two side-pieces corresponding in length to the roof, and on which the four legs are hinged in such a manner that when folded each pair lies close together and along its respective side-piece. Unfolded they are kept in position by hooks, the sort of thing one uses to keep windows and shutters open in this country, and also by four 5-inch pins which are inserted in turn through eyelets in the four corners of the drill roof, then through corresponding holes in the wooden side-pieces and end-pieces of the frame and finally down the centre of the legs themselves. These pins therefore not only help to fix the legs, but

also clamp the separate pieces of the frame together, and keep the roof tight and in position.

It will be seen, therefore, that the whole affair can be erected in a few moments. This frame is made of teak but any other strong wood will of course do, as teak is certainly rather heavy. Still the complete tent only weighs 15 lbs., and rolls up into a cylindrical bundle some 3 ft. long and about 7 in. thick. It could probably be made to weigh less as both the side and end-pieces are considerably stouter than is absolutely necessary, being $1\frac{3}{4}$ in. wide and 1 in. thick. I prefer to have it really strong, however, as one does not want to risk breakage at a critical moment, and often, owing to the slope or unevenness of the ground, considerable strain is imposed upon it. For spy-holes one can either squint through the corners or else cut slits where required in the drill. After a few tears have been made, one will find that the practice of apparently ruining the tent may be stopped, as only a limited number are required in any one wall. For the lens, slits are likewise cut, just big enough to stretch over the lens seating, over which is pushed a rubber ring to prevent the flap slipping back and covering the lens up again.

When at work in the tent, it will be seen that one is seated in an oblong box 3 ft. high, 3 ft. long and $2\frac{1}{2}$ ft. wide. This may not appear altogether palatial, but it is ample for me with a half-plate camera, a 5 by 4 reflex and their necessary impedimenta and since starting to use it, I have never had occasion to endure the agonies one reads of through sitting in a cramped position for hours on end, for the simple reason that I have found movement quite easy in this amount of space. The middle of the last sentence is rather suggestive of an advertisement for a quack medicine. However, to continue; the only improvements I can think of at present are that the legs might be lengthened by about 6 inches. However the actual details of size rest with the person who is to use the tent, and he will of course adapt it to his own bulk. Another possible improvement is that it might be made sunproof. Now, it certainly is not, and one has often to endure a topi in consequence, and as topis are bulky this is a nuisance, since one is liable with such an ungainly thing on one's head to bump it into the sides or roof of the tent, and so upset the equilibrium of the bird one is attempting to photograph.

As I have described the hide, it would not be out of place to describe the cameras I use in it. Undoubtedly where one's means limit one to a single camera, a half-plate field camera is indispensable. For all round general work both in and out of the hide this type and size is without doubt the best, and is what I think most bird-photographers use. My half-plate is an Ensign Triple Victo with a battery of Zeiss and Ross lenses of different focal lengths, which I have picked up at odd times and which give me a choice of varying the size of the picture without necessarily having to find a new position for the camera, an undertaking which is often impossible or nearly so in a restricted place or in trees. The lenses in question are a 4 inch Ross wide-angle, a Ross-Zeiss convertible anastigmat with focal lengths of $7\frac{1}{4}$ inches for the com-

bination and $11\frac{1}{2}$ inches and 14 inches for the separate components, and also a Zeiss Protar giving focal lengths of 17 and 29 centimetres. I have in addition a Zeiss telephoto attachment for use with the Protar, but so far it has proved itself quite an unnecessary possession. Still, if it comes in useful even once, I shall consider it has justified its existence. Actually only one good anastigmatic lens, preferably of the convertible type, is necessary; though to possess two is more convenient, namely one of long focus for use in the hide, as this enables one to get a decent sized image without having to work uncomfortably close up; and the other of fairly short focus say 6 inches or 7 inches at the outside, for photographing nests and eggs, etc. The long focus lens should work at a very large aperture, so that short exposures may, where necessary, be given in a bad light. Only lately I expended 14 plates on a bird standing near its nest in a mud bank in a pretty thick forest, yet with a lens working at f 5.4 and using exposures of one-third second, one and all were underexposed, even though the sun was shining brightly directly overhead, and within a few feet of the nest was a large glaring patch of sunlight. A couple of attempts at short time exposures of about one second were even on the underexposed side, and these latter negatives were of no use as the image of the bird was in both cases blurred through movement. The ordinary photographer, used to taking landscapes, etc., will be surprised at the comparatively long exposures required to avoid underexposure in the photography of birds and their nests. Hence the necessity for the lens with a very wide stop.

With a bird seated on its nest the stop and exposure problem is not so acute, as many sitting birds remain pretty still, and I have at times used exposures of the best part of a minute. A Nilgiri Quaker Babbler (*Alcippe poioicephala poioicephala*) obligingly sat still for such a period, when I was once obliged to stop down the lens, a Ross telecentric, as far as it would go to get the necessary depth of focus, as the only possible position for camera and tent was so close that the limited extension of the reflex I was using was insufficient to bring even the furthest portion of the bird into sharp focus. However, by slightly unscrewing the front combination of the lens and stopping down as I have said, I succeeded in getting a quite passable negative. The nest was in a particularly dense patch of forest, and as it turned out, double this exposure would have produced a better result. Very often the click of the shutter has the most convenient effect of making the bird sit as still as a rock, while it listens with strained attention for further repetitions of the unknown sound. I have found this to be particularly the case with the Thrushes. Everyone knows the look of annoyance and the outraged demeanour assumed by the common Song Thrush at home when one presumes to approach it when seated on its nest. However, I have found that even with a nervous bird, which is continuously shifting its head or which jumps up seemingly the moment one presses the release, exposures of up to about half a second don't so very often show movement, the bird apparently not being able to get on the move, after being startled by the opening of the shutter, until after the exposure is

finished. Of course one needs to judge the correct moment to take the photograph so as to get the bird during a period of stillness, and this is largely a matter of practice.

Another necessary adjunct in the lens is a surplus of covering power, and sharpness to the very edges of the surface it does cover, that is to say that in a half-plate camera, a lens which covers a whole plate without showing any loss of definition round the edges should at least be used. Especially does this necessity arise in taking photographs to show off the contents of nests, as with the ordinary tripod the camera cannot usually be depressed as a whole at a sufficiently deep angle, and the lens has therefore to be still further tilted, with the result that its axis no longer passes through the centre of the plate. To tilt the nest instead of the camera is very bad policy indeed and only results in an unnatural-looking photograph. Besides, one wants one's reproductions to show the subject exactly as it was constructed and used by the owners. Most field cameras have a tilting back but unfortunately not all of them tilt backwards sufficiently far, that is, well beyond an angle of 90° with the camera base. However, this defect can very easily be remedied as a rule by removing the locking devices and rescrawing them on to the camera body a little lower down. To be able to swing the back of the camera in this manner comes in especially in photographing nests on open patches of ground, such as beds of shingle or on the sea-shore, open commons, in heather, etc., in fact anywhere where it is an advantage to get the foreground and distance both in focus in order to give one a fuller impression of the bird's habitat without having to stop down greatly and so unnecessarily increase the exposure. The near foreground of the picture to be in focus requires a longer extension than does the distance, and as the image cast on the plate by the lens is, of course, inverted, this state of affairs is consequently arrived at when the back is tilted beyond the vertical, as the top of the plate is then further from the lens than is the bottom of it.

Lately I have been using a 5 by 4 reflex in the hide with most excellent results, but, before I go any further, let me say that though a great convenience a reflex is not an absolute necessity, whereas a field camera, and preferably a half-plate one, most certainly is. The reflex does, however, save one from making the mistake of photographing the bird when it is out of focus, and also of course one has a chance of putting a bird rapidly in focus and getting its photo, where the time wasted and the noise made with an ordinary camera in removing the plate, refocussing and replacing the plate, renders such a course out of the question, and all one can do is to wait for it to condescend to alight on the exact spot on which one has previously focussed. The lens I am using in this reflex is an 11 inch Ross Telecentric working at *f.* 5.4 and the main reason why I invested in the 5 by 4 size is that it naturally racks out further than does a quarter plate of the same make.

The obvious course when buying a reflex is to choose one with a long extension, but I bought mine in India when reflexes were both hard to get and infernally expensive, so took the first I


could get. There are now many good cameras of this type on the market, and some of them are probably more suitable for bird-work than mine, which incidentally is an Auto-Graflex, with the performances of which, however, I am perfectly satisfied, and I see from the current Kodak list that there are many improvements in their present Graflex cameras, which put mine clean out of date. The low-powered telephoto lenses seem also to have been much improved of late.

When using the reflex in the hide, I always remove the focussing hood for convenience, as the restricted height of the tent makes it very awkward to work with it on, and the tent itself is quite dark enough to render the image on the screen easily visible. It is quite safe to do this, and even when using the mirror as the shutter with the focal plane shutter open, I have never had a plate spoiled by extraneous light entering through the screen and passing round the edges of the mirror. Using the mirror in this way gives one a pretty silent and comparatively long instantaneous exposure, probably of about half or one-third of a second. If the shutter is put out of action and at open, and one regulates the raising and lowering of the mirror with the right hand, while holding down the release with the left, one obtains a means of giving an absolutely silent time-exposure of any duration and this is of considerable use at times when working on a sitting bird, which shows signs of sleepiness or is naturally still.

When taking small birds, I invariably use quarter plates in wooden carriers in the plate holders of both these cameras, since, as the image is small and generally only a comparatively small area is required from which the finished enlargement will be made, it is only a needless expense to use the larger plate. I have already mentioned why it is better to get a small image all of which will be in focus, so it is unnecessary to go further into that point.

As regards the best plates for use in bird-photography I will not say much, as most photographers have their own particular fads about them, but I must say the ordinary special rapid type by any of the well known makers take a lot of beating in this country. Of course orthochromatic plates give a better impression of the true grading of the colours of the subject. Recently I have taken to the Ilford Iso-Zenith plate wherever I have anticipated the necessity of high speed, and have found it most excellent. Its H & D speed number is 700, so one can naturally cut down exposure considerably. The Hoopoe photo was taken at $1/40$ second at *f.* 8 on one of these plates, when the sun was not shining directly either on the bird or its immediate surroundings. The Southern Grey Tit is also an effort on one of these plates and was taken at $1/20$ at *f.* 8 on quite a dull day. The only rub is that they do not keep for very long, and once the tin in which they are sealed has been opened, it is advisable to use them up pretty quickly.

An enlarger, by the way, is a most useful possession, and one which I consider the bird-photographer cannot well do without. I for one, know that many of my earlier photographs would be infinitely better enlarged than they are at normal size, but one is very chary of sending hard earned negatives through the post,



THE CEYLON HOOPOE
(*Upupa epops ceylonensis*)



THE SOUTHERN GREY-TIT
(*Parus major mahrattarum*)



THE BLACK AND ORANGE FLY-CATCHER
(*Ochromela nigrorufa*)

especially the Indian one, let alone the feeling one has that other people do not always treat one's negatives with the care with which one would like them to be treated. It is somewhat annoying, to say the least of it, for a negative to come back with a large scratch right across it, far more annoying in fact than if one had smashed it altogether oneself. To do one's own enlarging will eventually turn out an economy, too, and the initial expense need not be great. The half-plate camera, with the lens with which the negatives were originally taken, again comes in, and can be used with daylight as the illuminant by having a hole cut in the dark-room window or door with a reflector behind it. Over the hole the camera is closely fitted so that no light can enter the room other than that which passes through the negative and lens, and the bromide paper on which the enlargement is to be made is pinned on to an easel placed in front of the lens on a table or other convenient support.

(The End)

ON A COLLECTION OF MOTHS OF THE FAMILY *GEOMETRIDÆ*
FROM UPPER BURMA MADE BY CAPTAIN A. E. SWANN

BY

LOUIS B. PROUT, F.E.S.

PART II

(With a plate.)

(Continued from page 146 of this Volume)

* 71. *Perizoma schistacea* (Moore)

Anticlea schistacea Moore, Lep. Coll. Atk., p. 273 (1888) (Darjiling).

Htawgaw, April-May, 1923, 1 ♂; Hpimaw Fort, June 1923, 3 ♂♂, 1 ♀.

A small and rather aberrant form, perhaps worthy of a name. Typical *schistacea* belongs almost exclusively to Sikkim and Assam, though it has also been recorded from Dalhousie.

* 72. *Perizoma cerva* (Hmps.)

Cidaria cerva Hmps., Journ. Bombay Nat. His. Soc., xiv, 515 (1902) (Sikkim).

Htawgaw, September-October 1923, 1 ♀.

Only two specimens hitherto known to me, the second being a ♂ ex Coll. Elwes, presumably also from Sikkim, labelled by him 'unnamed in Coll. Atkinson'.

* 73. *Perizoma lacernigera* (Butl.)

Idaea lacernigera Butl., Ill. Het., vii, 109, Pl. cxxxvi, Fig. 15 (1889) (Dharmasala).

Htawgaw, April-May 1923, 1 ♂; 1 ♀; Hpimaw Fort, June 1923, 1 ♂.

Also known from Sikkim.

* 74. *Perizoma hockingii* (Butl.)

Eupithecia hockingii Butl., Ill. Het., vii, 115, Pl. cxxxvii, Fig. 12 (1889) (Dharmasala).

Htawgaw, April-May 1923, 1 ♀, September-October 1923, 1 ♀.

Both this and *lacernigera* may prove racially differentiable from the North-west Indian forms, but Sikkim examples of both seem fairly typical.

* 75. *Perizoma puerilis* sp. n.

Perizoma puerilis Warr. MS. in Coll. Tring Mus.

♂, 20 mm. Face whitish (partly abraded). Palpus longish, 2nd joint with rather long loose scaling directed forward, 3rd joint moderate; pale, with some blackish admixture above. Vertex white. Antennal ciliation minute. Body slender, pale, above mixed with fuscous.

Forewing with apex acute, termen oblique, slightly sinuous; areoles large, R¹ from middle of the outer, SC⁵ from its apex; whitish, suffused with cream-buff; markings fuscous; basal patch small, obsolete posteriorly; cell-mark black, slightly elongate; median band narrow (circ. 1.5 mm.), not intense, darkest at edges, slightly constricted about cell-mark (which is on its proximal edge) slightly outbent in middle; subterminal consisting of rather large white spots, that in cellule 4 elongate proximally (tapering); some weak dark irroration bordering the subterminal, developing a fairly strong proximal spot close to costa and a larger, somewhat elongate one in cellule 5, just crossing into cellule 4; a slight oblique apical dash.—*Hindwing* rather narrow, termen slightly prominent at R¹ and very slightly at M¹, faintly sinuate inward between R¹ and R²; DC strongly oblique, rather weakly biangulate; white.



Geometridæ from Upper Burma.

GEOMETRIDÆ FROM UPPER BURMA.

EXPLANATION OF PLATE II.

Fig.	1.	<i>Krananda orthotmeta</i> , Prout sp. n.	
„	2.	<i>Heterolocha segregis</i>	„
„	3.	<i>Cleora coniozona</i>	„
„	4.	<i>Boarmia virguncula</i>	„
„	5.	<i>Loxotephria perileuca</i>	„
„	6.	<i>Ectropis pulvicopia</i>	„
„	7.	<i>Cleora cryptogonia</i>	„
„	8.	<i>Ectropis embolochroma</i>	„
„	9.	<i>Micrabraxas anisonoma</i>	„
„	10.	<i>Enantiodes consanguinea</i>	„
„	11.	<i>Medasina lasiochora</i>	„
„	12.	<i>Ctenognophos methoria</i>	„
„	13.	<i>Hirasa plagiochorda</i>	„
„	14.	<i>Orsonoba æthocrypta</i>	„
„	15.	<i>Semiothisa ageta</i>	„
„	16.	<i>Ocælophora agana</i>	„
„	17.	<i>Ectropis hiulca</i>	„
„	18.	„ <i>æthregenes</i>	„
„	19.	<i>Sysstema aulotis</i>	„
„	20.	<i>Ectropis zotica</i>	„
„	21.	„ <i>lophomeris</i>	„
„	22.	„ <i>chrysoteucta</i>	„

Forewing beneath more coarsely irrorated, except distally; the pattern of the upper side faintly showing through. Hindwing rather less white than above; a sinuous postmedian line, scarcely beyond middle, strongest posteriorly; traces of white subterminal spots, with some fuscous proximal irroration.

Htawgaw, June 1923, 1 ♂ (type); 'West China', 1 ♀ in Coll. Tring Mus.

Curiously like some South American *Psaliodes*, e.g. *olivaria* Warr. Probably nearest to *ochreotincta* Willem. (*The Entom.*, xlviii, 60) which is rather smaller, less *Psaliodes*-like, with SC⁵ of forewing arising proximally to apex of areole, DC of hindwing rather more angled, etc.

* 76. *Perizoma decorata quadrinotata* Warr.

Perizoma quadrinotata Warr., Nov. Zool., iii, 122 (1896) (Khasis).

Laukhaung, April-May 1923, 1 ♂; Htawgaw, April-May 1923, 2 ♂♂, 1 ♀, June 1923, 1 ♂; Hpimaw Fort, June 1923, 5 ♂♂.

Probably Hampson (*Journ.*, Bombay Nat. His. Soc., xii, 77)—who, however, commonly ignored geographical variation—was justified in regarding *quadrinotata* as a synonym of *decorata* Moore from Sikkim; but as it has been provided with a name and is generally smaller and more sharply marked than the Sikkim specimens I have conserved it provisionally. The Laukhaung and Htawgaw examples are nearly like the Khasi, the Hpimaw (though also small) rather variable and on the whole paler-banded.

* 77. *Perizoma conjuncta* Warr.

Perizoma conjuncta Warr., Proc. Zool. Soc. Lond., p. 381 (1893) (Karen Hills).

Htawgaw, July 1923, 1 ♂.

The specimen is small and pale, but a little worn and may be either *ab.* or a local modification, rather reminiscent of *albofasciata* Moore. The species has been taken in numbers in the Khasis but is probably very local. As the abdomen is crested throughout, the subfamily Keys of Meyrick and Turner would throw it out of *Perizoma*.

* 78. *Perizoma latifasciata* (Warr.)

Epirrhoë (?) *latifasciata* Warr., Proc. Zool. Soc. Lond., p. 375 (1893) (Sikkim).

Hpimaw Fort, 9th to 13th August, 1923, 2 ♀♀, 14th to 18th August 1923, 1 ♂, 1 ♀.

* 79. *Perizoma maculata* (Moore)

Cidaria maculata Moore, Lep. Coll. Atk., p. 277 (1888) (Darjiling).

Htawgaw, April-May 1923, 1 ♀ (large).

Known from Sikkim, Bhutan and Assam.

* 80. *Perizoma albofasciata* (Moore)

Cidaria albofasciata Moore, Lep. Coll. Atk., p. 277 (1888) (Darjiling).

Htawgaw, April-May 1923, 1 ♂; Hpimaw Fort, June 1923, 3 ♂♂.

Perhaps a new race, but only the Htawgaw specimen is in good condition. All are small, especially two of the Hpimaw.

* 81. *Perizoma lucifrons* sp. n.

♂, 25 mm. Like *variabilis* Warr., except as follows:—

Head and palpus pale (or whitish) ochreous-brown. Abdomen dorsally with a transverse white line at base (often reduced to a spot but always conspicuous), followed on tergite 2 by a bright, somewhat tawny line or narrow band; anal tuft less mixed with ochreous-brown. Forewing relatively slightly broader; scarcely so strongly glossy. Hindwing slightly less glistening, in some lights rather less white (more tinged with grey); apex faintly darker-clouded. Hindwing beneath a little more powdery (white, with fine but copious dark irroration).

Htawgaw, March 1923, 1 ♂, 4th to 10th April 1923, type and another ♂; Sikkim, August 1909 (F. Möller) 8 small ♂♂ in Coll. Brit. Mus., 3 ♀♀ undated (O. Möller and J. F. Pilcher) in Coll. Tring Mus.

There has been a good deal of confusion over the species named *variabilis* by Warren and it is perhaps less variable than was supposed. True *variabilis* (as Elwes notes, *loc. cit.*) is a high altitude form (10,000–13,000 feet) and it is conceivable that *lucifrons* may come to be regarded as a race from more moderate elevations, though I consider it unlikely. Warren himself mixed with it other near allies and indeed his allotype ♀ as labelled by himself (but not a legitimate one, as the recorded series was '4 ♂♂, 4 ♀♀ from Tonglo') is Möller's ♀ *lucifrons* registered above. Others (Hampson, *Faun. Ind. Moths.* iii. 373, and, I regret to add, myself in preparing vol. iv. of 'Seitz') allowed the 'lumping' to stand and judged the species from material in the British Museum instead of from Warren's originals. My differentiation of *mediangularis* Prout (*loc. cit.* 259 = *mediangularia* in err. Pl. 12 c) is in part vitiated by this confusion, but it is at the least another very distinct race, larger and relatively longer winged. In any case the V-shaped midsubterminal mark cannot be used differentially in any of the three; it is developed in Warren's four Tonglo ♂♂ and a ♀ from Jongri but not in Tonglo ♀♀, developed in *mediangularis* allotype ♀, but weak in the type ♂, obsolete in type *lucifrons*, but strongly developed in Capt. Swann's other examples, its posterior arm thick and tinged with buff. As to the further errors regarding ab. *albimacula* Prout, see *infra*.

* 82. *Perizoma variabilis* Warr. (?)

Perizoma variabilis Warr., Proc. Zool. Soc. Lond. p. 377 (1893) (Sikkim).

Hpimaw Fort, June 1923, 1 ♂.

This specimen is not quite fresh, but appears to be evidently a further development of the form described by me (*Seitz Macrolep.* iv. 259) as *variabilis* ab. *albimacula*, by Warren as 'a third varietal form' (and chosen for figuring, Pl. xxx. Fig. 18, wrongly cited on pp. 377 and 434 and by Hampson as 'Fig. 17'), by Hampson as the principal form, relegating the type, together with whatever else he lumped therewith, to a brief supplementary paragraph. The Hpimaw example has the large white subterminal spot continued posteriorly, though more narrowly, almost to the hind margin. I am still inclined to allow *albimacula* to rank as an aberration of *variabilis*; and I do not think the present specimen can be a parallel variation of *lucifrons*.

* 83. *Perizoma seriata* (Moore)

Cidaria seriata Moore, Lep. Coll. Atk. p. 278 (1888) (Darjiling).

Hpimaw Fort, June 1923, 2 ♂♂.

This species is known from North India and Formosa.

* 84. *Perizoma bipartaria* (Leech)

Cidaria bipartaria Leech, Ann. Mag. Nat. Hist. (6) xix. 650 (1897) (Pu-tsu-fang, W. China).

Fenshuiling Pass (4 miles from), early July, 1923, 1 ♀.

On the hitherto unique type (also a ♀) see further *Seitz Macrolep.* iv. 259, Pl. 7 k.

* 85. *Trichoplites latifasciaria* (Leech)

Cidaria latifasciaria Leech, Ann. Mag. Nat. Hist. (6) xix. 644 (1897) (Wa-Shan, W. China).

Hpimaw Fort, June 1923, 2 ♂♂.

Both are darker, especially in median area, than Leech's unique type, which is a female. The discovery of the male proves that it belongs to Swinhoe's genus *Trichoplites*, though the discocellulars of the hind wing are not, as in the type species, biangulate.

* 86. *Trichoplites cuprearia* (Moore)

Anticlea cuprearia Moore, Proc. Zool. Soc. Lond. p. 656 (1867) (nom. praecoc. ?) (N. India).

Hpimaw Fort, 24 September, 1922, 1 ♂.

Worn, but according to the structure, etc., belonging here as a darker form, apparently not unlike an also worn specimen from the Nagas in the Tring Museum. Westward the species reaches Sikkim, eastward, I believe, West China, but it seems nowhere common.

* 87. *Lampropteryx opistholasia* sp. n. (Pl. 1, Fig. 15)

♂, 39 mm. Face broad, slightly prominent. Palpus little over 1; somewhat blackened. Antennal ciliation short (less than 1). Abdomen rather elongate, the lateral pencils not developed. Head and body light fuscous.

Forewing with cell less than $\frac{1}{2}$; glossy dark fuscous, the distal area paler; band between basal patch and median band as dark as they, separated from the former by a curved whitish line, from the latter by two slender, approximated whitish lines, the outer the sharper, bluntly angled at both folds; cell-spot elongate, black; median band traversed, as in the allies, by faint lunulate lines, its breadth at costa 7 mm., at hind-margin 4 mm., the fine white line which bounds it gently excurved between costa and R^2 , bluntly bilobed between R^2 and M^2 , acutely indented at M^2 ; sub-terminal formed of small detached lunules (posteriorly spots), the accompanying dark proximal marks anteriorly cuneiform; an oblique white line from apex.—*Hind wing* relatively rather small, abdominal area beneath folded, containing a pencil of brown hair, a clothing of lighter hair also occupying a great part of this wing beneath, strongest about M and proximal half of M^1 and M^2 ; glossy pale greyish fuscous, almost uniform.

Forewing beneath with weak cell-dot and traces of postmedian line and sub-terminal dots, both obsolete towards hind margin. Hind wing with elongate blackish cell-mark, indistinct dentate postmedian line (incurved between the radials) and traces of whitish subterminal dots.

Hpimaw Fort, early July 1923, the type only.

Very near *rotundaria* Leech (*Ann. Mag. Nat. Hist.* (6) xix. 659), of which it might even be the hitherto unknown ♂ but that the discocellulars of the hindwing are not biangulate. Forewing less hoary between postmedian and subterminal, apical dash stronger, antemedian less angled behind middle, postmedian more angled at M^2 , cell-mark large. Unique in the genus in the transference of the sexual hair to the hindwing.

88. *Electrophaës aliena* (Butl.)

Cidaria aliena Butl., *Ann. Mag. Nat. Hist.* (5) vi. 230 (1880) (Bhutan).

Htawgaw, June 1923, 1 ♂ (ab.), September–October 1923, 1 ♂.

The June specimen is a pretty aberration, rather small and bright, the hindwing above less white than in the great majority of specimens of this species, beneath with the postmedian line more sinuate. One Darjiling ♂ in coll. Tring Mus. resembles it in size and coloration. The other Htawgaw specimen is typical. This form inhabits North India and I believe *tsermosaria* Oberth. from West China and an unnamed *Electrophaës* from North Tibet to be races of it.

* 89. *Ecliptopera lativittaria* (Moore)

Scotosia lativittaria Moore, *Proc. Zool. Soc. Lond.* p. 657 (1867) (Sikkim).

Kangfang, June 1923, 1 ♀.

This fine species is scarcely an *Ecliptopera*, but I do not at present see where better to refer it.

90. *Ecliptopera silaceata* (Schiff.) (?)

Geometra silaceata [Schiff.] Schmett. *Wien.*, p. 113 (1775) (Austria).

Htawgaw, early July 1923, 1 ♀.

Most of the Asiatic forms referred by Hampson (*Faun. Ind. Moths*, iii. 357) to this widely distributed species have proved distinct, though closely allied; but there remains a residuum which would seem to be really races of it and to these belongs perhaps the present specimen, which has both wings darkened, the central band of the forewing rather solid, the markings of the distal area somewhat reduced and the postmedian line of the hindwing rather distally placed. More likely, however, it is a distinct, but at present unnamed species, as has proved to be the case with a somewhat similar ♂ from Omei-Shan of which I have had the genitalia examined by Mr. Burrows.

* 91. *Ecliptopera relata* (Butl.)

Cidaria relata Butl., *Ann. Mag. Nat. Hist.* (5) vi. 229 (1880) (N. E. Himalayas).

Htawgaw, June 1923, 1 ♂.

92. *Eustroma inextricata* (Walk.)

Cidaria inextricata Walk., List Lep. Ins. xxxv. 1691 (1866) (Northern India).
Hpimaw Fort, August 1923, 2 ♀♀.

This species ranges from Sikkim to West China, presumably in suitable places in South China and certainly on Formosa and is represented in Japan by a race or very close ally, *aerosa* Butl.

93. *Photoscotosia miniosata* (Walk.)

Scotosia miniosata Walk, List Lep. Ins. xxv. 1354 (1862) (Sylhet).

Hpimaw Fort, 24th September, 1922, 1 ♂, June 1923, 3 ♂♂; Htawgaw, August 1923, 2 ♀♀.

Range: N. India to W. China, Formosa. An unnamed race on Luzon.

* 94. *Chartographa fabiolaria trigoniplaga* (Hmps.)

Callabraxas trigoniplaga Hmps., Tr. Ent. Soc. Lond., p. 312 (1895) (Nepal).

Hparè, end of August 1923, 1 ♂.

From the slender material accessible to me, I judge that *fabiolaria* Oberth. (China) and *trigoniplaga* Hmps. (North West and Northern India and Tibet) are races, both variable, of a single species; the present specimen is an aberration—or perhaps slight local modification—of the latter.

95. *Dysstroma cinereata* (Moore)

Cidaria cinereata Moore, Proc. Zool. Soc. Lond., p. 662 (1867) (Sikkim).

Htawgaw, 4–10 April 1923, 1 ♂, April–May 1923, 1 ♂, early July 1923, 1 ♂, August 1923, 1 ♂; Hpimaw Fort, June 1923, 2 ♂♂.

Distributed from Sikkim to West China.

* 96. *Dysstroma subapicaria* (Moore)

Cidaria subapicaria Moore, Proc. Zool. Soc. Lond., p. 663 (1867) (Sikkim).

Htawgaw, June 1923, 1 ♂; Hpimaw Fort, June 1923, 1 ♂, 1 ♀.

Perhaps a race, the hindwing rather pale, inclined to develop a darker border. Typical *subapicaria* is known from Sikkim and Bhutan.

* 97. *Dysstroma citrata dentifera* (Warr.)

Polyphasia dentifera Warr., Nov. Zool. iii. 387 (1896) (Darjiling).

Lankhaung, 7th March, 1923, 1 ♂; Hpimaw Fort, June 1923, 1 ♂.

This race is distributed in North India and Tibet. The species has an enormous range in the Palaearctic Region and even in North America.

* 98. *Thera cyphoschema* sp. n. (Pl. 1, Fig. 6)

♂ ♀, 24–27 mm. Head reddish brown. Palpus about 2; pale at base, darkened on outer side. Antenna in ♂ with rudimentary pectinations (scarcely 1), surmounted by short ciliation. Thorax reddish brown, paler beneath. Abdomen rather pale, especially beneath. Legs partly blackened, with white irroration and extremities of joints.

Forewing reddish brown, about as in average British *obeliscata* Hb. (less glossy than in *cognata* Thnb.); basal patch subobsolete, indicated by two faint lines (the proximal sometimes hardly traceable, the distal weakly outbent on both folds) and in the most sharply marked specimens by some slight band-like shading between them; cell-mark elongate, oblique, black; antemedian equally oblique, generally absorbing the cell-mark, sometimes just proximal, blackish anteriorly, right-angled at M, very faint posteriorly; postmedian from costa 3 or 4 mm. beyond antemedian, gently incurved at first but mainly oblique outward to a projection at R², thence very feeble, lunulate-dentate; veins in median area blackened, generally strongly; some further dark shading usually present, commonly becoming blackish in the radial lobe; subterminal line indicated, at least in anterior half, where it is accompanied proximally by some dark shading, an elongate mark in cellule 6 commonly conspicuous; the characteristic apical dash of *Thera* rarely noticeable.—*Hindwing* whitish grey, with very faint traces of strongly excurved postmedian line,

Both wings beneath glossy greyish, the forewing a little suffused; both with weak cell-dot and traces of curved postmedian line; forewing with pale costal spot just outside postmedian.

Htawgaw, March 1923, 2 ♂♂, 1 ♀ (1 ♂ dated 20. iii), April-May 1923, 7 ♂♂, 5 ♀♀, June 1923, 1 ♂, early July 1923, 3 ♂♂; Blackrock, early July 1923, 1 ♂.

The June-July specimens are on an average smaller, probably a second generation.

The genus *Thera* is Holarctic, straggling into the Himalayas.

* 99. *Triphosa consona* sp. n. (Pl. 1, Fig. 11)

♀, 49 mm. Coloration and general distribution of the markings almost exactly as in well coloured examples of *dubitata* Linn., though with slightly more dark admixture, at least in the paler parts. Palpus blackish, with third joint rather longer than in *dubitata*. Head and body a good deal clouded with black-grey.

Forewing with the central band rather less noticeably divided into two than in most *dubitata*, the vinous shades well distributed in it; antemedian lines more deeply excurved anteriorly than in *dubitata* and in addition with an acute outward projection at fold; the lines between this and basal patch likewise exangled at fold; postmedian rather straight from costa to SC⁵, the double projection at R¹ rather weak and even; the band beyond distinctly tinged with olive; subterminal expanded into a larger white subterminal spot than in *dubitata*; the vinous band proximal to it rather strongly bisected.—*Hindwing* with termen appreciably less deeply dentate than in *dubitata*; the lines and vein-dashes stronger; some vinous suffusion in abdominal region.

Underside rather darker than in *dubitata*, otherwise similar.

Htawgaw, April-May 1923, the type only.

From the Indian species of the *rubrodota* group also the angular antemedian and the vinous præsubterminal band will readily distinguish it.

100. *Anaëtis affinis* (Warr.)

Docirava affinis Warr., Nov. Zool. i. 398 (1894) (Central China).

Htawgaw, early July 1923, 1 ♀.

Previously known from Sikkim and West and Central China. The specimens quoted by Warren from 'Japan' were certainly—like a good number of Chinese specimens in the early days of the Tring Museum—erroneously labelled.

* 101. *Chætolopha incurvata* (Moore)

Eupithecia incurvata Moore, Lep. Coll. Atk., p. 268 (1888) (Khasis).

Htawgaw, August-September 1923, 1 ♂, 1 ♀.

The taxonomic position of this very distinct species is problematical, but in any case it is certainly no *Eupithecia*. Its nearest relative is probably *rubicunda* Swinh. (*Tr. Ent. Soc. Lond.*, 1902, p. 647), from Perak, Sumatra and Borneo, which can also be provisionally included in *Chætolopha*.

* 102. *Hyphenorhynchus erectilineata* (Moore)

Lygranoa erectilineata Moore, Lep. Coll. Atk., p. 272 (1888) (Khasis).

Htawgaw, July 1923, 1 ♂, 3 ♀♀, August 1923, 1 ♀; Laukhaung, July 1923, 1 ♀.

103. *Collix ghosha* Walk. (form ?)

Collix ghosha Walk., List Lep. Ins., xxiv. 1249 (1862) (Ceylon).

Langyang, 14th September 1922, 1 ♂.

A worn specimen, but almost certainly referable to a *Collix* which at present stands as a possible form of *ghosha*, considerably larger and with the maculation of the underside slightly less heavy, but will probably demand separating as a species. *C. ghosha* is spread over the greater part of the Indo-Australian Region, the large form (or ally) perhaps restricted to Assam, Burma, the Malay Peninsula and Formosa (vide *Ent. Mitt.* iii. 248, No. 77).

104. *Collix* sp.

Htawgaw, September–October 1923, 1 ♀.

Likewise worn, but apparently nearest to *hypospilata* Guen, though larger and with the subterminal spots beneath less sharp. Probably a species unknown to me, scarcely possibly a remarkable ab. (still larger) of the preceding.

* 105. *Horisme flavofasciata* (Moore)

Collix flavofasciata Moore, Dep. Coll. Atk., p. 270 (1888) (Darjiling).

Htawgaw, 4–10 April 1923, 1 ♂, 14 October, 1923, 1 ♀.

A very fine pair of this apparently rare species, hitherto known only from Sikkim and Assam. Both are large, perhaps racially.

* 106. *Horisme hyperythra* (Hmps.)

Phibalapteryx hyperythra Hmps., Faun. Ind. Moths iii. 347 (1895) (Nilgiris).

Htawgaw, July, 1 ♀.

A fairly common species in India and Ceylon and reappearing in Japan, the Liu Kiu Islands and Formosa.

Parazoma gen. n.

Face without cone. Palpus rather long, second joint with long projecting hair-scales beneath, third joint well developed, somewhat drooping. Antenna in ♂ minutely ciliated. Metathorax with metallic-mixed crest. Pectus scarcely hairy. Femora glabrous. Hind tibia with all spurs. Abdominal crests scarcely developed; terminal segments in ♂ well developed, valve broad, uncus pointed, labides long, their lower arms perhaps embracing the juxta, which appears to be tripartite, apparently no coremata. *Forewing* with much the shape, scaling and pattern of a *Perizoma*; areole very large, undivided, its anterior margin approaching C, SC⁵ well stalked with SC²⁴, R¹ from areole. *Hindwing* moderately ample, not continuing pattern of forewing; termen very little waved; DC oblique and (in the type species) generally angled (partly through the incurving of DC²), C anastomosing to near end of cell, SC² stalked, R² about central or very slightly behind.

Type of the genus: *Parazoma ferax* sp. n.

Probably related to *Eupithecia* but quite different in habitus, in details of venation, in lacking the 'body-plate' of the 8th sternite of the ♂ abdomen, etc. From *Perizoma* it differs in the single areole, labides with heads not united, etc.

* 107. *Parazoma ferax* sp. n.

♂ ♀, 19–24 mm. Head and body fuscous, more or less sprinkled with black and white. Foreleg above blackish, marked with white at ends of joints.

Forewing rather glossy, coloured about as in dark examples of *Perizoma bifaciata* Haw. or as *P. variabilis* Warr., etc.; a bright orange-brown subcostal line (conspicuous with the lens) almost from base to postmedian; basal patch brown, irrorated with black, limited by a very slightly curved pale line; intermediate band brown; median band black, about 3 mm. in width (rather variable), not broadening anteriorly, its proximal edge more or less curved, its distal with small subcostal and (double) central projections; cell-spot deeper black, but quite inconspicuous; fine pale lines bounding this band, the distal one white, at costa double; distal area brown, clearest just beyond median band; then more or less irrorated and clouded with black (variable); a pale mid-subterminal spot, as in many *Perizoma* and equally variable both in size and colour—pure white or light brown; terminal line interrupted by the veins, which are here tinged with brown; fringe mottled.—*Hindwing* glossy dark grey, almost without markings; a small cell-dot sometimes distinct.

Forewing beneath much more blurred, the median band more dissolved into lines. *Hindwing* beneath variable, but always with alternating pale and black-grey lines, the postmedian generally the strongest.

Hpimaw Fort, August 1923, 5 ♂♂, 5 ♀♀ (including the type ♂); Htawgaw, August–September 1923, 2 ♂♂; Hparè, September 1923, 7 ♀♀.

Also known to me from Darjiling, July–August 1886 (H. J. Elwes), 3 ♂♂, 1 ♀ (determined by Hampson—presumably without examining the forewing

venation—as ‘*Larentia? bicolor* Warr. var.’), August 1904, 1 ♀ in coll. L.B. Prout; ‘Sikkim,’ 7,000 feet, 5 ♂♂, August 1895 and September 1909, and 1 ♀ in Coll. Brit. Mus.; and? Khasis, August 1894, 1 ♂ in Coll. Tring Mus. with DC of hindwing not biangulate.

* 108. *Parazoma semifusca* (Warr.)

Perizoma semifusca Warr., Nov. Zool. iii. 123 (1896) (Khasis).

Htawgaw, March 1923, 3 ♂♂.

All three are aberrant in the obsolescence of the dark border of hindwing. This may prove racial, but I have seen at least one Khasi ♂ of nearly the same form.

Except for the non-biangulate discocellulars of the hindwing and some very secondary differences in the genitalia, this species fits well into *Parazoma*. It is not a *Eupithecia*, as Hampson (*Journ., Bombay Nat. Hist. Soc.*, xii. 80) made it, lacking the body-plate of the 8th sternite, as well as differing in shape and facies.

109. *Physetobasis dentifascia* Hmps.

Physetobasis dentifascia Hmps. Faun. Ind. Moths., iii. 386 (1895) (Dharmasala).

Htawgaw, July 1923, 1 ♂.

Small and dark, perhaps a subspecies. The West Chinese race (?), *mandarinaria* Leech, is, however, larger and more rufescent or (a Yunnan ♂ in coll. Joicey) smallish and rufescent. Slightly different forms, again, occur in Lower Burma and Central China. A Simla specimen in coll. Brit. Mus. agrees with the type. More material is badly needed for working out the variation.

* 110. *Eupithecia russeola* sp. n.

♀, 21 mm. Frontal cone sharp. Palpus moderate, second joint heavily scaled above. Antennal ciliation very short (well under $\frac{1}{2}$). Head and body brownish fuscous, paler beneath.

Forewing with costa scarcely arched, apex rather sharp, termen moderately long, rather strongly oblique, straightish from apex about to R^2 , thence very gently curved to SM^2 , straighter again to tornus; areole undivided; hazel to russet, more reddish tinged than *rajata* Guen., darker than *albispumata* Warr. and without the white admixture; costal margin more fuscous, cut by still darker marks, denoting the commencement of the lines; cell-dot scarcely raised, rather small though slightly elongate, black, with a few whitish scales at its edges; lines in proximal area weak, in central area scarcely noticeable, all nearly as oblique as the postmedian; postmedian rather thick and fairly conspicuous from its anterior angle almost to hindmargin, straightish, parallel with termen, its anterior angle acute, but scarcely noticeable, the costal (inwardly oblique) arm being obsolescent, while the costal spot of the subterminal looks almost like a resumption of the postmedian; the usual pale band outside the postmedian scarcely at all developed, its bisecting line fine and not strong; subterminal line slight, even near the tornus only developing a small and fine white spot; terminal line weak, interrupted; fringe paler, especially in distal half, the proximal half feebly chequered.—*Hindwing* with costa slightly longer and straighter than is *rajata*, apex moderately rounded, termen straightish between R^1 and M^1 ; pale grey to whitish grey basally and costally, nearly concolorous with forewing distally and posteriorly, the abdominal margin also mostly darkened with broad fuscous shading; cell-dot minute; postmedian markings (the line itself, narrow pale band beyond, with very fine limiting line or vein-teeth) indicated except at costa; terminal line rather more conspicuous than on forewing; fringe as on forewing.

Forewing beneath rather paler, less reddish, much as in *rajata*; cell-mark strong; costal margin slightly darkened; four costal spots present, the antemedian the weakest; postmedian line and some faint lines in outer area indicated. Hindwing a little paler; cell-dot stronger than above; the curved postmedian line complete, fairly distinct; lines of outer area well indicated.

Hparè, September 1923, 1 ♂ (type); Kangfang, September–October 1923, 1 ♂.

The Kangfang specimen is slightly paler and (especially on hindwing) more distinctly marked than the type. Another Hparè ♂, of the same date

as the type, may be a greyer aberration, but seems to have the antennal ciliation slightly less short, the postmedian line of forewing slightly more proximal, with the pale band beyond it more noticeable and associated with a pale oblique streak from apex. More probably a separate species, but not very fresh.

111. *Eupithecia* sp. n.

Htawgaw, September–October 1923, 1 ♀ (torn).

Larger and paler than the preceding, rather narrower, cell-spot of forewing rounder, central area narrower, its boundary lines more regularly developed. An extremely worn ♀ (Htawgaw, September 1922) may possibly connect it with the greyer relative of *russeola* (supra).

112. *Eupithecia* sp. n.

Hparè, September 1923, 1 ♂.

Smaller and rather more glossy than *russeola*, in shape and markings nearer to No. 110.

113. *Eupithecia robiginascens* Prout

Eupithecia robiginascens Prout Nov. Zool. xxxiii. 9 (1926) (Bhutan).

Hpimaw Fort, 14–18 August, 1923, 1 ♀.

Not very fresh, but apparently referable to this species, which inhabits N. India.

* 114. *Eupithecia tricrossa* sp. n.

♂ ♀, 17–19 mm. Palpus moderate, second joint triangularly scaled above, third joint short, pointed. Antenna in ♂ minutely ciliated. Head grey. Body above glossy blackish, with a slight admixture of metallic bluish, beneath very pale grey. Foreleg above darkened, spotted with white at ends of joints.

Forewing with apex not acute, termen gently curved, not exceptionally long; areole undivided; glossy white-grey, the markings dark grey tinged with brown (or, under a strong lens, grey-brown with minute longitudinal blackish irroration); some metallic bluish scales at base; a large black cell-spot of raised scales; basal patch and antemedian band sometimes fused together, the latter touching the cell-spot, about as oblique as termen; a postmedian band about 1.5 or 2 mm. beyond cell-spot, angled outward just before R^1 , then almost parallel with termen, very slightly sinuous; fine parallel lines proximally and distally hereto; subterminal line of the ground-colour, fine, slightly crenulate anteriorly, interrupted between the medians; three dark patches proximally to it, costal, radial and tornal, the middle one tapering anteriorly to accommodate an oblique pale streak directed towards apex, the posterior one about reaching M^2 ; the narrow dark distal shade only interrupted at the veins; fringe weakly clouded.—*Hindwing* with apex rounded, termen rounded-prominent at SC^2 and M^1 , straightish or almost concave between; light grey, slightly darkened at abdominal margin and base of M ; a blackish cell-dot; a dark grey, slightly curved postmedian line (band); a slight dark shade at tornus; terminal line and fringe as on forewing.

Underside less glossy; white-grey, the forewing proximally and anteriorly more suffused, darkest on proximal part of costal margin; cell dots and postmedians distinct; proximal subterminal shade of forewing indicated, strongest anteriorly; hindwing sometimes with anterior indications of a similar subterminal shade.

Hpimaw Fort. 9–13 August, 1923, type ♂ and 1 ♀, August 1923, allotype ♀.

Also occurs in Sikkim (Darjiling, Nagrispur and Kurseong) and Bhutan (Buxa), but—like very many Indian *Eupithecia*—seems never to have been named. Hampson confused it with *atrisignis* Butl. from North-West India, which is browner, less strongly glossy, less variegated, but with the veins in middle of forewing reddened, the cell-spot generally less large, less oblique, the median area broad, antemedian line strongly curved and less oblique than termen, the oblique band-like shading obsolete, the postmedian more angled, the subterminal shade not distinctly tripartite.

* 115. *Eupithecia acyrtoterna* sp. n.

♂, 23 mm. Face without distinct cone below; dark grey. Palpus $1\frac{1}{2}$, second joint heavily scaled above and especially beneath, third joint distinct; pale, with blackish irroration. Vertex pale grey. Antenna evenly ciliated, the ciliation fully 1. Body grey, apparently black-mixed on side (discoloured), paler beneath. Foreleg blackened, the joints with pale tips.

Forewing moderately elongate, costa straightish except at base and near apex, termen long, very oblique, very gently curved; areole undivided; pale grey; slightly (less than in *virgaureata* H. Dbld.) tinged with brown; dark irroration; cell-dot black, fairly large (about as in *virgaureata*); costa with indications of dark antemedian, median and postmedian spots and of dark dashes between; lines indistinct, excepting the double postmedian (which is angled near costa, much as in *virgaureata* or scarcely so strongly) and the pale subterminal (with some incomplete dark shading proximally); subterminal only very slightly expanded near tornus, here rather markedly dark-bordered proximally; terminal line black, interrupted at the veins; fringe with large but weak spots opposite the veins.—*Hindwing* rather narrow, the termen from SC^2-M^2 almost straight (only very faintly oblique outward to R^1 , incurved between the radials and excurved between R^3 and M^2); rather paler than the forewing, except in abdominal region and at extreme termen; markings obsolescent anteriorly, nowhere strong; cell-dot small; subterminal spot near tornus rather large, but very little paler than its surroundings; terminal line and fringe as on forewing.

Both wings beneath paler; forewing with fairly large cell-dot, weak costal spots, faint postmedian and rather faint proximal-subterminal band; hindwing with smaller cell-dot, shadowy sub-basal and median bands (the latter just outside the cell-dot, interrupted in middle) and fainter subterminal; both with the terminal line indistinct.

Hpimaw Fort, 14-18 August 1923, the type only.

Perhaps near *virgaureata*. The shape of the wings, though foreshadowed in several Indian species, is characteristic and I think only rivalled by *quadripunctata* Warr. (1888, N. W. India), which is a large, browner species, with rather more elongate forewing, larger cell-spot, the other markings weaker, etc.

* 116. *Eupithecia anasticta* sp. n.

♀, 23 mm. Face and palpus dark fuscous; frontal tuft rather thin but sharp; palpus with second joint heavily scaled above and beneath, third joint rather strong. Occiput and collar paler. Antenna minutely ciliated. Thorax and abdomen fuscous, the thorax above rather darker; abdomen robust, with the first tergite paler. Foreleg blackened, with pale rings at ends of joints.

Forewing moderately broad, apex not acute, termen gently curved, oblique (normal), a little straighter between M^2 and SM^2 ; areole elongate, double (*Eucymatoge* in sensu Mey.); glossy dark fuscous (not quite as dark as *lucigera* Butl., Ill. Het. vii. 115, Pl. cxxxvii, Fig. 11, more like average *vulgata atropicta* Dietze); the dark irroration very fine and even, not conspicuous; cell-mark rather elongate, not intense; lines almost entirely obsolete; postmedian just discernible, a little less angled anteriorly than in *rajata* Guen., accompanied distally at costa by a small whitish spot and at the radials by almost invisibly minute dots; subterminal faintly indicated, developing two very conspicuous white spots—a small one in cellule 3 and a large one behind M^2 .—*Hindwing* concolorous in posterior half, paler in anterior; a small cell-mark, followed about middle of wing by a rather broad but very weak postmedian line; whitish and dark marks near abdominal margin indicating beginnings of other (obsolete) lines; two white subterminal spots as on forewing, but rather more equal in size; fringe very weakly spotted.

Both wings beneath very slightly paler than hindwing above, the posterior part of forewing a little paler still; marked with cell-spot, rather broad but rather weak postmedian line, fine and weak subterminal, with a faint band-like dark shade proximally, and very faint indications of a bisected pale band between the latter and the postmedian.

Hpimaw Fort, 9-13 August 1923, the type only.

I have pointed out on other occasions (*The Entom.* xl. 169; *Seitz Macrolep.* iv. 274) that the splitting up of this genus on the single character of divided or

undivided areole gives unnatural results. The character in question is a crux of geometrid classification and must be used with great caution. After many years' work at the family I can only agree with Pearsall (*Canad. Ent.* xxxvi. 210) that as 'Nature follows no hard and fast lines' we must submit to the anomaly of using a character as generic in one case while rejecting it in another. Every intermediate stage is found in *Nyctobia* (proximal areole minute or dividing wall very weak, etc.), insistence on the character divides species which in all other respects appear to be closely and truly related (e.g. *Eupithecia pini* Retz. and *bilunulata* Zett., etc.), the Palaearctic group associated by Meyrick (*Tr. Ent. Soc. Lond.* 1892, p. 68) as *Eucymatoge* has no character of homogeneity as a whole (as compared with *Eupithecia*) and precisely the same variation of structure occurs in other allied genera (*Lobophora* and *Mysticoptera*, *Dasyuris* and *Dasyternica*, *Cartaletis* and *Aletis*, *Cyllopoda* and *Flavinia*, etc.). Of course much the same may be said of other artificially accepted characters—structure of ♂ antenna, number of hindtibial spurs, etc.—but as Meyrick (*Tr. N. Z. Inst.* xlix. 249-50) has laid impossible weight on the above arguments in connection with a generally more fundamental one (cf. Forbes, *Journ.*, *N. Y. Ent. Soc.*, xxv. 45) it seems necessary to refer to them here. For an able discussion of the subject see Turner, (*Tr. R. Soc. S. Austral.*, xlvi. 226-8). As regards the Indian species of *Eupithecia*, Hampson was evidently so satisfied as to their homogeneity that he neglected to examine the areole, for it is divided in three species which he refers here—*eupitheciata* Walk., *rigida* Swinh. and *acutangula* Hmps. (*Faun. Ind. Moths* iii. 398-400.)

* 117. *Eupithecia albispumata* (Warr.)

Eupithecia albispumata Warr., *Proc. Zool. Soc. Lond.*, p. 384, Pl. xxxi, fig. 23 (1893) (Khasis).

Htawgaw, September-October 1923, 1 ♂, 1 ♀.

Both are rubbed, but they seem to agree with Warren's species.

* 118. *Eupithecia leucenthesis* sp. n.

♂ ♀. 18-20 mm. Face fuscous. Palpus shortish (about $1\frac{1}{4}$), rather browner, Vertex whitish. Antennal ciliation in ♂ rather short (about $\frac{1}{2}$), in ♀ rudimentary. Body very pale brown, the wing-tegulae almost white.

Forewing rather narrow, apex rather sharp, termen long, slightly bowed, strongly oblique; cinnamon to orange-cinnamon, the costal margin feebly spotted with fuscous; an ill-defined whitish longitudinal streak, filling the cell and somewhat diffused behind it (notably in proximal area), distally mingling with the ground-colour but traceable at the base of the radials and obliquely forward to apex; cell-dot minute, blackish; lines very feeble, traceable in fuscous shading especially at M, where they tend to form vein-spots or dashes; postmedian sharply angled at R¹, thence parallel with termen, accompanied distally by indications of the usual pale band with fine thread-line; sub-terminal faint, accompanied proximally by a band of a very slightly darker shade than the ground-colour; terminal spots not very strong; fringe weakly mottled.—*Hindwing* rather narrower, costa rather long, apex rounded, termen straightish to R³ or M¹ at M¹ very bluntly bent; whitish proximally and anteriorly, then nearly concolorous with forewing; traces of waved fuscous lines; abdominal margin darker spotted.

Underside with small cell-dots and (at least on hindwing, which is a little paler than forewing) traces of waved lines.

Hparé, September 1923, 1 ♂, 2 ♀♀.

None of the specimens is perfectly fresh, but they are well recognizable; *asema* Hmps. (*Ill. Het.*, viii, 107, Pl. CLII, fig. 23), known from the Nilgiris and Palnis only, is very similar, but more whitish, the forewing rather acuter, etc., and has the antennal ciliation fully 1.

* 119. *Eupithecia leucostaxis* sp. n.

♀, 22-23 mm. Head white, tinged with ochreous; face nearly smooth, terminating in a minute ochreous cone below. Palpus rather more ochreous; moderate, second joint rough-scaled above and beneath, third joint small, not distinct. Thorax above whitish, mottled with ochreous; abdomen belted with ochreous; both whiter beneath. Fore and midlegs more or less darkened, except at the extremities of the joints.

Wings fairly broad, in shape perhaps comparable with *subnotata* Haw., the termen of *forewing* scarcely even so oblique as in that species; white, irrorated and copiously marked with ochreous, much as in *ochracea* Warr. (*Proc. Zool. Soc. Lond.*, 1888, p. 321)¹ but rather brighter; cell-dot minute or obsolescent; a number of fine waved lines, somewhat as in *subnotata*; boundary lines of median area slightly stronger and with slightly stronger white lines outside them; antemedian and the correlated lines bisinuate; postmedian right-angled at R¹, thence slightly wavy; a very characteristic row of interneural white subterminal spots, slightly dark-edged proximally; no terminal line; fringe rather paler.—*Hindwing* moderately ample, termen almost regular; slightly whiter than fore wing, at least proximally to the postmedian the lines in this area being almost obsolete except at abdominal margin; postmedian and subterminal more as on forewing, but weakening near costa.

Forewing beneath more suffused, but with hindmargin white; an elongate but weak cell-mark; the white line beyond the postmedian indicated; subterminal spots moderately developed. Hindwing with small cell-dot, alternations of lines (not very sharp) in proximal part, curved whitish postmedian band and weak, more or less macular subterminal.

Hpimaw Fort, 9th to 13th August, 1922 (type) August 1923 (paratype). Also known to me from Yatung (Indian Tibet), Nainital, etc.

Hitherto confused with *ochracea* Warr.; larger, more *Laciniodes*-like, with narrower white areas, more macular subterminal, etc.

* 120. *Eupithecia ustata* Moore

Eupithecia ustata Moore, Lep. Coll. Atk., p. 268 (1888) (Darjiling).

Htawgaw, April–May 1923, 1 ♂.

Differs chiefly from a figure in the British Museum (evidently crude) of Moore's type in having a proximal projection of the brown distal coloration into the median area (at base of M¹ and M² and entering the cell). Might conceivably be a race of the Japanese *sophia* Butl. (*Ann. Mag. Nat. Hist.* (5) i, 444).

121. *Eupithecia eupitheciata* (Walk.)

Phibalapteryx eupitheciata Walk., List Lep. Ins., xxvi, 1720 (1862) (Australia).

Htawgaw, July 1923, 1 ♀.

One of the most widely distributed Indo-Australian species, perhaps a migrant. I can add to the localities given by Hampson (*Faun. Ind. Moths*, iii, 398) Java, Sangir, Celebes, Dutch and British New Guinea.

* 122. *Chloroclystis consueta* (Butl.) (?)

Eupithecia consueta Butl., Ann. Mag. Nat. Hist. (5) iv. 449 (1879) (Japan).

Hpimaw Fort, June 1923, 1 ♂.

The specimen is worn, but seems to agree with my series of *consueta* from Yachiaolin, Central China, which in its turn may perhaps be a race of *rectangulata* Linn.

* 123. *Chloroclystis filicata* (Swinh.)

Eupithecia filicata Swinh., Tr. Ent. Soc. Lond., p. 1 (1892) (Khasis).

Htawgaw, June 1923, 1 ♂.

* 124. *Chloroclystis chlorophilata* (Walk.)

Eupithecia chlorophilata Walk., List Lep. Ins., xxvi. 1768 (1862) (Bhutan).

Htawgaw, early July 1923, 1 ♂.

* 125. *Chloroclystis papillosa* (Warr.)

Aëtheolepis papillosa Warr., Nov. Zool., iii. 124 (1896) (Khasis).

Htawgaw, July 1923; 1 ♂; Laukhaung, April–May 1923, 1 ♀.

¹ Unaccountably referred by Warren and Hampson to *Asthena* and *Hydrelia* respectively, really a *Eupithecia* pure and simple.

* 126. *Chloroclystis inaequata* (Warr.)

Sesquiptera inaequata Warr., Nov. Zool., iii. 126 (1896) (Khasis).

Htawgaw, April-May, 1923, 1 ♀; Hpimaw Fort, 14-18 August 1923, 1 ♀.

These seem to agree perfectly with a Khasi ♀ in the British Museum which I believe to be correctly mated with a ♂ from the same source. The ♀♀ in this group are, however, confusingly alike, keeping pretty closely to the facies of the genotype, *coronata* Hb., while the ♂♂ vary so remarkably in the secondary sexual modifications of the hindwing that Warren has founded upon them a number of genera.

* 127. *Chloroclystis rubroviridis* (Warr.)

Gymnopera rubroviridis Warr., Nov. Zool. iii. 127 (1896) (Khasis).

Htawgaw, June 1923, 2 ♀♀.

Known also from Sikkim and Indian Tibet and I believe from Borneo.

* 128. *Rhinoprora xanthocomes* sp. n. (Pl. 1, Fig. 24)

♂♀, 17-19 mm. Face pale green; frontal cone small. Palpus $2\frac{1}{4}$ to $2\frac{1}{2}$, second joint heavily scaled, third joint less than $\frac{1}{2}$ second; pale green, marked beneath with black. Vertex, thorax and abdomen green, fading to yellowish, beneath whiter; abdomen above variably sprinkled with black, in the strongly marked aberrations with a large black band which in repose probably meets that of the forewing. Fore and midleg partly darkened, leaving pale spots at ends of joints.

Forewing pale green, slightly bluish when fresh, but easily discolouring to dirty yellowish or whitish; basal patch scarcely darkened, but bounded by a straightish black line which is thickened at costa, weakened in middle, distinct at hindmargin; median area moderately broad, bounded proximally by a rather oblique, anteriorly gently curved blackish line, which is distinct throughout or strengthened at the veins, distally by a weaker line, which is very slightly angled on R^1 and R^3 , thence oblique inward but almost obsolete; a second line in the median area, parallel with the antemedian, often distinct, sometimes obscured by a more or less broad, band-like, brown and black shade which occupies the proximal part (in its narrower posterior half nearly the whole breadth) of this area; other median lines and shades only developed costally; subterminal line obsolete, its position indicated by small proximal spots between the radials, a larger, rather more proximal, costal spot and some very slight dark scaling in posterior half; terminal line black, interrupted at the veins; fringe dark-spotted opposite the veins and faintly bisected between.—*Hindwing* rather small, termen slightly prominent at R^1 , then slightly concave to R^3 ; in ♀ greyish, with traces of darker lines, in ♂ whiter, narrowed, the abdominal margin with some long ochreous hair distally, at tornus becoming somewhat distorted, an oval ochreous androconial patch near the end of M^2 and some less well-defined ochreous scaling before.

Both wings beneath pale, weakly marked excepting the costal area of forewings; sometimes with indications of a dark subterminal band on one wing or both.

Htawgaw, June, July and August 1923, 8 ♂♂, 2 ♀♀, also a large, discoloured, rather aberrant ♀ September-October; Sikkim, a ♂ in Coll. Brit. Mus. misidentified by Hampson as *trichophora* Hmps. n.; Khasis, a few in Coll. Tring. Mus. apparently mixed by Warren with *nigroviridata* Warr. (?); ? Hpimaw Fort, 9-13 August 1923, a rather dark ♀, more regularly marked across the forewing and with the median area scarcely narrowing behind.

Nearest *R. trichophora* Hmps. n. (*Faun. Ind. Moths*, iii. 393) but with the structure and colouring of the ♂ sexual modifications of the hindwing quite different.

129. *Calluga costalis* Moore.

Calluga costalis Moore, Lep. Ceyl., iii. 480, Pl. 206, fig. 1 (1887) (Ceylon).

Laukhaung, April-May 1923, 1 ♂, 10 ♀♀, July 1923, 1 ♀.

Probably the July specimen represents a second brood; only one of the early generation is as small.

The species is widely distributed and I am not sure that *cissocosma* Turn (Proc. Roy. Soc. Vict., xvi. 232, Queensland) is more than a form of it,

130. *Brabira artemidora* (Oberth.)

Melanippe artemidora Oberth., Et. Ent. x., 33, Pl. 1, Fig. 6 (1884) (Askold).
Htawgaw, March 1923, 1 ♀.

Distributed from Japan and Formosa to Northern India. Should the Indian form prove racially tenable, the name of *pallida* Moore will be available for it.

* 131. *Brabira atkinsoni* Moore.

Brabira atkinsoni Moore, Lep. Coll. Atk., p. 271 (1888) (Darjiling).

Hpimaw Fort, 9-13 August 1923, 1 ♂, 14-18 August 1923, 2 ♀♀.

All are weakly marked. The species (Sikkim, Assam) seems very variable, or perhaps embraces two or three close allies, as yet not differentiated.

* 132. *Microloba bella diacena*, subsp. n.

♂ ♀. Distinguishable from *M. b. bella*, Butl. (*Ann. Mag. Nat. Hist.* (5) i, 448, Japan) by the entire, or almost entire, suppression of the dark markings of the central area of both wings, leaving only the costal markings of the forewing, cell-spots and terminal patches of both wings and sub-basal mark of hindwing. Even on the underside the postmedian line and its band-like proximal shading are much less strongly developed than in the Japanese race, though the hindwing generally shows rather strong costal spots.

Hpimaw Fort, June 1923, type ♂, 14-18 August 1923, allotype ♀. Also known from the Khasis and West China, its racial distinctness hitherto overlooked.

The name-typical form inhabits Japan, Corea and East Siberia and, so far as I have yet discovered, the Formosan form is not separable. I have not yet seen examples from Central China where more or less intermediate forms might be looked for.

* 133. *Heterophleps clityogramma*, sp. n.

♂, 37 mm.; ♀, 35 mm. Closely similar to *sinuosaria* Leech (*Ann. Mag. Nat. Hist.* (6) xix. 548; Prout in *Seitz Macrolep.*, iv. 188, Pl. 11 c, Ta-chien-lu)—Forewing slightly less warm brown, yet without the violet-grey tone of the allied *grisearia* Leech; cell-mark and the dark spot which precedes it on costa enlarged; antemedian line much more oblique, more swollen into spots at its two outward angles. That on M not quite 4 mm. from base, that on SM² 5-6 mm. from base.—Hindwing paler than in *sinuosaria*, at least anteriorly.—Forewing beneath with the cell-mark elongate.

Tali, Haut Yunnan, type ♂, Hpimaw Fort, 9-13 August, 1923, allotype ♀.

The ♂ was sent me several years ago by my late friend Thierry-Mieg as '*grisearia* Leech' and no doubt others from the same source will be found in his collection, now in the Paris Museum. Both my examples are somewhat rubbed, but their agreement with one another and distinctness from *sinuosaria* are abundantly manifest.

* 134. *Cryptoloba aerata* (Moore)

Larentia aerata Moore, Proc. Zool. Soc. Lond., p. 654 (1867) (Bengal).

Hpimaw Fort, 9-13 August, 1923, 1 ♀.

* 135. *Cryptoloba togulata* sp. n.

♀, 23 mm. Face dark fuscous. Palpus about 2, terminal joint concealed; brown, mixed at tip with whitish. Antennal pectinations rather long and lax, vertex and thorax fuscous, mixed with brighter brown; abdomen and underside paler and greyer.

Forewing with costa arched, apex acute, termen straightish or very faintly sinuate anteriorly, rounded about R³, thence more oblique; areole very large, undivided; pale indefinite buff, with slight dark irroration (small strigulation); rather large blackish spots beyond $\frac{1}{3}$ and beyond $\frac{2}{3}$ costa, the former giving birth to a straightish, band-like shade which limits a distinctly darkened proximal area, the latter marking the commencement of slighter and more ill-defined terminal shading, which is narrower posteriorly than anteriorly and has its proximal boundary curved; a much smaller and weaker dark spot on costa near apex; a moderate terminal spot behind SC⁵ and faint indications of others at midtermen and near tornus; cell-spot faintly indicated near proximal side

of pale median area.—*Hindwing* narrower than in *aerata* Moore, rather approaching that of the *cinerea* group, the angles rounded (especially tornus), the termen straightish or very slightly sinuate between SC^2 and R^3 ; DC strongly oblique posteriorly, SC^2-R^1 well stalked, R^2 central (or slightly before middle), M^1 about connate, SM^2 wanting; median area at least as pale as on forewing, proximal area little darkened, with a large dark cell-dot at its outer edge, terminal dark shade rather broad but not solid, only its proximal line (narrow band) and apical part well developed.

Underside similar, but with the forewing more blarred.

Hparê, September 1923, the type only.

Possibly related to *olivescens* Hmps (Journ., Bombay Nat. Hist. Soc. xiv, 511) but very distinct. In spite of differences in size, shape, colour and hindwing venation it is just possible that it will prove to be the ♀ of *trinotata* Warr. (Proc. Zool. Soc. Lond., 1893, p. 344, Sikkim), of which the ♂ type remains unique.

* 136. *Cryptoloba cinerea* (Butl.)

Lygranoa cinerea Butl., Ann. Mag. Nat. Hist. (5) vi, 228 (1880) (Darjiling).

Htawgaw, 22 August, 1922, 1 ♂, April–May 1923, 2 ♂♂, 1 ♀, June 1923, 1 ♂; Hpimaw Fort, June 1923, 3 ♂♂, 14–18 August, 1923, 1 ♀.

Capt. Swann's examples do not appear to differ racially from the Sikkim form. On the other hand Hampson (*Faun. Ind. Moths*, iii, 337) differentiates a very marked 'form' (race) from the North-West Himalayas.

137. *Cryptoloba etaina* Swinh.

Cryptoloba etaina Swinh., Ann. Mag. Nat. Hist. (7) vi, 310 (1900) (Khasis).

Hkamkawn, June 1923, 1 ♀; Hpimaw Fort, June 1923, 1 ♀.

Perhaps a race, or even a new but very closely related species. The ground-colour of the forewing slightly more recalls that of *olivaria* Swinh. (*Ann. Mag. Nat. Hist.* (6) xix, 165) and the postmedian line has both the central curves slightly strengthened, the fine black line (the true postmedian) which accompanies the brown shade rather more interrupted than in Swinhoe's type of *etaina* and the few other examples (likewise ♂) which I can compare. The unicolorous forewing beneath, with the postmedian line costally indicated in black, the different fringes, etc., remove it from *olivaria*.

138. *Atopophysa indistincta* (Butl.)

Scotosia indistincta Butl., Ill. Het., vii, 118, Pl. cxxxvii, Fig. 19 (1889) (Dharmasala).

Hpimaw Fort, June, 1923, 1 ♂.

Widely distributed in the Himalayas and across China to Ningpo. A race or close relative is described as *A. i. opulens* Prout (*Ent. Mitt.*, iii, 247, Formosa).

(To be continued)

THE COMMON BUTTERFLIES OF THE PLAINS OF INDIA
(INCLUDING THOSE MET WITH IN THE HILL STATIONS
OF THE BOMBAY PRESIDENCY)

BY

T. R. BELL, C.I.E., I.F.S. (Retd.)

PART XXXVI

(Continued from page 837 of Volume XXX)

Sub-family (7)—ERYNNINÆ.

Imago.—Fast flying insects with the exception of *Baracus* and all keeping generally to the trees, flying high: again with the same exception. Medium-sized insects too, but there are some that are well on the small side such as *Telicota mæsoïdes* and *Padraona gola*. and *Baracus* again—although this last is larger than either of the other two mentioned. This last is in colour and markings like *Taractrocera nicévillei*: uppersides brown with small subapical subdiscal spots. *Telicota*, *Padraona* and *Cupitha* are black or blackish-brown banded transversely with yellow, either orange or paler. *Halpe* are nearly all dark-brown above with subapical and discal semihyaline white spots at end of cell, in interspaces 2 and 3 and 6, 7, and 8. On the undersides of the hind wings *Halpe* and *Baracus* are brown or yellow with some spotty variegation; *Telicota*, *Padraona* are yellow with bands as on upperside; *Cupitha* is a pure yellow with large black spaces.

Antennæ.—Generally with the club having the apiculus bent at right angles and short, sharp; in *Baracus* bent slightly in a curve and in *Cupitha* not quite at a right angle.

Palpi.—With the second joint half-laxly scaled; the third joint rather minute and hidden in *Cupitha*; somewhat prominent in the others except in *Baracus* where it is most prominent; in all it is porrect or erectly-porrect.

Hind tibiæ.—Fringed in nearly all, but sometimes only slightly (seemingly naked in some species of *Halpe*); with two pairs of spurs.

Fore wing.—Vein 12 reaches costa before end of cell; cell less than two-thirds the length of costa; 3 from close to lower end of cell; 2 very variable.

Hind wing.—Vein 7 emitted a little before upper end of cell; 5 obsolete; disco-cellulars faint; 5 absent or very faint; 3, 2 from close to lower end.

Egg.—The egg of this division or sub-family is generally a dome-shaped structure, much higher as a rule than half the breadth; without any basal band or foundation; generally without meridional ribs, quite smooth to the eye, often somewhat reticulate or rough under the microscope; generally whitish in colour but never purely white, always slightly soiled-white, sometimes with a pinkish shade of a very pale intensity. The egg of *Baracus* is meridonally ribbed with some 17 very fine ribs; that of *Cupitha* is of the same description as those of the *Platingiinae* and does not really fit in here at all—it is very strongly ribbed and is red in colour; all the rest being whitish, either plain or with red blotches (*Baracus*) or tiny red dots.

Larva.—The shape very ordinary, thickest in middle, fining to anal end which is broadly rounded; and to the neck which is narrow compared to the head; the head however really rather small, round or semi-elliptical, hardly bilobed with the surface rough and covered with appressed, fine, minute hairs not in any way obscuring the sculpture or the colour. Surface covered with very minute, erect, light hairs, longest round dorsoventral margin and, especially on anal segment; this anal segment is nearly all roughened on dorsum with little tubercles, each bearing a hair. *Spiracles* of segment 2 particularly large. Colour some shade of green, the skin thin and the tracheal tubes showing through; generally without longitudinal lines except in *Padraona gola* and *Halpe honorei*—this last has a broad, longitudinal, red, lateral band; always a collar to segment 2.

Pupa.—With the front broad, somewhat rounded; a slight boss between the eyes on frons; thorax slightly humped; cremaster generally with two points behind between which is a line of longish shafts with hooked ends. *Surface* covered up with tiny hairs, sometimes bristle-like and longer on eyes; cremaster with the sides hollowed out, the extensor ridges dorsally and ventrally strong, the dorsum between these concave. *Spiracles* of segment 2 with very large, funnel-shaped, generally brown or red-brown expansions. *Colour* generally that of an old bone, sometimes with a green shade, sometimes strongly brown; covered with a cereous, white powder as a rule.

Habits.—Eggs laid on under surface of leaves. Larvæ make cells by joining the edges of the leaf or blade at the point. The cell eventually cut free and falls to the ground; always lined inside with silk and, generally, white cereous powder from the pupa dusts the sides. Larva eats bamboos and grasses and, exceptionally, only in the case of *Cupitha*, deciduous trees which is altogether abnormal. But, any way, *Cupitha* makes a bad fit in the sub-family whatever way it is looked at. The only thing that can be said is that the genus fits better into it than into any other.

Genus 25.—TELICOTA.

There are three species of this genus in British India according to Watson: *bambusæ* (M.); *augias* (L.) and *palmarum* (M.). Swinhoe reverts to Mabilie's *Corone* for *palmarum* owing to the position of veins 2, 3 of the fore wing; and he gives twelve other species (of *Corone*) as existing in the Malayan sub-region. There are four *Telicota* from the same area. The three Indian species are all dark-brown or blackish insects with golden-yellow or orange, transverse bands of contiguous spots on the fore and hind wings on the upperside and with the undersides suffused with yellow in the males, the bands showing through it more or less; the females having narrower bands in two of them while the female of *palmarum* is all brown with the markings duller, not yellow at all. They are all medium-sized insects of powerful flight, very quick on the wing; they rest on the tops of leaves, taking short flights, returning constantly to the same place.

Imago.—As above.

Antennæ.—More than half the length of the costa of fore wing; club stout, with a short, pointed, terminal crook.

Palpi.—Uprturned, the second joint somewhat laxly scaled; third joint sub-erect, somewhat prominent, bluntly conical.

Hind tibiæ.—Fringed and with two pairs of spurs.

Fore wing.—Vein 12 ends on costa some distance before end of cell; upper discocellular minute, middle and lower in an inwardly-oblique line, the middle the longer; vein 5 from their junction rather close to lower end of cell, vein 3 nearer to the end of cell, quite close to it, in the female, further from it in the male (in *Corone* closer still to end in both sexes); vein 2 a little beyond the middle in the male, rather nearer the end in the female for *Telicota*, before the middle in *Corone*; costa very slightly arched before its middle; cell rather long, very little less than two-thirds the length of costa; apex sub-acute; outer margin convex, somewhat shorter than the hinder margin; wing narrow, the oblique outer margin giving the outer portion of the wing a narrow, triangular shape.

Hind wing.—Vein 7 from well before upper end of cell, the cell-margin bent down from base to cell-end; discocellulars outwardly-oblique, faint; 5 absent; 3 from close to lower end of cell, 2 from twice the same distance from margin of 3 in *Telicota*, close together in *Corone*. All male *Telicota* have a linear discal stigma on the upper side of fore wing, extending, in *bambusæ*, *augias* and *palmarum*, from the lower end of cell to the middle of the sub-median vein with veins 2, 3 passing across it, dividing it often into three sections; this stigma bluish-greyish in colour, running through the black band between the basal and discal orange markings. In *maesoides*, the male has a short, glandular, black streak along the sub-median vein across the black between the same markings.

Egg.—High dome-shaped. The surface unsculptured. Colour whitish with a slight pink tint.

Larva.—Of ordinary shape, broadly rounded at anal end, the dorsum of anal segment roughened; the neck small, compared to the head which is round and yellow or black with a rugose surface covered with tiny, appressed

hairs that obscure neither sculpture nor colour. Surface of body covered with minute, erect hairs, longer round the dorsoventral and anal margins; the skin thin showing tracheal tubes underneath. *Spiracles* of segment 2 very large; of segment 12 smaller; the rest smallest. Colour green; unmarked.

Pupa.—Broad in front, with a well-developed cremaster; of ordinary shape; the cremaster with strong extensor-ridges and ending in a truncated extremity with a small tooth at each corner and with a fringe of longish shafts, either hooked or not, at ends; head with a slight convexity between eyes; proboscis slightly produced free beyond ends of wings. Surface of pupa covered with short, erect hairs, denser and stronger on eyes. *Spiracles* of segment 2 with very conspicuous, large, strainer-shaped expansions. Colour brownish and covered with a cereous, white excretion.

Habits.—Egg laid on underside of leaf. Larva makes a cylindrical cell at point of leaf by joining edges; small at first; new cells as required; eventually, when about to pupate, cuts it free upon which the cell falls to the ground. Butterfly is a powerful flier, keeping to the higher parts of trees.

215. *Telicota bambusæ*, (M.).—(Pl. M, figs. 75♂, 75a♀).—Male. *Upperside*. Fore wing with the ground colour black or very dark-brown that is nearly black; but every interspace between the veins from base outwards to stigma, continued along costa above stigma and along inner margin below filled in with light golden orange leaving a band of ground colour all along the outer margin through which the ends of the veins that come through the orange are slightly orange-bordered at where they leave it; the vein forming the bottom of cell and vein 1 and often, also, the vein forming top of cell are black through the orange from base and all veins from the stigma outwards through the orange (but not inside stigma) are thinly black, including 8 to 11 on the costa; the black band containing the stigma is slightly broader than the stigma itself and is continued beyond it outside the cell to vein 6, embracing the inner halves of interspaces 4, 5. Hind wing. Ground colour black; the markings also golden-orange—a large spot towards end of cell disconnected from much larger, quadrate—elongated spots forming a transverse band in interspaces 2, 3, 4, slightly extending upwards into interspace 5 and confluent across interspace 1 by a broadish neck with a yellow streak that borders the outer edge of vein 1b from base to outer margin—the transverse band about 2.5 mm. in width. On the fore wing, under the median vein there is a slight covering of decumbent, orange hairs from base outwards to vein 2 and a similar covering over the whole of the orange mark along inner margin under vein 1 but there is no fringe, or very little, along the inner margin. On the hind wing there are quite a lot of longish, orange hairs in the cell, underneath it in interspace 1b right out to outside edge of the transverse orange band and in interspace 1a still longer, denser hairs, not quite so decumbent, right out to outer margin. *Underside*. Fore wing golden-orange but the posterior spots of the discal spot-band much paler, showing through; the base of the wing in the cell to about one-third its length and below it right out to the inside edge of the discal band including the whole of interspace below vein 1 along inner margin and outside discal band upwards to vein 2, all black; the bases of interspaces 4 and 5 are very slightly blackish up to half way out, the costa is thinly black, the marginal black band of upperside shows through also slightly blackish and there are blacker small spots, one in each interspace 2 to 8 limiting the blackish outer border from the orange discal band. Hind wing orange, the discal band showing through golden-orange, little blackish border-spots in each interspace limiting the discal band outside and inside. *Cilia* above yellow with basal third brown, the actual border of the wing finely black, showing very distinctly on the undersides; below the very bases of cilia are yellow and the middle rather broadly tinged brown.—Female. *Upperside* not so black; the orange markings reduced a good deal in size. Fore wing with the base of cell black extending outwards along bottom of cell to middle of it which is again black; the black band that bears the stigma in male extends right in to the base in interspace 1 leaving only a streak of orange above vein 1 and touching it; the edge of costa black right down to base whereas, in the male, it is black only in the outer half. Hind wing as in male. *Underside* like the male but the black more extensive, following the upperside in the fore wing; in the hind wing the colour is more dull-yellow than in the male, the discal band perhaps rather more conspicuous. Antennæ black above, orange beneath except a narrow black band at edges of joints,

the club with the end half, including point dull-red below; palpi, head, thorax and abdomen above concolorous with the orange of wings, the long hairs of thorax mixed with brown ones imparting a greenish shade, the abdomen banded black; below uniform bright-ochreous, the tibiae and tarsi orange. Expanse 35 mm.; the male rather less.

Egg.—The *shape* is that of a high dome with a very short, cylindrical base of 0.05 mm. in height above which is the broadest part; top somewhat flattened and 0.44 mm. in diameter, a circular area. *Surface* only slightly shining with some minute, irregularly-disposed and irregularly-shaped rugosities; the micropyle small, apical, nearly smooth, darker than the rest. *Colour* white with a delicate pink shade. B: 1.05 mm.; H: 0.85 mm.

Larva.—The *shape* is that of *Telicota*, *Halpe*, etc.: body cylindrical but with the ventrum rather flattened as the legs and prolegs are short and the caterpillar lies rather closely applied to the resting-surface; the anal end broadly and semicircularly rounded, the anal segment rather large, somewhat flattened on dorsum, very slightly bordered tumidly round the free edge and considerably roughened dorsally on the whole surface, with the usual six comparatively long hairs from the free edge all round as well as others much shorter (which cover the whole body); segment 13 about one-third the length of segment 12 but quite well-distinguished and rather waved as to hinder margin; segment 12 about equal to segment 14 in length, perhaps even shorter; segment 11 coequal with 12. *Head* nearly round with a distinct, triangular sinus on vertex, the face not very convex; the true clypeus just about one-third the height of head, an isosceles triangle, acute at apex; the false clypeus outside it with straight margins quite parallel to those of the true clypeus all the way to base, reaching about two-thirds the height of face, acute-apexed; surface of face very rugged seen under the lens, covered with more or less decumbent, very short, fine and lightish hairs, with some slightly longer about mouth-opening; the labrum transverse, reddish-shining; the ligula white, black-bordered, not very large with a frontal, triangular sinus; the antennae more or less colourless in both joints; the mandibles dark-brown with reddish bases when exposed; the eyes brown; the general colour of the head is soiled whitish obscured, however, by a dark red-brown band along the hinder border though not quite reaching it, this band widening out below to include the whole of the eye-area and joining on to a broad, similarly coloured band down the middle of face (dorsal) that splits down the false clypeus, including it but not the true clypeus which remains the ground-colour except for a somewhat bold dorsal line from apex to base; thus all that remains of the white ground-colour is a large oval space on each lobe from vertex to base and the true clypeus. *Surface* of body covered with extremely minute, erect, light, fine hairs all over amongst which the subdorsal one and a spiracular one, arising each from a small, dark tubercle, only are seemingly twice as long; the usual transverse fine impressed lines towards hinder margin of each segment; otherwise surface dull and smooth. *Spiracles* somewhat small, light yellow or yellow; those of segment 2 very large, outlined very thinly dark; those of segment 12 smaller than those considerably but larger than the rest; all flush or nearly so and rather broad. *Colour* grass-green of a medium shade with segment 2 rather whitish, segment 14 somewhat soiled; the ventrum and legs, both true and prolegs, lighter. The skin is translucent and shows all the tracheae through as well as the pulsating, paired, yellow bodies dorsally on segment 10 (in the male?). L: 24 mm.; B: 4 mm.

Pupa.—In *shape* the body is cylindrical, slightly broadest at shoulders whence the decrease in diameter is extremely gradual to the anal end, segments 11-14 decreasing cone-wise, the apex of the cone, segment 14, being composed of a very short, transverse, annular piece, (i.e., very much shorter than broad) with a large, trapeze-shaped, flattened piece proceeding from its end horizontally backwards, broadening to end, the dorsum of this trapeze with a strong ridge on each side (sub-dorsal or dorsolateral), these two ridges diverging backwards and each ending in a flat deltoid tooth—or it would be more correct to say that the trapeze-shaped piece bears two ridges, diverging, losing themselves towards the extremity of the piece or pupa and not quite forming the lateral margins of it; the square or straight end of the piece is fringed with a close row of stiff, hooked bristles which are reddish in colour and nearly one-third as long as the whole trapeze; segment 13 a very short, transverse piece; very much shorter than 12; 12 a little shorter than 11; *head* with the vertex parallel to the axis of body, half as long again as segment 2 which, with the front

slope of thorax and head-vertex lies in a plane inclined to the longitudinal axis of the pupa at an angle of about 30° ; the frons of the head in a plane at right angles to the longitudinal axis, rather broad and little convex forming the front of the pupa; the eyes prominent; the margins of head and segment 2 quite straight; thorax hardly humped at all, nearly parallel to axis of pupa in its hinder two-thirds with the rest of the dorsal line of abdomen; the hinder margin of thorax a quarter-circle curve meeting the wing-lines in a not very broadly rounded angle of nearly 90° ; segment 4 slightly longer in dorsal line than 5 which, again, is only half the length of 6; proboscis produced free beyond wings to not quite the middle of the succeeding segment 9. *Surface* of pupa distantly shallow-punctate and fine-aciculate-lined transversely; the bevelled edges of segments 8-11 non-existent in front (segments 9-11) and longly-sloping, smooth behind (segments 8-10), the sloping surfaces over-hung by a single row of reddish bristle-like hairs projecting from their anterior limits backwards over them and about as long as the surfaces. *Spiracles* of segment 2 indicated by a thick, raised half-funnel (semicircular) occupying the front margin of segment 3 and as broad as half length of segment 2, as long as the same segment (twice as long as broad that is) and the same colour as pupa with a plush-like bloom on it in certain lights; the rest of the spiracles small, oval, nearly flush, slightly lighter in colour than the rest of the pupa. *Surface*, besides, covered with short, erect, reddish hairs all over, the eyes fringed before and behind, the hairs all extremely short, longer on anal segment and eyes. *Colour* burnt-sienna brown, reddish all over, the cremaster darker, the thorax duller, the wings like the thorax, L : 18 mm. : B : 4.5 mm. nearly.

The head is slightly bowed and little less in transverse diameter than the shoulders which are the broadest part of the pupa; the apex of thorax is the highest point.

Habits.—The egg is laid always single, near the point or at the very point of the leaf on the upperside. The little larva makes a cell by lying on the midrib at the point and drawing the edges of the leaf together over it; and at each stage, the larva makes a similar cell, each time to suit it in size, lining the inside with silk. The pupa is formed in a cell of two or three leaves drawn together to make a cylindrical house which fits closely to the pupa, and is dusted with a cereous excretion from the larval body that makes its appearance first between the first pairs of prolegs and before the first. The pupa is of course covered with this and is attached by the tail inside lightly to a thicker pad at the closed end of the cell—one end remains, though closed outwardly, unfastened by silks through which the emerging butterfly can easily force its way. The different stages and states of the larva and egg, larva and pupa are normal in duration at all times as far as one has been able to observe.

This is mostly an inhabitant of regions of fairly heavy rainfall; it is common wherever there is bamboo-growth which is practically confined to places where the monsoon is strong. It thus comes about that it does not exist in the absolutely open country and mostly is found in hills and jungles. Swinhoe, in his *Lepidoptera Indica*, vol. x, p. 249, gives the habitat as India, Burma, Ceylon, Andamans and the Malay Peninsula; says that the types are from Calcutta and that it is a very common species all over India up to the Himalayas and has been recorded from many localities, adding, "our figures of the larva and pupa are from Davidson's original drawings not previously published". These together with the butterflies, are contained in the book on Pl. 813, fig. 3, the male, 3a the female and 3b the underside, with 3c the larva and pupa, these last two very bad and sausage-like reproductions of the originals.

The figures in coloured Plate M accompanying these papers of the male, 75 and the female, 75a are good enough in the shape and accuracy of the markings, but the female is too pale as compared to the other; the *cilia* are too pink in places, or red; the underside of the male hind wing is too grey—it should be yellow; the spots too red and the body is altogether too dark.

216. *Telicota augias*, (L.).—Male, female. Like those sexes of *Telicota bambusæ* except that they are paler all round and, perhaps, a trifle larger. The main difference appears to be that the golden-orange of *bambusæ* is here golden-yellow, extending from the outer edge of the discal band along upper sides of the veins to practically the outer margin on the fore wings.

Swinhoe states in *Lepidoptera Indica*, vol. x, p. 249, that, in *bambusæ*, there “is a blackish streak near the anal angle on the underside of the hind wing in each of the two anal interspaces.” That is certainly not present in a large series bred in the Kanara District. The two species are so extraordinarily alike that anybody might be excused for mistaking them for one and the same if it were not for the difference in genital armature discovered by Elwes and Edwards (*Trans. Zool. Society* of the year 1897, p. 251, Pl. 25, Figs. 62, 62a and 63) confirmed lately by Mr. W. Ormiston, F.E.S. in a paper entitled “Genitalia of some Ceylonese Hesperiadæ” published in the *Report of the Proceedings of the Third Entomological Meeting* held at Pusa, February 3, 1919, vol. iii, p. 1024, Pl. 171, Figs. 25 and 26. All that appears to be known of the earlier stages of *augias* are the remark that “the larva feeds commonly on cane-leaves and is also said to feed on bamboos and rice rarely, but is scarcely a pest” contained in the same Report, vol. I, p. 116 in the second paper read, entitled *Annotated List of Indian Crop-Pests by the Imperial Entomologist*, Mr. T. Bainbrigge Fletcher, R.N., F.L.S., F.E.S., F.Z.S. The ‘cane’ is sugar-cane, a grass; not *Calamus*, the cane that is of the *Palmaceæ*.

Swinhoe gives the habitat as Sikkim, Assam, Central and Southern India, Burma, Siam, Andamans, Ceylon, Hongkong, Formosa, Malay Peninsula, Malay Archipelago and says “A common species in most of the above-mentioned localities; we took many specimens in Mhow, Poona and Bombay; Elwes records it also from E. Pegu and the Naga Hills; Manders from the Shan States; Druce from Siam; Watson from the Chin Hills; Aitken from Nasik and Betham from the Central Provinces.” It is common above the Western Ghats in the Dharwar District also, but it is not to be found in the N. Kanara District either below the Ghats or where there is very heavy rainfall—the rainfall varies from 400 inches on the crest of the Ghats to 100 and over on the sea coast, and 30 inches or even less on the eastern limits. It is probably altogether confined to the open country where rainfall is light.

217. *Telicota* (=Corone, Mabille) *palmarum*, (M.).—Male. *Upperside*. Fore wing—golden-ochreous, colour very similar to that of *augias*, the costa and all the veins and a line through the middle of the cell blackish-brown; outer-marginal border blackish-brown, broadest at the apex and hinder angle, its inner margin sinuous and curving inwards opposite the cell, then gradually widening hindwards; a blackish-brown streak from the base running immediately below the median vein and expanding gradually outwards, narrowing at

the cell-end and expands into a patch outside the cell which reaches the costa, being somewhat constricted in its middle; the submedian vein heavily blackish-brown. Hind wing blackish-brown, some golden-ochreous hairs at the base and along the abdominal area; a somewhat large, golden-ochreous, oval spot at end of cell and a somewhat broad, short, discal band of the same colour, divided by the blackish-brown veins, from a little above vein 6 to near the internal vein; with a thin, golden-ochreous streak running from this end to the margin. *Cilia* of fore wing blackish-brown; of the hind-wing golden-ochreous. *Underside* orange-ochreous. Fore wing with the basal half of the cell and all the basal space below it dark blackish-brown; this colour running narrowly to the cell-end and along the hinder margin leaving a costal streak, the outer half of the cell and a discal band from the costa orange-ochreous curving well outwards, then inwardly-oblique to near the hinder margin, divided by the veins into elongate, subquadrate spots which increase in size hindwards; costa and outer-marginal lines finely blackish-brown. Hind wing uniformly orange-ochreous, darker than the fore wing, the discal band as on the upperside, divided by the veins and finely edged on both sides with blackish-brown. Antennæ black, the upper half of shaft with minute, ochreous dots on its underside and a smear of the same colour on the lower half of the club; palpi ochreous and blackish-brown; head dull-ochreous with a transverse brown band in front and another behind; thorax with ochreous setæ; abdomen with indistinctly similar-coloured bands; on the underside: palpi, body and legs are of the colour of the hind wing.—Female. *Upperside* chocolate-brown, the markings dull brownish ochreous. Fore wing with a large spot at end of cell and a narrow band divided into spots by the veins, commencing very narrowly on the costa by the apex, angled almost acutely outwards, then oblique and almost straight to near the middle of the hinder margin, gradually broadening hindwards. Hind wing with indications of the small cell-spot and with a discal band, much narrower than the discal band of the male. *Cilia* of fore wing brown, of hind wing ochreous-white. *Underside* paler and duller in colour; markings similar. Expanse: male 40-42 mm.; female 45-50 mm.

Larva.—Feeds on date and cocoanut palms (*Indian Museum Notes*, V, No. 3, p. 126, Pl. 9, larva) ' . . . there is no picture of the larva in that plate, only the male and female butterfly are shown.

Habitat.—Sikkim, Assam, Bengal, Burma, Nikobars.

Distribution.—The types come from Calcutta; it is a widely spread but rare species; our description and figures are from a male in our collection from the Khasia Hills and two females from Maulmein in the Hankook Museum, Newcastle, taken by Adamson in January 1879; Elwes records it from the Karen Hills, Margharita, in Upper Assam and from Perak; de Nicéville from Sikkim and Cachar; Watson from Beeling in Burma and the Nikobars.'

In the *Indian Museum Notes* under the reference noted above, it is definitely stated that this insect was bred in Calcutta in December 1900. Hence it may well be found in the Plains somewhere else.

218. *Telicota mæsoïdes* (Kollar).—Male. *Upperside*. Fore wing with the cell ochreous-orange, rather lighter than the golden-orange of *bambusæ*, the costa above it similar as far as the basal half of interspace 9, the rest of that interspace blackish-brown, the upper and lower borders of cell also brown at base of wing; interspace 1 below the cell from base sprinkled rather thickly with ochreous-orange scales out as far as to and below the junction of vein 2; interspace along the inner margin below vein 1 pale-yellow at base becoming ochreous-orange outwards and finishing about three-quarters the way to tornal angle; a postmedial band of ochreous-orange spots filling the interspaces 1 to 8 from vein to vein, the lowest spot longest, triangular, its outer border very slightly excavated and more or less in a line with the end of the yellow streak along the inner margin of wing, its apex on vein 1 and reaching to about 1.5 mm. of the basal, ochreous-orange powdering of scales; interspace 2 with the spot quadrate, about the middle of interspace, outer side slightly excavated, inner slightly triangular, about 1.5 mm. broad; the one in interspace 3 rather more than half the width of last and about as long as last but very slightly dislocated outwards, square on inside edge, slightly excavated

on outer; in interspaces 4, 5 narrower, shorter, their inner edges in a straight line and dislocated outwards about two-thirds the length of the spot beneath (in 3), their outer edges also in a straight line and also very slightly excavated; above this, in 6 and 7 the spots fill the basal halves of interspaces so that their broad outer edges only reach slightly beyond the inner edges of those in 4, 5, their pointed inner ends not quite reaching end of cell; above again in interspace 8 there is a shorter, quadrate spot filling more than the middle third of the space; all this leaves the ground colour showing as a broad, brown outer border and a thinner discal band broadest below, narrowing in base of interspace 2 and still more in base of 3 to widen out considerably in 4 and 5 which is connected by a very narrow neck at upper, outer angle of cell with a small triangular brown portion on costa composed of the base of interspace 8 and the outer half of 10 with a little of the outer end of 11; there are a few ochreous-orange, decumbent hairs under the bottom of the cell inside the junction of vein 2 but there is hardly any fringe of hair along inner margin. Hind wing: similar in colour to fore wing with the markings of a similar colour:—a more or less blurred, smallish spot in base of interspace 7 and another slightly larger half way out; a larger clear one in end of cell but not touching discocellulars or upper or under margin; a broad, discal band, also clear consisting of a spot in each interspace 2, 3 not quite reaching their bases and stretching out to nearly two-thirds their length, these the largest with, above, another quadrate, smaller, dislocated well outwards in interspace 4 and a bit of 5 and, in interspace 1*b*, another about half the length of the largest, placed about the middle of the lower side of the largest and connected with a similar one in interspace 1*a*; finally a streak below vein 1*a* a good deal longer than the last spot; in the cell there is a tuft of ochreous-orange hairs in the base and some more, decumbent, in the rest of cell; also some sparsely distributed over the surface below the cell and outwards and a ridge of longer hairs below vein 1*b* from base to the end or outside of the discal band. *Underside*. Fore wing similar to upperside but duller, the whole base of cell, the whole of interspace 1 and the whole of that below vein 1 along inner margin blackish-brown except the lowest spot of the discal band which shows through; this brownish-black continued upwards to occupy the base in interspace 2 inside the discal band-spot and further up still into the base of interspace 3 to be continued as a mere line along the discocellulars to upper end of cell; outside this, further, there are slight brown borders limiting or indicating the inside and outside limits of the band-spots in the interspaces 2 to 8, the whole of the blackish-brown intervals of the upperside being here covered with duller orange scales right out to termen and right up to costa except in interspace 2 where the outside limit of the band-spot is a triangular continuation of the black of inner margin and end of interspace 1; the termen of the wing is finely black. Hind wing. Ochreous-orange but the discal band as well as the cell-spot and spots in interspace 7 paler than on the upperside but just as large, all the rest of the wing powdered thickly with orange scales, each inside and outside border of all spots rather less thickly than the remainder of surface producing a mottled-spotted appearance; but the two anal interspaces beyond the discal band immaculate blackish-brown in a patch not reaching termen nor reaching abdominal margin and the termen is finely black from costa to vein 2. *Cilia* of fore wing above brown with tips of those in the vein-interspaces orange becoming more orange towards tornal angle; of hind wing nearly completely orange, except the extreme base and at ends of veins 2, 3 which are brown; underside of hind wing orange without brown bases but ends of 2, 3 very slightly brown; forewing with the cilia orange in interspace 1 and at angle, the rest with the basal half yellow, the outer half brown except extreme tips which are alternately brown and yellow. Antennæ black, banded broadly yellow below on shaft and having the whole of club yellow below except the extreme crook which is dull-reddish; palpi yellow below, the third joint and top of the second brown; head and thorax above mixed orange and yellow with some brown hairs; abdomen brown mixed with some orange hairs; beneath: yellow with a slight green shade, end segment of abdomen and legs more orange.—*Female*. *Upperside*. Both wings exactly like the male but with the blackish-brown more extensive, the spots therefore rather smaller, the cell of fore wing with upper border more prominently brown and with a brown line from base half way towards its end through the middle. All the rest as in male. Expanse 25 mm. for female; the male a little less.

Larva.—The body is of the *shape* of that of *Telicota bambusæ* except that the anal segment is rather longer-semicircular and overhangs the anal claspers behind, is about the same length as segment 12, is roughened on the dorsal surface though not tuberculate and is covered with minute, erect, soft, light hairs and there are eight much longer hairs (about as long as segment 13) round the free margin with a few shorter ones between them, all these marginal hairs pointing straight out at right angles to the margin; segment 13 is a parallel-sided, transverse piece between half and one-third the length of segment 12; the body is thickest in the middle but is rather slight throughout; there is a distinct neck and the head is, as usual, large in comparison to it; the body reminds one of the larva of *Parnara kumara* because the colour is whitish light-green and there are seven parallel lines across each segment starting at hinder margin and ending at about three-quarters the length forwards: these lines all parallel to the segment margins—transverse to the length of the body as usual; *head* triangular in shape—another difference to *Telicota*—and thick, the surface shallowly rugose but not honeycombed; the true clypeus reaching less than half the way towards vertex of head, longly triangular, the false clypeus reaching slightly more than half way up, also triangular, both with the apices acute, the strip formed by the latter nearly as broad as the true clypeus; the labrum narrow, short, red-brown; the ligula nearly circular with a triangular indentation in front margin and depressed from point of this sinus backwards, colour soiled, watery white; antennal, basal joint like the second joint: colourless tinged with red-brown; mandibles red-brown margined black, the cutting edges straight; eyes: four very close together in a curve, the 6th in continuation of 3, 4 but spaced twice as far from 4 as 4 is from 3; 5 forming an equilateral triangle with 4 and 6, behind them, glass-like; colour of head whitish or very light yellow, the hinder border narrowly black, a black cheek-stripe from each eye-curve up to vertex (this often coalescing with marginal black), another black stripe running down from vertex on each side of dorsal line (often coalescing with each other) to apex of false clypeus and splitting down its sides, the false clypeus itself often all black separated by a thin, light line from the last stripes and from the true clypeus, a dorsal, brown line on true clypeus. *Surface* of larva covered with extremely minute, black, pointed tubercles, each surmounted by a tiny colourless hair, except anal segment and segment 2; transverse, thin, impressed, parallel lines as already mentioned to each segment; segment 2 with a thin, black, chitinized, transverse collar. *Spiracles* oval, very light yellow, slightly raised, small; those of segments 12 and 2 twice the size. *Colour* glaucous whitish-green allowing the dorsal vessel (appearing darker green), the thin, white tracheæ and (in the male) the orange paired bodies near hinder margin of segment 10 to be seen through it; the anal segment not coloured in any way, segment 2 white; there are a few (2-4) minute disc-like tubercles of a light brown colour dorsally in the middle of segments 12 and 13—in a transverse row. L: 25 mm.; B: 3 mm.

Pupa.—More or less of the *shape* and *facies* of *Telicota bambusæ*; the colour is also similar but the main difference that strikes an observer is that, here, the cremaster is always directed slightly upwards; the *head* is square in front though slightly convex (frons) between the eyes which are prominent, the frons is in a plane at right angles to the longitudinal axis of the pupa, the vertex is inclined thereto at an angle of 30° ; segment 2 is parallel-sided, in the same plane as vertex of head and half its length; the thorax is only slightly humped, stout, evenly convex, highest a short way behind the front margin with evenly rounded shoulders which have a small, slightly roughened, low tubercular spot on their points, the front third of thorax sloping up at the same angle as segment 2 then gradually changing to the long hinder slope at a smaller angle in the opposite direction to the hinder margin which is a somewhat long, parabolic curve meeting the wings in a rounded, deep, rather broad angle of about 45° , this hinder margin flush with segment 4; segment 4 equal in length to segment 5, its hinder margin strongly curved backwards, that of segment 5 much less strongly curved; segment 6 one-third longer than 5; segments 11, 10, 9 telescoped into segments 10, 9, 8 respectively; segment 13 the same length as segment 12 which is half 11; segment 14 about the same length as 11 but falling away ventrally, the dorsal portion square bounded on each side by a strong, thin extensor-ridge ending in a point or short tooth-like spine at each hinder corner, the space between these ridges (dorsal area) rather deeply concave, the whole segment turned up at an angle of about 35°

to longitudinal axis of pupa; greatest breadth of pupa at shoulders; transverse section circular; proboscis produced free to end of segment 9 though pressed against surface; antennæ reaching two-thirds of the way towards ends of wings. *Surface* of pupa very nearly smooth, very shallowly and distantly pitted, covered sparsely with semi-decumbent, short, reddish, stiff hairs from, each, a minute, darker spot or dot; the cremaster fringed along the hinder margin and outside each tooth on each lateral margin by a dense fringe of reddish, porrect hairs as long as segment 11; a sparse tuft of hairs in front of each eye as well as on the back of each eye and a similarly thin tuft on convexity of frons—these hairs rather shorter than the cremastral ones. *Spiracles* of segment 2 have a very slightly raised, large semi-circular 'expansion' behind each, lying on the surface of thorax and shelving very gently forwards towards the angle made by the hinder margin of segment 2 and the antenna where the perforation of the spiracle is situated, this expansion being as broad as segment 2 is long; other spiracles small; oval, raised, light red-brown. *Colour* of pupa dull, light orange, darker on thorax and head; the cremaster shining red-brown. L: 16 mm.; B: 3.75 mm.

In the male of this Kanara species, above described and named *mæsoides* by Swinhoe (see *Lepidoptera Indica*, vol. x, p. 258), there is a short bar of whitish-grey, glandular hairs, all directed outwards along the vein, just above, or partly even on, vein 1 of the fore wing filling the interval between the basal yellow marking and the edge of the discal band-spot in interspace 1. This male mark was evidently overlooked in the original description of *mæsoides* by Butler and, subsequently by everybody else who examined the species—if it is that species at all. Certainly Swinhoe overlooked it, as there is no doubt at all that this and no other was examined by him. Both Watson and de Nicéville pronounced it long ago to be *dara* but did not even then notice the male-mark. The above is a very careful description of both sexes of the imago in case it should happen to be something new. In the meantime, on Swinhoe's authority above quoted, notwithstanding the presence of the male glandular mark, it will be called *mæsoides* and this, notwithstanding also, that it is not very exactly the same as the figures of that species in *Lepidoptera Indica*, vol. x, Pl. 815, Figs. 2, 2a, 2b. The larva and pupa there depicted, figure 2c, are those of the Kanara insect as far as they go, but as usual, they are bad reproductions of the originals.

Habits.—The egg is laid above or below the leaf of bamboo; the little larva eats the shell, gets to the tip of the leaf and makes a cell by joining the edges, often, at first, for quite a short distance, thus manufacturing a more or less cylindrical (long-conical) house; later on it eats the leaf above the cell, leaving only the midrib by which the cells hang; at the end, pupating inside this; often, though not always, it eats through the midrib so that the cell falls to the ground and pupation takes place there. This cell is strongly closed and thickly clothed or carpeted with silk inside; covered more or less by a white, cereous powdery excretion from the larva as is also the pupa which is not fastened by the tail inside. The cell fits closely to the pupa; its mouth is closed by silks spun across and across with a plentiful supply of white powder. The butterfly is plentiful in the Kanara District both above and below the Ghats, at actual sea level as well as in dense jungle on the hills. It is a very strong, fast flier, likes the shade with sunlight filtering through in patches; plays about for short periods

with others, always returning to a chosen seat on the top of a leaf, often near the ground, where it sits with its wings closed over the back. It is hardly ever met with in the open and is not an insect of the Plains. It goes to flowers sparingly but may be seen at bird-droppings, on leaves and, occasionally, at damp spots in the beds of nallas. It basks with the wings slightly opened. The food-plant of the larva is Bamboo of any sort: *Bambusa Oxytenanthera*, *Dendrocalamus*, *Teinostachyum*. Swinhoe gives the habitat as India, Burma, Andamans, Ceylon, Hongkong, Malay Peninsula. Under the distribution he says 'A common species throughout India. We have it also from Ceylon and the Andamans; Elwes records it from the Naga Hills, Sikkim and Perak; Watson from the Chin Hills; Doherty from Kumaon; de Rhé Philipe from Misuri and Fergusson from Travancore; J. J. Walker from Hongkong and Moore from Mergui.' He then adds a note stating that Doherty distinguished it from *dara* chiefly by 'the rich, dark, tawny-ochreous colour of the underside, *dara* being greenish-yellow set with dark scales. The markings are almost exactly alike. The prehensores are singularly different; seen from above, the uncus of *dara* is gradually acuminate while that of this is abruptly truncate and slightly bilobed. . . '

In these later days Mr. Ormiston has apparently discovered by renewed study of genitalia that *mæsoides* and *pseudomæsa* are different species—see his paper in the *Report of the Proceedings of the Third Entomological Meeting* held at Pusa, vol. ii, p. 1022. Until now *pseudomæsa* was considered to be synonymous with *mæsoides*. Up to 1898 *dara* included them both.

Swinhoe has treated the above species as a *Padraona*. The genus being distinguished from *Telicota* by very little as shown below, it is here treated as a *Telicota* because the larva is different from that of *Padraona gola* and the pupa is naked, the cell being open; the pupa is rather more like that of *Udaspes*, *Notocrypta* than like anything known to occur in the *Erynninae*. It is curious that Swinhoe states that the hind tibiæ in both *Telicota* and *Padraona* are naked. They are not, for they have fringes of hair. The palpi are likewise stated to be laxly scaled on the second joints while the scaling is actually rather dense. These conclusions being, in the present case, based upon an examination of *Telicota augias*, *bambusæ*, *mæsoides* and *Padraona gola*. Other species may possibly conform to Swinhoe's specifications.

Genus 26.—PADRAONA.

Imago.—These are all butterflies of the colouring and facies of *Telicota*. Also with similar habits; although, judging by *P. gola*, somewhat weaker in flight and more confined to the ground-vegetation in the hills. None are, probably, insects of the Plains. There are eight species occurring in British India (inclusive of *dara* and *mæsoides* following Swinhoe); these are *dara* (Kollar); *satra*, Swinhoe; *mæsoides* (Butler); *rectifascia* (El. and Edw.); *concinna* (El. and Edw.); *gola* (M.); *otala*, Swinhoe and *nala* (Plöts). The last two are from Burma and the N.W. Himalayas respectively; *concinna*, *satra* respectively from S. India and Ceylon; *rectifascia* from Sikkim, Burma; the habitat of *gola* is given below; of *dara* is the N.W. Himalayas, Sikkim, Assam and N. Burma and of *mæsoides* as stated above. Besides these there

are 24 species enumerated by Swinhoe as occurring in the Indo-Malayan and Chinese countries.

Antennæ.—As in *Telicota* but with the club moderate, not nearly so stout as in that genus, the terminal crook short, pointed.

Palpi.—As in *Telicota*.

Hind tibiæ.—As in *Telicota*.

Fore wing.—As in *Telicota* for vein 12 with the cell shorter, but not much; middle discocellular not so long in proportion to lower; vein 5 from their junction being much less curved from base; vein 3 closer to end of cell, 2 from the middle in both sexes; wing shorter, not nearly so produced apically; costa more arched, outer margin more convex, less oblique.

Hind wing.—Vein 7 a little before end of cell, less than in *Telicota*, the upper margin of cell not curving down at end; vein 2 emitted from close before base of 3.

Egg.—See below under *gola*.

Larva.—See below under *gola*.

Pupa.—See below under *gola*.

Habits.—See below. It is unfortunate that nothing appears to be known of the earlier stages of any species except *gola*.

219. *Padraona gola* (M.).—Male.—*Upperside*: very dark-brown with a slight purplish tint in certain lights; markings dark orange-ochreous, much darker than any of the *dara* group. Fore wing with a streak on basal half close to the costal line, filling interspace 12 completely but leaving bases of vein 12 and upper border of cell unmarked, orange-ochreous with the following orange-ochreous lines and spots:—a thick streak in cell along upper and bottom marginal veins developing into a spot at their ends, these spots however not actually coalescing, the streaks also not reaching base of wing, the lower streak rather undeveloped in the middle allowing the brown ground colour in middle of cell more or less to coalesce with that in base of interspace 2 and so with that in interspace 1; a discal broad, somewhat irregular band from middle of vein 1 to costa joined by a line from base along the upperside of 1 another line or band filling interspace between vein 1 and inner margin, the vein, however, remaining brown; the discal band consists of spots in each interspace 1 to 8; that in 1 occupying the middle third (really less than a third of the length of the whole interspace) with its outer edge excavated, its inner edge only very slightly so; the spot in interspace 2 somewhat longer, situated with its middle at junction of 2 with bottom of cell, its inner edge just before inner end of interspace, its outer end extending slightly further out than outer end of the one under it, also excavated; the spot in 3 occupies the base of the interspace and reaches just to its middle, slightly further than the outer edge of the next lower spot; that in 4 similar but slightly longer outwards, leaving the discocellular of cell brown and connected at its very outer, top corner by a thin chevron-mark with its angle outwards across interspace 5 with the lowest of the three subapical spots, striking it at its lower, outer corner; these three apical spots are quadrate, each excavated triangularly on outer edge, the lowest occupying the middle of interspace 6 and about half as long as the spot in 4; that in 7 dislocated inwards very slightly, about the same length, that in 8 dislocated outwards again and shorter; all this leaves the apex and a broad outer marginal area with the costal interspaces 9 and 10 connected with interspace 5 (except for the orange chevron-mark) all brown as well as a long base in interspace 1 and a short one in 2, the discocellulars and the veins crossing the discal band narrowly. Hind wing with a streak inside cell along the median vein covered by longish, decumbent hairs of the same orange colour; a broad, discal band consisting of quadrate spots in interspaces 1 to 4 and a small, triangular continuation into 5 from the inner, upper end of the uppermost, each spot quadrate, excavated on both sides, the band about as broad as the one on the fore wing and joined to a line along the lower side of vein 1 also covered with longish hairs. *Underside*. Nearly as dark as upperside, saturated richly (the expression is Swinhoe's) with orange-ochreous. Fore wing with the whole base of the wing including the bottom half of cell and everything inside the discal band below and including base of interspace 2, right out to outer margin below the band and beyond it in interspace 1 and the area beneath it and for half the distance to outer margin from the band in 2 and a slight continuation into 3 all black, together with a prominent mark along the discocellulars, the costa from vein 11—end to apex narrowly

and some blurred spots marking the outer borders of the spots of the band in interspaces 6 and 7 and still more slightly in 4. Hind wing is all orange except the whole of interspace 1a and the lower half of the next above, 1b, outside the discal band and even this is sprinkled with orange scales on the blackish ground; only the discocellular spot is pure-black, an elongated small lunule in the middle of the end of cell; the outer marginal area outside the discal band is marked with round blackish spots in interspaces 2 to 7 blurred owing to the plentiful sprinkling of orange scales; and each spot of the discal band is bordered inside obscurely but brownish; the band itself completely covered evenly with scales and, therefore, lighter and brighter than rest. *Cilia* on upperside of fore wing brown above vein 1 to apex, orange below it; of hind wing all orange; underneath, the fore wing ochreous brown above vein 1, but as on upperside below that vein; on hind wing orange. There are some brown, decumbent, longish hairs on upperside of hind wing in the upper half of cell besides the orange ones in lower half.—Female. Very likely the male in everything except that all the orange markings are smaller. Fore wing *upperside* with the base altogether unmarked except for the spots in the end of the cell—the ends of the streaks; the basal streaks and the costal one are only just indicated by a very sparse sprinkling of orange scales. The hind wing is absolutely like that of the male with the band narrower. *Underside* identical with that of the male except that the bands showing through are narrower beneath in the male and with the underside of club orange, both those being very pale-yellow in the female; palpi brown above, orange beneath in male. Antennæ black above, the shaft banded orange very pale in female; head and thorax above orange and brown mixed in male paler in female; abdomen above black with orange hairs in the male, but hardly any in female; below the colour of chest and abdomen is very pale in the male, nearly white in female; the legs with orange tibiae and tarsi in male, very pale in the female. Expanse 28 mm. or slightly more occasionally in the female, less for the male.

The above description of the butterflies is from bred specimens in the N. Kanara District of Bombay.

Larva.—The *shape* of the body is circular in transverse section with the ventrum somewhat flattened, thickest in the middle from segment 5 back, thinning to the neck and much more gradually back to the anal segment; the neck with a shining, somewhat swollen collar behind it to the short segment 2; the anal segment longly semicircular, convex transversely, somewhat sloping in dorsal line, overreaching the claspers in a thin flange and fringed all round with very fine, light hairs about 0.15 mm. in length—it is about half the width of the middle of the body; the *head* is slightly longer than a semicircle in shape, the vertex hardly emarginate in dorsal line, the face convex; the surface shining and rather coarsely reticulate-rugose, covered with extremely minute, light, erect, fine, 0.025 mm.—long hairs, these hairs over 0.05 mm. about the mouth-opening; the true clypeus longer than broad, about half the length of face or head, somewhat depressed in dorsal line, the apex very acute; the false clypeus a broad piece outside it of similar shape, reaching more than half way up the face, both plain green and smooth, shining like the head generally; the labrum is transverse, so thin as to be hardly perceptible, longer laterally than in the middle; the ligula also extremely transparent, about the same length as labrum or longer, triangularly emarginate in dorsal line of front margin which is whitish, otherwise colourless; the antennal, basal joint shining green, the 2nd somewhat rusty-soiled; the mandibles very large with the cutting edge entire, yellow with a terminal, dark red-brown edge; the eyes black, numbers 1-4 and 6 forming a semicircular curve, 1 and 2 very close together, 3 and 4 slightly further apart and 6 more than twice as far from 4 as 2 is from 3, number 5 far behind the curve forming a triangle with 4 and 6; the colour of the head is light grass-green with the dorsal line from the hinder margin over vertex to apex of false clypeus somewhat narrowly black, the line splitting there down each side of the false clypeus and forming a black, horse-shoe shaped curve on middle of face, the ends of legs reaching only half way to base of clypeus, also a broader, black band through eyes and up each side of head to meet on the vertex; segment 13 only about one-third the length of segment 12, decreasing in the length laterally on each side; prolegs short, green; true legs shining

green. Surface of the body dull, covered all over by minute, erect, very fine, white, 0.025 mm. long hairs which are about 0.075 mm. apart; each segment transversely folded with six or seven parallel folds from hinder margin forward for more than half its length; the segments quite distinctly marked. *Spiracles* very small, longly oval, nearly flush, white in colour, those of segments 2 and 12 very much larger. Colour is light glaucous-green, sprinkled plentifully with dark-green dots, especially in the spiracular region, the transverse folds often yellowish; a broad, more whitely-glaucous subdorsal band and a lateral, white line the whole length of the body, the actual dorsal line appearing dark-green (the dark-green dots, however, still darker green on it), a dark-green, interrupted line along the bases of prolegs; ventrum darkish-green. L: 16 mm.; B: 3.5 mm.

Pupa.—The type is that of the genera *Parnara*, *Udaspes*, formed naked on the underside of a leaf or blade of grass, the head more or less gently rounded in front outline, the eyes prominent, the frons inclined ventrally, the vertex produced out into a short, upturned, long, conical snout which is as long as the vertex-surface behind it which is again rather large, the hinder margin straight; segment 2 shorter than the head-vertex (without the snout), with both margins straight the dorsal line inclined to the longitudinal axis of pupa at an angle of somewhat over 30° and in the same plane as head-vertex and front slope of thorax; thorax only slightly humped, reaching its highest point just before hinder margin, this being also the highest part of pupa, the hinder margin a semicircular curve with a tendency to the triangular, meeting the wing-line in an open, rounded angle of about 60° , the dorsal line of pupa from thorax-apex backwards gradually sloping to end, the end generally slightly turned up in a light curve from segment 9 or 10; the lateral outline of pupa constricted very slightly at segment 2, the shoulders evenly rounded, the sides parallel from shoulders to segment 8, then gradually converging to the extremity of cremaster; segment 4 of equal length to segment 5, the hinder margin of former curved convexly backwards, that of 5 less so; segment 6 double the length of 5, the hinder margin straight, segments 7, 8 successively very slightly longer, segment 9 shorter again and coequal in length with segments 10, 11, 12; segment 13 a narrow, transverse ring, about one-third the length of segment 12; segment 14 rather longer than 12, deep at base, a thin plate behind, triangular in shape, the end rather broadly rounded, the lateral edges thickened and recurved dorsally leaving the dorsal centre hollowed out longitudinally; the proboscis reaching only just beyond the wings as a short point; the antennæ reaching only half way, pointed, the legs rather shorter, the coxal pieces large, nothing visible between the leg-limits. Surface somewhat shining, smooth, without hairs of any sort, segments well-marked. *Spiracles* of segment 2 oval, small, very slightly raised, hardly perceptible, the colour of the pupa; the rest similar, smaller, nearly the colour of the pupa. Colour grass-green with indistinct subdorsal and lateral, longitudinal lines the whole length of the body, an interrupted spiracular thread—the tracheæ seen through the integument; the head is lighter, the wings slightly also, the cremaster is whitish. L: 16 mm. over cremaster and snout, 14 mm. otherwise; B: 3.5 mm.

Habits.—The egg is laid on the tip of a grass-blade as a rule, and always single. The little larva makes a cell at the extreme point immediately on emergence, joining the edges of the blade with cross-silks. As it grows it increases the length of this; after reaching the third or fourth stages it lives on the undersides of the blades and just draws the edges loosely together, lying, thus, half exposed as often as not. The pupa is formed on the underside of a blade in such a cell generally formed expressly for the purpose. The larvæ are sluggish creatures but are rarely parasitized, and generally sit with segments 2-4 hunched up, the head lying back. The pupa is attached by the tail and a body-band, and rolls round and round when disturbed. The butterfly inhabits the undergrowth, and its habits are those of the genus *Telicota* for it is a strong flier. It does not, however, ever fly at any height from the ground;

the males bask on the uppersides of leaves with the wings half open, the lower wing more depressed than the upper as often as not; when completely at rest they close the wings completely over the back. The insects frequent flowers, generally such as grown low amongst grasses on the ground. The food-plants of the larva are grasses of various sorts, the softer ones being chosen for preference. Swinhoe gives as the habitat 'Sikkim Assam, S. India, Ceylon, Burma, Andamans, Malay Peninsula and Archipelago'. It is a common species in the Kanara District, both in rice-fields and in the jungles but, in that district, the rice-fields are never any distance from the jungles; indeed, they are generally surrounded by them. It is probably really an insect of jungles and heavy rainfall.

Genus 27.—HALPE.

Imago.—Insects of medium size, brown above and below, the fore wing with three small, subapical, semihyaline spots, generally two more at the end of the cell on the discocellulars, sometimes only one; and a discal series in interspaces 2, 3 and 1, the last occasionally wanting. The upperside of hind wing is immaculate in nearly all species. On the underside the hind wing may have a white or whitish or yellow band with or without darkish spots; or may have a row of darkish or even white (rarely) spots or in a few cases may be washed with yellow with or without dark spots; in these yellow-washed species there is some yellow on the upperside of hind wing too and, occasionally, on the upperside of fore wing. *Halpe* is a large genus, containing twenty-four species in British India and twenty-six more Indo-Malayan, Japanese and Chinese representatives. Swinhoe has divided the genus into two, according as to whether there is a sex-mark in the male or not; calling those that have none *Thoressa* and retaining *Halpe* for the remainder. There are three species of *Thoressa* in British India and one Indo-Malayan. The former are named *astigmata* (Swinhoe); *masoni* (M.) and *honorei*, de Nicéville. The first and last occur in S. India, the other is only found in Burma and Tonkin. In the genus *Halpe* proper two species *moorei*, Watson and *hyrtacus*, de Nicéville are found in the Bombay Presidency and their transformations are given below together with those of *astigmata* and *honorei*, also occurring there. The names of others of the genus in British India are *ceylonica*, M., from S. India, Ceylon, Mergui; *homolea* (Hewits.) from Sikkim through Burma to Java and Sumatra; *sikkima*, M., from Sikkim to Burma; *gupta*, de Nicéville from N.W. Himalayas to Assam and China; *burmana*, Swinh., from Burma; *knyvetti*, El. and Edw., from Sikkim; *zema* (Hewits.) occurring in Sikkim to Burma and in Borneo; *cerata* (Hewits.) in Sikkim, Assam, Burma; *debilis*, El. and Edw., in the Khasia Hills; *fusca*, Elwes in Upper Burma; *decorata*, M., in Ceylon; *sitala*, de Nicéville in the Nilgiris, Anamalais, Ceylon; *brunnea*, M., in Ceylon; *kumara*, de Nicéville in Sikkim, Bhutan; *aina*, de Nicéville in N.W. Himalayas, Sikkim, Assam; *hyrie*, de Nicéville in the Naga Hills, Sikkim, Assam; *evershedii*, Evans in S. India; *albipectus*, de Nicéville in Burma.

Antennæ.—A little more than half the length of the costa of fore wing; club moderate, long with a sharp crook, its tip short, pointed.

Palpi.—Porrect; second joint laxly scaled; third joint minute, obtusely conical.

Hind tibiæ.—Slightly fringed and with two pairs of spurs.

Fore wing.—The male in all except the *Thoressa* section with a linear, discal stigma on the disc running obliquely from the lower end of cell to below vein 1 before its middle; see also below for the hind wing. Vein 12 reaches costa before upper end of cell which (cell) is about two-thirds the length of costa; 6 from upper end of cell, so close that there is no upper discocellular vein; 5 nearer 4 than to 6 because the middle discocellular is nearly twice the length of the lower; the discocellulars in an inwardly-oblique, straight line; 3 from about one-sixth before lower end of cell in the male, in the female its base a little nearer end; 2 beyond the middle; costa arched; apex subacute; outer margin evenly convex, shorter than hinder margin. Hind wing in all except *Thoressa* with a tuft of hair on upperside attached along

costal vein; vein 7 from well before upper end of cell, bent upwards at its base in male with the upper margin of cell bent inwards and downwards to cell-end; in female 7 and upper margin of cell are both nearly in a straight line; discocellulars faint; 5 absent; 3 from quite close to lower end; 2 from about an eighth to one-fifth before end; costa and outer margin evenly curved.

Larva.—More or less of ordinary shape with the anal end rather broadly rounded, sloping to end; the head rather small for larva but of course much broader, higher than the neck, generally yellow but sometimes much suffused with black, rugose as to surface, nearly quite round. *Surface* dullish, the skin thin showing the tracheal tubes through; covered with minute, erect hairs, those round the dorsoventral margin longer; those on anal segment besides, from small, prominent tubercles. *Colour* generally watery green; no longitudinal lines except in *Thoressa honorei* where there is lateral, longitudinal band above which the dorsum is yellowish and underneath which the colour is white, the anal segment being black.

Pupa.—More or less of the same shape as that of *Telicota bambusæ* with a slightly convex frons to head and a well-developed, strong cremaster with very distinct extensor-ridges, the suspensory shafts hooked in some species, simple in others, resembling hairs; the spiracle of segment 2 with a very large, ear-shaped or funnel-like expansion behind it. *Surface* covered with fine hairs, not easily visible however except under a lens, those on eyes generally longest—the pupa of *Thoressa honorei* has the hairs so minute as to be difficult to see even with a lens.

Habits.—The butterflies are strong fliers, settling to feed upon flowers but only for a short time, very quick in settling and rising; resting with the wings closed over the back, the front ones rather much drawn down between the hinder; they are fond of shade but fly mostly round the tops of high trees and may occasionally be seen suckling at bird-droppings on leaf-surfaces. They are all perhaps confined to jungle country. The eggs are laid on the underside of leaves as a rule; the larvæ make a cell by joining the edges of the leaf at the point at first and generally through their lives but sometimes double a leaf transversely; they finally close it strongly at both ends as well and cut it free so that it falls to the ground. All the pupæ have a covering of cereous, white powder which is excreted by the larva and gets all over the inside of the cell. Most of the larvæ occasionally lie over in the cell for several months in the dry season; then pupate and the butterfly comes out in the normal ten days or so. The food plant of all the caterpillars known is Bamboo of any kind.

220. *Halpe moorei*, Watson.—Male. *Upperside*: blackish-brown when fresh with a purplish gloss or bloom in certain lights; fading paler. Fore wing with seven hyaline-white spots: three small, subapical in interspaces 6, 7, 8, the first two near the bases of the interspaces, the uppermost about the middle; two spots in the cell, one above the other in a line with the sex-mark; two larger, sub-quadrate in interspaces 2, 3, the lower the larger slightly, the upper with its inner edge in a line with outer edge of the other, both a bit outwards from bases of their interspaces; the costal space up to half way out, interspace 1 from base to the stigma and the inner-marginal interspace half way out covered with golden-yellowish, decumbent hairs; the stigma itself black, swollen. Hind wing unmarked except for an obscure paleness about the inner half of the interspaces 4 and 5 which may spread downwards into interspaces 2 and 3 slightly and upwards into 6; the lower half of cell and this paleness with decumbent, longish, pale-yellow hairs; the upper half of cell with a tuft of blackish hairs producing a distinct blackish appearance; a ridge of longer, only semidecumbent hairs along vein 1b from base to outer margin. *Underside* paler; brown. Fore wing with the semihyaline spots showing through; a sprinkling of pale-yellow scales along the costa above vein 12 and to apex and thence somewhat narrowly down to vein 2 along

outer margin with small, blurred-whitish submarginal spots in interspaces 2 to 7, one in the middle of each. Hind wing with a conspicuous white band along and including vein 1b expanding generally downwards at end more than half way to anal angle and from this arises, just about half way towards the outer margin, a broad, conspicuous, white band that runs across the whole wing to the outer margin just below the end of vein 8; this band being made up of a large, quadrate, white spot in interspace 1b about 1.5 mm. long, the inner edge straight and immediately below junction of vein 2 with the cell, its outer edge slightly emarginate; followed by a much narrower one in the base of interspace 2, its outer edge in a line with outer edge of the other; a third nearly filling the whole of the basal half of interspaces 4, 5; a fourth above, more or less oval, little more than half the size of the third (which is the largest of all), its inner edge in a line with outer edge of third in interspace 6 and, finally, a small, blurred one in interspace 8 situated just about over the middle of the fourth; every spot divided from the others by the brown veins—the very extreme base of interspace 3 may also have a few white scales in it; the base, costa and along outer margin have a sprinkling of pale-yellow scales and there is just an indication of a small, white spot, submarginally in interspaces 2, 3 and 4 with, always, a larger one amounting to a little patch in interspace 1b that may be joined by a line with the end of the white streak along vein 1b; a few light-yellow, decumbent, long hairs in the cell. *Cilia* of fore wing above and below brown with tiny white intervals in middle of each interspace, the interspace 1 pure white; hind wing with them checkered white and brown equally and with the bases brown. There is very little sign of a fringe of hair along inner margin of fore wing. Antennæ black, the shaft with the edges of joints thinly white below and the club white below from very base to half way towards tip and up along the sides of the dull-orange upper half; palpi brown on top, pale ochreous below and set with black, short, bristle-like hairs, the third joint all brown surrounded completely at base by pale-ochreous; head brown mixed with grey; thorax brown with yellow and greyish hairs and a greenish sheen; abdomen brown with grey segmental bands at edges of segments; beneath, the chest pale-ochreous with great jet-black patches on the coxae, the tibiae and tarsi light-orange; abdomen below also brown but the segmental bands very broadly whitish-ochreous.—Female like the male but with an extra white spot of small size in the middle of and just above vein 1 of fore wing; the outer edges of the large discal pair of spots always excavated. Expanse 30-35 mm.

Larva.—The body is of the shape of that of *Telicota bambusæ* or any of the genus *Halpe*; the head about as broad as segment 3; segment 13 convex backwards, becoming shorter towards the sides, not quite half as long as segment 12 and rather higher at its posterior margin than the front surface of anal segment; this latter about as long as segment 12, rounded at end broadly—an elongated semicircle in shape; a distinct neck, the prolegs and legs short; the body flattest in the middle but very much the same breadth really from segment 3 to segment 12; the head rather higher than broad, otherwise rounded, thick, the surface finely honeycombed—roughened with a clothing of extremely short, fine, light, erect hairs, longer about the mouth-opening and eye-regions; the clypeus about one-third the height of the face, longly triangular, the apex acute; the false clypeus outside it forming a narrow band all round, also triangular, about half the height of face or less, a slight raised dorsal ridge up the middle or the former; the labrum transverse, narrow, parallel-sided, brown-red in colour; the ligula pear-shaped the narrow side forming the base or stalk, also red-brown, shining as is the labrum; the antennal, basal as well as second joints light red-brown, shining; the mandibles also rusty with narrow black borders and without teeth; the eyes, four in a very gentle curve, equally spaced, the fifth well behind forming an equilateral triangle with sixth and fourth, also red-brownish. The surface dull, covered with extremely minute, erect, fine, light hairs which are much longer subspiracularly and still longer again on the anal margin and dorsum of the anal segment where each hair rises from a prominent, though still small (prominent under a strong glass), conical, rusty tubercle—the dorsum of 14 covered with these tubercles; the hinder margins of segments have the front margins of the succeeding segments telescoped into them and the surface of the segment just before the hinder margin is crossed by transverse, depressed lines forming four folds or interspaces parallel to each other and to the

hinder and front margins; some folding in spiracular region; *Spiracles* small, convex, broadly oval, very light yellow-brown or brownish-yellow, those of segment 2 much larger, those of segment 12 nearly twice as large but smaller than those of segment 2. *Colour* greenish with a slight rusty tinge especially dorsally; the male organs on dorsum of segment 10 very conspicuous, orange; the tracheæ slightly visible subcutaneously; the colour of the head is soiled very light orange with a broad, dorsal, dark shade from hinder margin over vertex to apex of clypeus, this shade being sometimes a broad, brown-red band which splits thinly down each side of the clypeus; the eyes are dark, the four in a curve placed on a light red-brown patch that is not extensive. L: 25 mm.; B: 3.5 mm. (not quite full-fed).

Pupa.—The pupa, too, is much the same shape as that of *Telicota bambusæ* in every way: it has the same sort of spiracular expansions to segment 2; the colour is also the same; the body is widest at the thorax; there is a slight, light lateral constriction at segment 2 behind the rather prominent eyes, the vertex of head is long and inclined at an angle of somewhere near 45° to the longitudinal axis of the pupa, the frons is perpendicular to that axis, rounded, short and prominent though only lightly so, convex; segment 2 is in the same plane as the vertex of head, short, only about one-third the length of the vertex of head and the margins are parallel, straight and quite flush with dorsal surface; the front slope of thorax is between 45° and 60° to the longitudinal axis of the pupa, short, the general surface of thorax even, lowly convex, the dorsal portion after the short frontal slope gradually ascending to the hinder margin, the whole dorsal line of pupa from about the front third of thorax to segment 9 being parallel to the longitudinal axis; the hinder margin of thorax is more or less a low, broad triangle, the apex of which, however, is narrowly rounded and it meets the wing-line on each side in an angle of between 60° and 90° which angle is rounded, broad and shallow, segment 2 is about one-fourth as long as thorax, the hinder margin curved convexly backwards and is equal in length to segment 5 whose hinder margin is also curved backwards but the curve is not nearly so convex (backwards); segment 6 is a good deal longer than 5 and equal to the succeeding segments up to 12; segment 13 is very short and the dorsal slope, with that of segments 11 and part of 10 and segment 14, is inclined at an angle of about 30° or less to the longitudinal axis of pupa; segment 14 is about equal in length to segment 12 but is, of course, narrower at the front margin, still narrower at the hinder margin—it is trapeze-shaped—thinning from the deep base at segment 13 backwards to a thin edge at extremity where the corners are produced out sideways into short, conical teeth, one to each, the edge between these teeth being concave forwards, the dorsal surface also has a fairly deep, gradual depression in the dorsal line and there is no sign of extensor ridges either dorsally or ventrally; the transverse section of the body is more or less circular from thorax to segment 13; the proboscis is not produced in any way beyond the wings and the antennæ reach to about two-thirds the length of wings from their bases. *Surface* of pupa is slightly shining and nearly smooth though somewhat granulated-looking, the segment margins are all distinct, those of segments 8-12 very slightly raised (the hinder margins), all of them have a narrow, quite smooth band along hinder margin; the whole surface covered with comparatively long, soft, decumbent, light hairs; longer, erect, golden reddish hairs which cover, in a tufty way, the sides of eyes and the frontal boss and the softer hairs are rather longer than usual and more erect on the front slope of thorax dorsally; the cremastral edge is set with the longest, golden-reddish hairs of any, sticking out backwards, straight along the sides as well as between the teeth and the hairs generally on segment 14 are longer than elsewhere. *Spiracles* of segment 2 are distinguished by large expansions which are semicircular, only very slightly raised from the general surface of segment 3 upon which they lie, straight edge being along the common margin of segments 2 and 3, the length slightly longer (the length being taken in the sense transverse to pupal length in this case) than segment 2 in dorsal line, the surface of the semicircle sloping down from the raised edges forwards to a deep slightly excentric depression or hole placed nearer the outside (next the antenna) edge of the semicircle than to the inside; this depression or hole is transparent-looking, light yellowish-green while the expansion semicircle containing it is dark red-brown so that it is very conspicuous and, being placed

up against the front margin of segment 3, that is against the diameter of the semi-circle, produces the appearance of a low or short horse-shoe of which the upper leg (the one nearest dorsal line of pupa) is thicker than the lower, the curve of the horse-shoe being thinnest as the hole is comparatively rather large; the two ends of the horse-shoe where they impinge on the segment margin are somewhat thickened also; the other spiracles are small, oval, slightly raised, rather light red-brown or yellowish; above each spiracle or in a corresponding position, dorsolateral therefore, is, on each segment, much nearer the hinder margin than the front margin, a small, oval, flat, shining tubercle which is red-brown in colour also. *Colour* of pupa that of a fresh bone, the wings, thorax, segment 2 and head somewhat translucent or watery looking; the cremaster shining, bright brown-reddish, at least at the extremity; later on these parts become darker; there is always a greenish, dorsal, subcutaneous line. L: 17 mm.; B: 4 mm.

Habits.—The young egg larva makes a cell at the point of a leaf by joining the edges, and this method is continued up to the end by joining longer lengths of edges, a new leaf being chosen whenever necessary; it feeds at first upon the leaf just in front of the cell entrance—which is of course always towards the base of the leaf—but later on, when grown larger, from the second stage onwards, perhaps, goes out to feed on other leaves close by. The cell is always fairly tight for the body of the larva, but is invariably of much greater length and circular in section; it is invariably lined inside thickly with silk, sometimes evenly, sometimes in steps. When about to pupate, the larva makes the cell some way away from the tip of the leaf—that is, the tip is not included—by eating a line from edge inwards on one side to midrib and another about 30 mm. further away, towards the leaf-base, also in from the edge to mid-rib and turns over the piece thus freed at both ends on to the opposite upper surface of leaf, fastens the edge down and, after shaping the two ends irregularly, also fastens them down; clothes the inside densely or thickly with silk, evenly and in steps, settles down and, after two or three days, changes into a pupa; this pupa is often lightly covered with a cereous excretion which is visible between the two front pairs (sometimes between all) of prolegs on the sides of the larva prior to settling down. The butterfly is a very strong flier and rises often high into the air; is not much seen except at the tops of hills where the trees are low and accessible (their crowns). There males come and bask on the tops of leaves, sitting with their wings closed over the back, motionless, the antennæ directed well out forwards, apparently in an attitude of expectancy. They dart at anything of their own species that comes along, and go off into the air in pursuit. They, however, nearly invariably return to the same perch time after time. Here at the hill-tops, they make their appearance about 9 a.m. and stay until about 11 when they apparently go somewhere else, may be to feed. They do not go much to flowers but may be seen sucking at bird-droppings on leaves occasionally. Their visits to hill-tops must be made in quest of females; out of a couple of dozen butterflies caught in some successive days, not a single one was female. The latter sex apparently does not bask and is not easily got. In breeding however as many males as females emerge so there is no real discrepancy in numbers. The little egg-larva emerges through the top

of the egg, leaving a round lid and sometimes eats the shell but sometimes does not; the egg is laid on the underside of a leaf; the little larva is honey-coloured, very pale and has a round, quite black head. The final cell is cut free from the plant or rest of the leaf (when the leaves are too large to be wholly employed as in *Ochlandra talboti*, Brandis) and falls to the ground where pupation takes place within it. The butterfly is an insect of jungles and, probably, fairly heavy to very heavy rainfall, and is plentiful in the Kanara District from the sea-shore up to 3,000 feet, the highest point of the Ghats within those limits.

In one description of the pupa made, it is noticed that there is 'a small, flat, shining, oval disc-like tubercle laterally on segments 2 to 12 as well as three under spiracles of segments 10, 12, two on 8, 9, 11, dark-brown in colour' and further that the cremaster is 'trapeze-shaped, short, broader than long, about as long as 13, its distal extremity the shorter, the lateral sides shortest with a dorsolateral tooth or short point directed out and upwards at end of each or immediately before the end; the dorsal extensor-ridges very short, the ventral ones nearly wanting altogether'. The head is sometimes much more suffused and, often, has a crescent-shaped blackish mark in the middle of face of each lobe, sometimes connected to the dorsal band at apex of clypeus.

Swinhoe gives the habitat of *Halpe moorei* as India, Burma, Ceylon, Hongkong. Under 'Distribution' he says, 'Watson records it from the Chin Hills, Rangoon, Berhampur and the Andamans; de Rhé-Philipe from Misuri; J. J. Walker from Hongkong; de Nicéville from Sikkim; Elwes from Trichinopoly in Ceylon and Tavoy in Burma; Moore from Calcutta; we have received many examples from the Khasia Hills and have it from Port Blair in the Andamans, Calcutta, Ranikhot, Karwar in N. Kanara; Davidson, Bell and Aitken bred it at Karwar; our figures of the larva and pupa are from Davidson's original drawing not previously published. Adamson says it is a common species in Burma.'

The figures above referred to are contained on Plate 817 of *Lepidoptera Indica*, Fig. 1c. The butterflies are also figured, 1 being the male upperside, 1a the female and 1b the underside—the stigma on the male is not indicated at all and the colour is not nearly black enough. Neither the representations of the butterfly or the larva and pupa are good.

221. *Halpe hyrtacus*, de Nicéville.—Male. Upperside dark-brown, the area above vein 12 of fore wing along costa sparsely sprinkled with yellow scales and the basal half of cell sparsely set with golden, decumbent hairs; there are a few scales towards apex on the costa. Fore wing with six very small, white, semihyaline spots, or seven; two in the cell, the upper against the upper margin at the origin of vein 9, the larger as a rule, the other in the very outer, lower corner, much smaller and often absent; three subapical, occasionally very tiny dots in interspaces 6, 7, 8, that in 6 near the base of the interspace, that in 7 moved very slightly inwards, that in 8 about the middle of its interspace and the uppermost may be absent in some specimens; a spot, the largest—but still small—in the very base of interspace 2 with another, very slightly smaller and a bit further out in 3. The brand on the disc dark, consisting of three parts, the uppermost circular in the very base of interspace 2, the other two crescent-shaped, the hollow inwards, the two

in a line filling the interval between veins 1 and 2 from the end of cell to the middle of vein 1 or slightly further out; this band with each section tumid, the crescent-shaped parts naked and shining. There is a very slight hair-fringe along inner margin. Hind wing quite immaculate with darker, short hair occupying top half of cell, longer, decumbent, yellowish hairs in the bottom half stretching outwards to about three-quarters the way to outer margin below vein 3 followed by brown hair in interspace 1b and dense, long, brown hairs in a thick ridge all along 1b to anal angle. *Underside*: much paler than upper side. Fore wing with the spots as on upperside and the whole base below the cell and out to a line joining the upper end of the stigma (therefore lower, outer end of cell) to the tornal angle pure-white or nearly so; pale, blurred, whitish spots in interspaces 4, 5, 6 and 7 submarginal with obscure darkish lunules bordering their outer edges. Hind wing with the whole base except above vein 8 and base of interspace 7 white out to the middle, extending a little more outwards in interspace 1a and the lower half of 1b, white with a spot of ground colour under the junction of vein 2 with cell, another at the junction of 6 with cell and a third under the middle of vein 8; some yellow scales on the brown, costal area above vein 8; some few white scales making obscure spots, postdiscal, in interspaces 3, 4, 5; there are some white, decumbent hairs in the cell from base to its end.—Female like the male in everything except that the semi-hyaline spots are always larger, the stigma is absent on the fore wing upperside; and the whole colour is paler. *Cilia* dark-brown at base and very little paler beyond. Antennæ with the top bent sharply at right angles and rather long, longer than the width of club; black above with the tip of club grey; beneath, the upper part of shaft and club are yellow, the apiculus dark brown-red. Palpi above black and yellow mixed, the third joint brown and somewhat porrect, not hidden by the hairs; below light-ochreous. Head, thorax and abdomen more or less concoloured with wings above; below, the pectus pale-ochreous in middle, whitish at sides, the abdomen grey with last segment brown; legs slightly ferruginous. Expanse 40 mm.; the female slightly larger than the male.

Larva.—Is of a more or less ordinary skipper *shape*: more or less cylindrical, the ventrum of course, however, depressed or flattened; the prolegs and true legs short; the anal segment sloping at about 60° to the longitudinal axis of the body, semicircular in shape, convex transversely as well as very slightly longitudinally, the whole surface of anal segment except the extreme anterior margin covered with small, star-shaped, black or deep red-brown tubercles from each of which rises a soft, short, light greyish hair: the ground-colour on which these tubercles are situated is greyish, different from the rest of the larva; the neck is slightly tumid owing to the presence of a somewhat thickened, light-yellowish collar from spiracle to spiracle; the *head* is large, somewhat oval in shape, higher than broad, the medial (dorsal) line very slightly depressed on vertex, continued down as far as apex of clypeus, the lower face, from just about the level of the true clypeus down, being slightly flattened and inclined at an angle to the upper part; the true clypeus reaches about one-third or slightly more up the face; the false clypeus about half way and both are triangular in shape with the apex acute, the surface somewhat lugose; the labrum is small and membranous; the ligula rather large, deeply emarginate and trapeze-shaped, the outer or emarginate side being the longest; the whole shining red-brown; the mandibles are large, shining, dark red-brown; the eyes are black, four in a short curve, the other two difficult to see; the surface is rugose-cellular, rather strongly so, and is covered with a clothing of erect, fine, soft, greyish hairs that are best seen against the light sideways: they are not short,—needless to say the head is much broader and higher than segment 2; the colour is dark brown or nearly black, somewhat shining, this black being confined to the whole of the hinder part of the head, thence over vertex in a broad dorsal band down to include the clypeus and the lower parts of face leaving the middle of each lobe-face soiled lightish yellow with a triangular point of brownish—a sort of flush from the black—coming up into its basal part parallel to the dorsal line for a short distance; there is a similar point coming out from the broad, dorsal line about half way between apex of clypeus and vertex which is directed diagonally out and upwards. *Surface* of larva dull, more or less smooth except for some very short, erect, minute, light hairs which are sparsely disposed over the dorsal parts above the spiracular line; there are some much longer hairs in the

dorso-ventral region and along anal posterior margin as well as ventrally, these longer hairs like those on the head: soft, light, more or less erect; each segment has a series of depressed, thin lines, parallel to hinder margin and nearer to it than to front margin: there are five of these, the first two much further apart than the others; then there are the small, hair-bearing tubercles on segment 14 mentioned above; segment 13 is very short. *Spiracles* not very small, light yellowish, rather longly oval, nearly flush; those of segment 2 very much larger, those of segment 12 not so large as those of segment 2 but considerably larger than rest. *Colour* of the larva is a very soiled-looking somewhat watery olive-green, with the ventrum green and the anal segment dorsally greyish; segment 2 yellowish. L: over 25 mm.; B: 3.5 mm. when not quite full-fed.

Pupa.—A stout pupa of the shape of *Telicota bambusæ*; thorax stout and the head transverse, bowed, the frons high, in a plane at right angles to longitudinal axis of the pupa, slightly convex, the vertex at an angle of 45° or more, broad between the bases of antennæ, the hinder margin straight, the eyes large, the outline in front of head square, slightly convexly rounded, the labrum a small, trapezoidal piece between the bases of proboscis, the ligula a small, very acute drop-shaped piece contiguous to it in front: this labrum and ligula ventral of course; segment 2 a short piece in the same dorsal plane as vertex of head, its hinder margin straight, laterally cut away to receive the spiracles and there very thinly raised over the opening of spiracle; thorax at apex the highest part of pupa, the front slope for half its length in the same plane as segment 2 and head-vertex, then curving round gradually to become more or less parallel to the longitudinal axis of pupa, then slightly inclined towards segment 4 which rises towards it and forms its even hinder slope-end, the hinder margin a parabolic curve rather narrowed in the pole, that meets the wings in a widely open, rounded, rather deep angle of 45°; segment 4 is as long as segment 5 in the dorsal line; segment 6 is one-third longer than either; segment 12 is shorter than 11; segment 13 is just represented as a base to the cremastral segment, very short; bevelled margins of segments 8-11 hardly noticeably developed; cremastral segment longly triangular though truncated at extremity, the dorsal extensor ridges thin, distinctly raised, the area between them nearly flat, the sides of cremaster (lateral faces) also triangular with a large, oval, shallow hollow bordered by a raised line or thin ridge at the ventral base under segment 13 or on the outer face of the anal claspers (represented ventrally by the scars) and *this large hollow or oval is characteristic*, the ventral extremity of cremaster hollowed out considerably, the suspensory hooklets small, bunched at the very extremity and slightly dorsally; the proboscis is not produced at all beyond wings; the body is of equal width from the evenly rounded shoulders to segment 6, the pupa is constricted slightly just behind eyes, the head nearly as broad as shoulders, the body from segment 6 thinning to cremaster gradually, the dorsal line thence to cremaster slightly convex, the ventral line straight. *Surface* of pupa moderately dull, transversely finely rayed with minutely waved, raised lines, the segment margins well enough marked, the extreme hinder margins of each segment narrowly smooth on segments 8 to 10; the whole pupa covered with a soft, more or less decumbent, short, light down which is longer and more erected in the spiracular regions and much longer and nearly erect on the head-frons, the front and posterior surfaces of eyes and along the sides of the cremaster; there is, besides, a very small, flattened, red-brown, chitinized disc subdorsally near the hinder margin of each segment and a subdorsal, depressed point on segment 2. *Spiracles* of segment 2 with an ear-shaped, deep-brown, rugose expansion on segment 3 front margin consisting of a slightly raised semicircular boundary behind, the whole surface sloping downwards from that boundary to the front margin of the segment and ending in a hole in its spiracular region under the thinly raised hinder margin of segment 2 alluded to above: this dark brown, rugose surface quite 2 mm. in breadth by over 1 mm. in length (dimensions corresponding with length and breadth of pupa); other spiracles rather small, narrowly oblong, raised, yellowish. *Colour* of the pupa that of a soiled fresh bone, the wings greenish, the cremastral extensor ridges and borders of the hollow red-brown; boss of head or frons and dorsal portion of thorax sullied with brown, the latter having it disposed in subdorsal bands; abdominal segment with two, parallel, transverse rows of brown dots reaching down to

lateral line and a postspiracular and some subspiracular ones. L: 20 mm.; B: 5 mm.

Habits.—This species was first caught amongst the hills in big jungle at a place in the N. Kanara District of the Bombay Presidency called Tarimallapur, an uninhabited village-site up the Kalinaddi river, forty miles from the coast. Shortly afterwards a larva was obtained at the same place, in the year 1895, on the large-leaved bamboo, *Ochlandra talboti*, Brandis, that grows in damp, even swampy places by nallas in the valleys. Many years afterwards specimens were sent to de Nicéville in Calcutta who said that he had just named it from two ragged specimens received from Travancore. Subsequently to the first discovery, a few years afterwards, more larvæ were obtained at another place on the crest of the Western Ghats in the same district, called Anshi; and on the same species of bamboo; and, since then too, it has been occasionally found at this last place when looking for caterpillars of *Halpe astigmata*; at least it would be more correct to say that, when looking for caterpillars of *hyrtacus*, those of *astigmata* were found and collected thinking they were the other. It is a rare species and, like *astigmata*, is hardly ever seen in the perfect state, flying. The butterfly has been seen thrice and *astigmata* once. Many caterpillars have occasionally been obtained when a violent effort was made; but the habit they have of lying over for months during the cold weather has prevented breeding any number of perfect insects. The best time for collecting is during the very heavy rains in August when it is difficult to get about. No carts can be had, the river is in flood, roads are impassable because unbridged; the bungalows are very damp and uncomfortable. Besides which the place where the larvæ have to be sought is swarming with leeches. The habits of the larva are those of *Telicota bambusæ* as far as the cell-making is concerned; it is just as sluggish or even more so in its movements. It does not appear to excrete any cereous, white powder in the cell but probably cuts the cell away from the leaf when about to pupate. The pupa is firmly attached by the tail inside. The habitat is the Western Ghats from Kanara to Travancore.

Another note about the larva is to the effect that the 'anal segment is covered with minute, star-shaped, low tubercles of a red-brown colour from each of which springs a short, erect hair'.

222. *Halpe (Thoressa) astigmata* (Swinh.):—Male. *Upperside* blackish-brown. Fore wing with the basal portion as far out as to half the length of cell and the whole of the inner-marginal space below vein 1 to outer margin set with golden-yellow hairs, sparsely except along the inner margin; and with six semihyaline spots, two subapical towards the bases of interspaces 6, 7, the lower slightly the larger and two discal, the upper towards the base of interspace 3, somewhat larger still, the fourth underneath, further in, just before the junction of vein 3 with end of cell, in interspace 2 largest, oblong but only slightly longer than broad, its edges straight, just 1 mm. broad; and two in the cell, coalescing in a point on the middle line of cell and situate so that the lower border of the lower one takes up a length of the bottom vein of cell extending from close to the outer, lower corner of cell as far in as the base of vein 2 where it joins the bottom vein, both these cell-spots about equal in size, smaller than the discal one in interspace 2 but larger than any of the others. There is only a slight fringe of hairs along the inner

margin. Hind wing: the lower half of cell extending outwards thence half way to outer margin covered sparsely with golden-yellow, decumbent, longish hairs; much denser, brown, longer hairs all along vein 1b from base to outer margin; but without markings. *Underside* paler than above because it is sprinkled all over sparsely by small, yellow scales all over the hind wing and along costa and the apex down to tornal angle in a narrowing band. Fore wing with spots as above and a very indistinct series of pale submarginal spots in interspaces 3 to 7; interspaces 4, 5 often pale at base and 1 and inner-marginal space pale at outer ends. Hind wing with a pure-white spot of scales in the middle of interspace 7 and at its base, the latter the larger (but rather small at that) and one beyond the middle of interspace 6 followed by very small ones in 5, 4, 3 somewhat further out and in a straight line with finally one in the very middle of 2; a pale streak along the upperside of vein 1b from base to half way out with a pale patch in interspace 1b before the margin. There are some shortish, yellow, decumbent hairs in cell and outwards below it and extending a bit beyond its end. Antennæ black above, the shaft black below too; the club with just the end ochreous as well as the whole of the underside, the apiculus brownish-red and this colour extending some way down the underside of club. Palpi brown above with some ochreous hairs, dull ochreous below because of a mixture of black hairs. Head brown; the brown with a greenish sheen; abdomen dark-brown with some yellow scales. Below, the pectus ochreous, the legs ferruginous, all pale; abdomen dark but with broad, pale-ochreous segmental bands.—Female like the male but somewhat paler; the spots no larger. *Cilia* on fore wing ashy-brown slightly paler at apex and more extensively above tornal angle; on hind wing all very pale throughout. Expanse 35 mm. for female; the male a bit smaller.

Larva.—Is of the ordinary shape, fattest in the middle, fining gently to both ends; the anal end rather longly semicircular, dorsally sloping in segment 14 at about 30°, the extremity slightly overreaching the anal claspers; segment 13 rather less than half 12 in length; 14 about as long as 12, segment 11 somewhat longer than 12; legs short; head rather small, broadly elliptical with the vertex broadly rounded and very slightly indented in dorsal line and depressed as far as the top of clypeus, the bottom of depression angular; the surface cellular-rough, the roughness somewhat accentuated and prominent on vertex, clothed besides with semi-decumbent, short, somewhat sparse, light hairs all over, these hairs somewhat longer about base of lobes and mouth-opening; true clypeus triangular, much longer than broad, apex very acute, reaching about two-fifths the way up face, with irregular ridges down its surface; the false clypeus reaching about half-way up face, narrow beyond the true clypeus, the strip on each side increasing in width from base to apex but fairly narrow even at apex, triangular with the apex acute; the labrum dark-brown like the head, transverse, rather short; the antennal joints both shining, whitish-soiled; the mandibles large and strong with the cutting edge toothless, quite even, the colour of head; the ligula the same colour too, rather large, circular fringed with sparse, fine, reddish hairs, with a rather large, rounded sinus in dorsal line on the front margin; eyes dark, the topmost much smaller than the next three lower, the colour very dark red-brown with a light, broad, subdorsal area on face; the fifth and sixth also very small; *Spiracles* of moderate size, those of segments 2 and 12 rather larger, all flush and yellowish or extremely light-brown, broadly oval. *Surface* dull, the segments quite distinct, covered all over with minute, erect, very light-coloured, numerous hairs which are very much longer round the anal margin where, also, they are reddish on the dorsal part. *Colour* light-green with a dorsal, darker, pulsating line. L: 25 mm.; B: 4.5.

Pupa.—Is more or less cylindrical in shape from the smoothly and evenly rounded shoulders to segment 9 after which it gradually fines down, the transverse section always a circle, until it reaches the hinder margin of segment 12; segment 13 is not apparent at all dorsally but is as a short band ventrally, the three segments 13-14 forming a conical piece, mostly slightly longer than broad at the base, ending, viewed dorsally, in a moderately broadly truncated extremity, this extremity being the cremastral portion, most of segment 14, flattened on top with fairly strong dorsal extensor-ridges limiting it laterally leaving the middle part sunken, the very extremity hollowed out ventrally and slightly bent down, each corner produced into a strong, chitinized, sharp, conical, short point about half as long as the distance separating

their bases, this distance slightly concave backwards, the edges of the whole piece as far back as segment 13 and between the points fringed densely with hairs which are twice or three times as long as the little teeth and stick out horizontally all round; segment 13 half as long as cremaster or segment 12 which are coequal (by the cremaster is meant the hard, flattened, chitinated end piece of chrysalis); segment 12 slightly shorter than 11; the thorax is twice as long as segment 8 including the bevilled margins of this latter segment, its dorsal line is mostly parallel to the longitudinal axis of pupa, ascending towards it from the straight front margin, descending from it to segment 4 in the posterior part, the hinder margin is a broad quarter-circle curve meeting the wings in a moderately deep, broadly rounded angle of about 70° ; segments 4 and 5 are coequal, about half segment 6 in length, segment 6 and segment 7 are again coequal; segment 8 is slightly longer; the head has the clypeus nearly ventral, rounded upwards and narrowed towards the base next the proboscis, the frons is in a plane perpendicular to the longitudinal axis of the pupa, much swollen convexly between the eyes, the vertex is inclined at an angle of 30° to the axis and is dorsal and broad—the whole head is broadly square in front of the shoulders—with a straight hinder margin; segment 2 is a transverse strip with parallel margins about the length of segment 5 and is hardly narrowed towards the spiracles; the proboscis reaches the ends of the wings but the antennæ only reach about two-thirds or three-quarters the way down, the midlegs ending between these two—the proboscis is *not* in any way free beyond the wings. Surface of pupa hardly shining, covered with a fine, short, reddish down which is decumbent and not nearly dense enough to constitute a pile; the abdomen being extremely minutely punctate rather closely, the thorax, segments 2 and head extremely finely aciculated transversely and irregularly. Spiracles of segment 2 are little perforations situated at the common margin of segments 2 and 3 and form the somewhat laterally ex-centric centre of a very shallow half-funnel formed by a slightly raised, red-brown semicircular surface lying on segment 3, a large semicircle with its height equal in length to the length of segment 2 and its breadth along the common margin twice that; the other spiracles are rather large, prominent, oval, with depressed, central slits, light brownish-yellow in colour. By the way, above and beneath each spiracle, slightly in front of it there is a small, oval, shining spot, very slightly raised from the surface and the abdomen is finely transversely shallowly aciculate as well as punctate, Colour is a soiled bone-colour with the cremastral portion red-brown and shining. L: 22 mm.; B: 6 mm.

Habits.—The habits of the larva consist in forming, at first, a cell by sewing together the edges at the tip of the leaf or blade; indeed this habit is continued right up to the end except that, at the end, when the larva is fully grown, it eats a great deal more of course and, always eating the blade on which the cell is formed, it finishes up by having the portion it lives in towards the tip hanging to the basal portion by just the midrib; it changes blades as required during this last stage. Finally, to pupate, it proceeds to turn the leaf over at right angles to the midrib, bringing the tip down towards the base, then eating both laminæ through to the midrib about a couple of inches from where it is doubled; it fastens the edges of the oblong piece thus formed together with web, rounding off the open end and, finally cuts through the midrib also so that the cell falls to the ground—much in the manner of *Halpe moorei* or *Plastingia submaculata*. It generally has been found close to the beds of nallas in the monsoon months and, sometimes, the cell falls into water; it mostly, indeed, gets washed down the hill-sides and gets swamped away down stream until caught up between stones at the side or it gets caught up before reaching the stream. The larvæ often make two or three attempts at a cell before finishing off a final one. This cell is

lined inside with silk and especially at the open end there is a dense covering or lax closure of feathery, cereous matter mixed in with the silks which makes a very effective bar to any water getting in. The pupa is not attached inside the cell and is generally—always—placed with its head *away* from the open end. The imago is rarely seen and is an insect of the damp, heavy rainfall ghat portions of the Western Ghats in Kanara. It flies fast, is fond of sitting on the ends of leaves high up, basking, and is very difficult to catch. The foodplant is *Ochlandra talboti*, Brandis.

Swinhoe pictures the species on Plate 322 in *Lepidoptera Indica* Figures 4 male and 4a female, 4b the underside, and says that it has been obtained in the Nilgiris, the type being in the B.M. from there. It was, it is believed, collected by Sir George Hampson; he says that he has it in his collection from the same locality and that Davidson, Bell and Aitken record it from Karwar. N. Kanara would be more correct because the insect was never seen in Karwar itself on the very sea shore. It came from Anshi on the crest of the Western Ghats at about 1,500 feet in the evergreen areas where the rainfall must be at least 200 inches in the year if not more. The first specimen seen was caught there with a net in the year 1895 and it has not been seen flying again. It was caught in the same nalla in which afterwards the larva was obtained, curiously enough. It is of course an insect of powerful flight but little or nothing is known otherwise of its habits.

223. Halpe (Thoressa) honorci, de Nicéville.—Male. *Upperside*: brown with an ochreous tint; spots dark orange-ochreous. Fore wing with the basal half covered with short, ochreous hairs in the interspaces and the following semihyaline markings:—2 mm. long spot in the cell with its outer edge just at middle of wing and filling the cell up to and down to its borders, quadrate or constricted into a neck of varying thickness in the middle along the fold in cell; a discal one in interspace 2, its inner edge below outer edge of cell-spot, about twice the length or somewhat less with another above it in interspace 3 that is half the length of it (equal to length of cell-spot), out-reaching the one below by half its length; with three subapical spots in interspaces 6, 7, 8; one above the other, the topmost a mere dot, the bottom one only slightly smaller than the one interspace 3, the middle one between the uppermost and lowest and the inner edge of the three in a perfectly straight line; besides an orange, not semihyaline mark in interspace 1 that is chevron shaped, the sharp end innermost, the top are linear and slanting obliquely from lower, outer corner of spot in interspace 2 to above the middle of vein in the middle of interspace, the lower arm much wider, thicker, slanting outwards to vein 1 at about three-quarters its length from base of wing; the yellow scaling leaves a bare space of ground-colour, oval in shape, bordering each spot in interspaces 6, 7, 8 and 2, 3 outwards. Hind wing. Extraordinarily variable as to the extent of the orange-ochreous patch. One specimen caught in the very wettest month of the monsoon has the whole of the interspace 4 including some of 5 above of that colour extending from the discocellulars (end of cell) outwards to three-quarters its length just to embrace a small longitudinally-oval spot of ground colour in its extremity: and also interspace 3 (the brown vein separating the two narrowly) for the same distance out embracing another, similar spot of ground colour: and, further, the basal half of interspace 2 with the spot just beyond its extremity—these coincide with the postdiscal, curved series of brown spots sometimes present on the underside; these spots covered with long, concolorous-ochreous decumbent hairs while the rest of the cell has them black and there is a line of similarly—long, not quite so decumbent, orange-ochreous hairs all along vein 1b to close to outer margin. In another

specimen also caught in the rains the cell underneath the (here) concolorous, long and decumbent hairs (none are black) is all orange-ochreous as well as the whole of interspaces 4 and 5 to near outer margin extending upwards slightly to top of interspace 6 beyond middle and downwards through interspace 3 to completely fill 2 to close to outer margin, the whole of the interval separating vein 2 from 1b with its fringe of hairs being covered with orange hairs also. In still another specimen caught in January the patch is as extensive as in this second one but has no sign of any brown spots. *Underside*. 'Fore wing with the costa increasingly, the apex broadly, the outer margin decreasingly dull, dark ochreous; all the rest of the wing blackish, the spots as on the upperside, the discal series with the chevron mark much paler, diffused, larger. Hind wing dark, dull ochreous; a thin, black bar closing the cell, a complete, discal, curved series of long-oval, blackish spots, one in each interspace commencing with one above middle of cell towards the base of interspace 7 and one beyond its middle, the last in the middle of interspace 1b; all veins blackish the abdominal margin and a streak in the basal two-thirds of the submedian interspace black.' (Swinhoe, *Lepidoptera Indica*, vol. x, p. 287.) Here, again, the marking is variable. In some specimens it is absolutely uniform dark-ochreous without a sign of any marking; in others Swinhoe's cell-bar and postdiscal spots are all there, only the black streak in the submedian interspace and on the veins is wanting; in others the submedian interspace is certainly darkish as is also the costal area above vein 8 and there is a submarginal series between the veins of dull-brown spots that may be quite obscure. And, which is peculiar, these variations are not dependent upon dry and wet seasons, nor upon sex. Antennæ with the shaft black but a yellow, thin streak on the underside towards top in some; in others slight indications of yellow on the undersides at base of each joint, the underside of club at base extending down along the shaft yellow, the apiculus dull red extending much further down on to the club above than below. Palpi brown above with some ochreous hairs, below pale-ochreous at base shading up to orange at end in some fresh specimens. Head and thorax orange above or ochreous-orange with a plentiful admixture of brown. Abdomen brown with very plentiful admixture of orange-ochreous scales on distal half. Legs orange-ochreous or pale-ochreous according to specimens. The fringe of hairs along inner margin of fore wing slight. Cilia of fore wing brown with darker base, the portion below vein 1 often ochreous; of hind wing ochreous with brown bases, altogether slightly brown in interspaces 2, 3 both above and below. Expanse 40 mm. or rather less for female; the male often as much.

The crook of the antennæ is bent back at right angles and short and the hind tibiæ are quite naked or with a very slight fringe of hairs.

Swinhoe says in his book: 'Female. *Upperside* coloured like the male. Fore wing with the spots similar, but there is only one spot below the two discal spots. Hind wing with the patch usually restricted to a conspicuous streak in interspace 4 and inconspicuous, thin streaks in 3 and 2. *Underside* paler than in male; all the markings similar but indistinct.'

Larva.—This caterpillar is, in *shape*, very like that of *Halpe moorei* but is altogether *sui generis* amongst this type of skipper larva as it is not concolorous; it is longitudinally marked with a dorsolateral, broad, rose-coloured band from segment 2 to segment 13, the dorsal area above it being more or less yellow, the spiracular area below it, white. The 13th segment is a narrow, transverse, parallel-margined piece, shorter than half the length of segment 12 and with a very slight dorsal slope backwards towards the anal segment; that segment, i.e., the anal one (segment 14) is semicircular in outline has a slightly greater slope than 13 and is about the same length as segment 12; the legs and prolegs are short; segment 2 is whitish and has a thin, transverse, black collar across its middle which is very slightly interrupted in the dorsal line; the body is thickest at middle and decreases slightly towards both ends; there is a distinct neck before the comparatively large (higher and broader than segment 2, equal about in diameter to segment 3), nearly round, thick *head* which is only very slightly indented in dorsal line on vertex between the lobes; the surface of head is shallowly and minutely honeycombed-rugose with a somewhat sparse covering of very minute, appressed, short, light hairs with some longer, erect, stronger ones on the edges of the ligula and around the mouth-opening; the clypeus is rather less than half the height of the head, a good deal higher than broad at base,

triangular with an acute apex, the false clypeus outside it reaching slightly further up than half the height of the head, the strip formed by it outside the true clypeus rather broad, broadening from base upwards to nearly three-quarters its own height in a gentle curve, then the sides converging to an acute apex in dorsal line; the labrum nearly white, shining, transverse, not short (really the same colour as clypeus and head generally), the ligula also the same colour, longer than broad, semicircularly rounded with a rather deep, triangular sinus occupying the centre of the front margin; the antennal, basal joint light, the second rather dark red-brown; the mandibles red-brown also with much darker edges, this cutting edge straight, not toothed; the eyes: 2, 3, 4 in a straight line pointing to 6 under the lower curve of head, 1 behind and above 2 at an obtuse angle, all the first four equally spaced, 6 twice the distance from 4 that 3 is from 4, the 5th behind 4 and 6 and forming an equilateral triangle with them, all dark. *Surface* of larva covered with extremely minute, erect, soft, light hairs, hardly visible under the lens; otherwise smooth, the skin thin with the tracheæ showing through; the anal segment largely chitinized on dorsum and very irregularly wrinkled-rugose, black; the thin, black collar across segment 2 also chitinized. *Spiracles* of ordinary size, nearly circular and yellow; those of segments 2 and 12 larger than the others. The *colour* of the body is peculiar for one of the genus: there is a spiracular white band above which is a dorsolateral, rose-coloured band of about the same width with a shade of green in it, above this a subdorsal, slightly narrower, yellow band and a dorsal, greenish line; all longitudinal reaching from segment 2 at the hinder margin to the front margin of segment 14; the head is light brownish-yellow with, generally, a round, black or blackish or brown, single, rather small spot near the top of the face on each lobe. L: 29 mm.; B: 4 mm.

Pupa.—The pupa is also, in *shape*, like that of *Halpe moorei* except for differences that will be evident below; it is, however, very peculiar in the excrescences of the head and in the formation of the cremaster; the body is thickest and stoutest generally about the middle though the difference between the stoutness there and at the thorax and shoulders is very small; the shoulders are evenly rounded, the eyes prominent, the front squarely blunt, the thorax slightly and evenly convex-humped, the abdomen decreasing in width backwards more quickly after segment 7 to the narrowly square-ended cremaster; segment 13 is more or less intimately soldered to 14 and is about half its length (slightly more) and also half the length of segment 12; segments 11-13 with a gentle dorsal slope, segment 14 much more strongly inclined downwards; the hinder margin of thorax is a short parabolic curve about equal to a semi-circle and meets the wing-line on each side in a broadly rounded angle of about 90°, segment 4 is about equal in length in dorsal line to segment 5, segment 6 is not quite twice as long, thorax is four times as long as 4, segment 2 is only slightly shorter than 4; the *head* is very peculiarly sculptured the whole frontal surface, inclined at an angle of about 80° to the longitudinal, axis of the pupa, is flattened between the eyes and raised very considerably above the pupal surface into a heart-shaped shield which is broader than long and has the heart-sinus posterior; this surface is very prominently and irregularly wrinkled-rugose and is separated from the head-vertex by a deep, linear depression of triangular section: the vertex behind this is also similarly raised and roughened in two pear-shaped pieces, one on each side of the dorsal line, the thin end of these pears touch the dorsal line and the broad ends reach down to the eyes, the breadth (in sense of length of pupa) is only half the length of the vertex, the rest of the vertex-surface being flush with the general pupal surface but roughened like the raised front pear-like portions; this flush vertex-portion is again divided into a dorsal, triangular part bordered in front by the pear-like, raised pieces with the base posteriorly formed by a fine line formed by the front borders of two longly oval, marginal pieces, one on each side of dorsal line and lying along the hinder margin of head; segment 2 is normal, not roughened, parallel-sided and in same plane as the head-vertex; the eyes are roughened in front; the cremaster is peculiar: it is composed of two pieces, the anterior nearly square, as broad as segment 13 (less than segment 12) and has its dorsal surface gently concave, its lateral edges raised considerably though somewhat thinly into sharp, low ridges leaning outwards with three or four backwards-directed, short, sharp, conical teeth; the posterior portion rather less than half the length and, at its

extremity which is quite square (the extremity of the pupa) rather less than half the breadth of the basal portion; this end piece has its sides gradually diverging forwards to bend suddenly out to the hinder corners of the anterior, square piece; this pupal end is furnished with a somewhat diffuse bunch of stiff, golden-brown, suspensory shafts which are somewhat ventral and hooked at the ends; ventrally the cremaster has nothing noteworthy; the antennæ reach half way down the wings, the proboscis reaches the ends but is not prolonged. The *surface* is moderately shining, the abdomen finely granulate-rugose, the thorax more or less regularly, transversely aciculary rugose in parallel lines, the wings irregularly aciculary lined transversely; besides the rugosities on head and eyes noticed above; the segment margins distinct, narrowly satiny-rugose along hinder borders, the margins of segments 8, 9, 10 more broadly satiny. *Spiracles* of segment 2 are guarded behind by a very prominent, 0.75 mm. long, more or less oval surface or protuberance which faces forwards and has another, oblong, similar, raised surface in front of it, separated from it by a depressed line, this anterior, raised body black, the posterior one red-brown; the other spiracles are very light yellow, moderately broad ovals, flush, rather small. The *colour* of the pupa is waxy-white all over with a green shade, the rugosities of head, and eyes are all nearly black (very dark red-brown); the ridges of cremaster and teeth are rather golden red-brown; besides which the greater part of the lateral region of segment 2, a short, subdorsal streak (longitudinal) from front margin of thorax backwards and a similar dorsolateral streak on segment 1 (head-vertex) are black. L: 25 mm.; B: 5 mm.

Habits.—The larva is very shy, eating in the evening. Pupa in a longitudinally joined bamboo-leaf and fixed by tail; coating of fairly thick silk inside of cell, sometimes steps of silk also; whole cell very tightly closed. The larva generally lives amongst the dead, yellow leaves and makes its cell of several joined together. Is found both above and below the ghats, at Siddalgundi and Karwar. Four or five were obtained in March 1919; the first came from Siddalgundi on March 15, 1893; another from Udamakki on August 1, 1893; all in the N. Kanara, District of the Bombay Presidency. The butterfly is a powerful flier, frequenting the jungles where bamboo grows; keeping much to the higher stories of the trees where it goes freely to flowers for the sake of the nectar in the mornings up to eleven o'clock if there be shade, not so late if the sun is too bright. It was much attracted to the flowers of the Dividivi or *Cæsalpinia coriaria*, Willd., the American Sumach, a cultivated tree with extremely sweet smelling flowers that come out in October and June. The insect is not often seen; neither are the caterpillars easy to come by. Swinhoe gives the habitat as South India and Ceylon and says 'The types in the Indian Museum, Calcutta, came from the Palni Hills; Hampson records it from the Nilgiris; Elwes from Trichinopoly; Davidson, Bell and Aitken who bred it in Karwar, did not figure the larva and pupa.' The larva and pupa have, as a matter of fact, been painted but never published.

(To be continued)

THREE MONTHS UP THE VALLEY OF THE SUTLEJ RIVER

BY

LT.-COL. R. W. BURTON, I.A. (RETD.)

PART II

(*With 2 plates*)

Continued from page 39 of this Volume

III

*' To sit on rocks, to muse o'er flood and fell,
To slowly trace the forests' shady scene,
Where things that own not man's dominion dwell.'*

—BYRON.

At 4 a.m. on May 30, the first day's *shikar* was commenced by an expedition up the main valley in search of ibex. Some climbing over precipitous rocks, with many a precarious hold above the raging stream below, opened out to better ground along the right bank of the river. About two miles from camp eight ibex were sighted. All had horns of upwards of thirty inches, and one buck appeared to be the owner of a fairly good head. Going further on more ibex were seen, eleven females and one small male. The party of eight came along and joined these. We followed the herd, they up the hillside we along the river, until there came a precipice which would force the animals lower down and so nearer to us. The best head appeared to be about thirty-eight inches, so I decided to take the chance that was offered. The distance was about 300 yards and I was that day, for the first and last time, using the .280 Ross Rifle. No dust flew to the shot. The animal galloped a few yards and disappeared behind some rocks. The remainder of the herd were carefully counted across an open stone shoot which lay in their way. The big one was not among them, therefore he must be hit, and was possibly dead. Ramdass staying with me, the two men were sent up the valley to cross by the first available snow bridge and make their way back along the opposite side. Watching them I envied the ease and speed with which they moved.

They found the stricken beast lying down where he had disappeared. He galloped across the stone shoot and stood among rocks; but the distance was now greater than before: quite 400 yards. The last of four shots brought him rolling down the hill to come to anchor against some bushes. The men got the carcass down to the edge of the river and, after several attempts, I passed across a rope

which, fastened round the neck, enabled us to haul the body through the rushing torrent of the icy stream. This saved much trouble: but he was very nearly lost, the weight of the body being almost more than old Ramdass and I could manage. The animal had been hit four times. Three shots were through the body—one of these through the shoulder—and the last had smashed the base of the skull. Having killed him I wished I had stayed my hand as the horns measured only thirty-six inches. He was in splendid winter coat. After this experience of the .280 rifle I decided never to use it again. The .375 would have killed the animal at the first shot. On the way back to camp tracks of a snow leopard were seen along the river bank. There was a curious yellow halo round the sun. The men said that it portended continuous fine weather.

The last day of May was spent in camp, seeing to the ibex skin, doing odd jobs of mending, and making preparations for a four days' bivouac further up the river.

The dawn of the first of June disclosed a cloudy sky and snow falling on the higher hills, so on this occasion the halo was at fault. We made an early start, the party consisting of the three *shikaris* and Panchram, one of the Chakrata coolies. We went far up the valley and, turning up a side ravine to within the snowline, the men settled themselves down in the ruins of a last year's shepherds' hut. My bedding was laid out at an angle between two low stone walls which kept off the wind, and a green waterproof sheet rigged up overhead furnished sufficient shelter for ordinary weather. On the whole the camp was quite a good one: fortunately there was neither mist nor snow. The hills round about were very rugged and precipitous but had no terrors for me. Wearing the woven grass shoes brought from Chakrata I could go over any sort of ground in safety.

The first day's excursion, still further up the main valley, was made over difficult country, steps having to be cut in several places across the steep slopes of frozen snow. The junction of the Spiti nala with the Thamin river—now reduced to but a miniature of itself—was reached, and it was made certain that no ibex had yet gone beyond that point. There were no tracks in the snow and no grass had yet appeared. Several lots of ibex were seen on the way up, but no head worth having. On the hillside opposite camp was the herd of eight ibex which had been reduced to seven. Our quest being now for bear and not ibex, to remain further at this bivouac was useless, and a return to the main camp was made the following day. Some wild onion was found, also two kinds of rhubarb, the variety with the smooth leaves being less bitter than the crinkled leaf kind which is mostly found at a higher elevation. Both of these, with plenty of sugar, formed an acceptable addition to camp fare.

The camp on the right bank of the river was a very pleasant one, well sheltered from the wind, which, coming as it did from regions of perpetual ice and snow, was bitterly cold. The sun gave us its grateful rays during sufficient hours of the day, and there was plenty of fuel. As is the case with all snow-fed streams, the river rose considerably each afternoon and became turbid with silt. The roar

of the waters and deep rumble of heavy stones being slowly ground to fragments by the rushing torrent was most soothing and conducive to sleep after a tiring day.

On June 4th the *shikari* coolies went to find a way across to the country east of the Pamachang stream and returned in the evening with news of fresh tracks of snow bear. A long round of some five miles brought us next day to bivouac on the opposite side of the river and but a stone's throw from the camp we had just left. On the way tracks of a snow leopard were again seen, as well as those of a small snow bear. Both of these animals had passed along the further side of the river, but a short sixty yards from my tent, during the previous night. The tracks of the bear led over our bivouac ground and away up the hill.

In the evening, following up these tracks, we came across the remains of an ibex, killed some five or six days previously by a snow leopard, as could be seen by the marks in the skin of the throat. It had been eaten by him, and a large red bear had also been at the carcass. The leopard had bagged a better head than I! The horns measured 42" by 10 1/2" in girth. Randless said that heads better than this are not to be found in that country, and he was probably correct. The night's tracks of the big bear were found higher up the hill, after we had tracked him to his lair of the previous day among some thick bushes, but no view of him did we get although we waited until it was nearly dark. Search next day showed that he had not again been about the hill so it was decided to move to a camp higher above Rupa. Had time permitted a few days' further stay would probably have enabled this fine bear to be bagged. That is the worst of not concentrating on one description of animal where animals of any kind are so few. Burihel were also wanted.

At dawn next morning it was found that the big bear had again been all over the hill. Every possible cover was searched but he was not to be found. *Shikari* men said he had smelt my footsteps and so decided that it was safer to sleep elsewhere during the day. Some ram-chukor were seen, and the tracks of the small bear followed the high-level path, leading towards Rupa for several miles. About twelve o'clock three burihel rams were sighted in a wide valley. Almost as we saw them the cries and song of a cattle herd far below caused them to begin to travel up hill. The leading ram had fair horns, the second one a small head, and the last of the three carried a good head: probably 26 inches. At first it was doubtful which way they intended to go, and how far. The leading ram was pretty to see as he pugnaciously rose on his hind legs and butted at the second one. He did this several times, on one occasion making a leap of about fifteen feet which the other fellow wisely let pass without attempting to meet the downward assault.

After five minutes' watching it became evident that the animals intended to cross over, high up on the same spur as we were on, so no time was to be lost in going up our side of it. The elevation was some 12,000 feet, and progress, on my part, somewhat slow. The impatient *shikari* cooly had to wait many times for me, and the burihel were a long three hundred yards' distant when I

pantingly gained the shelter of a bush at the hump of the spur. That five minutes' delay, watching the animals, had been fatal, and lost me the big ram. With these few minutes in hand, we would have been under cover waiting for the rams to file quietly past at short range. Having waited for the burhel to disappear over a ridge half a mile or more away, time was given to them fully to disappear; for sometimes a ram will return to the crest and re-survey the country just passed over. We followed as fast as possible. The *shikari*, leading this time by a couple of hundred yards, gained the crest and at once frantically beckoned to me—poor breathless and winded me—to hurry up! Doing my best I arrived much pumped, to see two of the burhel disappear round the next spur, probably for good, as the next sight of them would be at long range. The biggest ram was standing in the little valley below us, only a hundred yards distant, gazing up at the ridge. The *shikari* cooly had evidently been incautious as to how he looked over on his first getting there. Old Ramdass would not have committed such a fault. It had been necessary to let the cooly push on in order to keep the animals in sight. No time was to be lost: but I was in no condition to shoot straight. The wobbling .375 sent a bullet just under his chest. There was now time to gain breath, judge distance, and make ready in lying down position for the long shot which it would be necessary to attempt; and which was justified by the circumstances. Animals were difficult to find, and remaining days for *shikar* were becoming alarmingly few. Two of the rams soon appeared, making their way up the rocky spur which we knew must be their way of retreat. They were the small and the big ram. My rifle is oversighted, as are almost all rifles of the H. V. class. With the 200 yards lead, good enough in that rarefied air for any distance up to 400 yards I nearly got the big ram: missed him by inches only, just under his body. This gave me the exact range, but the ram disappeared among the rocks. Then the third ram, the former-leaping leader of this small party of gentlemen, came into view. The first shoot went just over him. He stood, fortunately, and the second taking him in the near hip, broke both his thighs. The distance was over 350 yards. He lay where hit, then slid down a small shale slope until checked by a large stone. Lying there, his colour blending with the shale, he was invisible to the naked eye, an object lesson of how marvellously wild animals can at times suit their environment.

The two other rams again came into view among the tumbled rocks. The big fellow bore a charmed life. The two bullets went exceedingly close: one, the first, must have been not two inches over his back. I saw the spurt of snow dust. This was with the 300 yds. lead. He made a standing leap of at least fifteen feet, obviously untouched, and again stood at gaze. On being missed a second time he quickly disappeared amidst the snow covered rocks behind which his young companion had already been lost to view. His agility was astounding and equal to that of an ibex.

To get to the fallen ram took me some time, there being much bad and difficult ground to be negotiated. He was in thin condition: horns 23" by 12" in girth. Having secured a

photograph of the prize we made our way downwards, meeting the remainder of the party eagerly climbing towards us to find out the result of so many shots. We got to camp above Rupa by five o'clock in the evening, having been twelve hours on the go and much of the time over vile ground. After a stay of one day in camp a move was made to a bivouac at a much higher elevation, about 13,000 feet, the climb being fairly steep. On the way up five burhel ewes were seen. The sight of three more in a valley beyond the bivouac raised hopes that perhaps more game would be found on the morrow.

The sky was now quite clear. On all sides nothing but snow was to be seen. The mountains extended in all directions, range after range, peak after peak. To the north, some two miles from camp, was the snow covered range dividing the valley of the Thamin from that of the Spiti river. To the north-east could be seen the Hangrang pass leading to Hăngō. The path to it, which started from the village of Sūinām, could be seen now and again as it wound upwards over the rounded spurs of the mountains, until it disappeared from sight above a stupendous precipice of reddish sandstone, streaked with strata of many hues. The glories of the near and distant mountains were spread out before me—high ragged ranges shooting up into the sky in all directions, the vision lost in a sea of peaks. No wind, not even a distant sound, disturbed the silence.

The next morning an early start was made, as it was intended to cover all possible ground in the search for burhel. Quite close to camp a lovely yellow fox was seen. He had a white tip to his tail, and stared unalarmed as we made our way along the hillside through the stunted gorse-like juniper. Valley after valley was searched without success and the *shikaris* confessed themselves unable to say where the burhel rams could be found. During the day two more yellow foxes were seen. They were a long way off, and the men insisted they were a small kind of panther and were the cause of our being unable to find burhel! My telescope showed very plainly what animals they were, but the men remained wilfully unconvinced. One of the foxes was playing very prettily in the snow after the manner of a happy dog.

It had now to be decided whether to remain in this valley and make an effort to bag a snow bear, or to retrace our steps to Pangi and try for burhel in the neighbourhood of Charong. A man was sent down the hill for baggage coolies; a decision could be arrived at in the morning when camp was struck and the loads packed: and so passed a second night under the glorious canopy of the starlit heavens.

'The 11th June, and Charong it is!' The coolies arrived at seven o'clock, so, starting down the hill by eight, Rupa was passed and Suinam reached by two in the afternoon. The *shikari* men were quite pleased with ten rupees each as pay, and two rupees eight annas each as *bakshish* for animals shot. Twenty rupees a month is good pay for *shikaries* in those parts. They asked me to come again, and were eager to take me further afield, so perhaps I overpaid them. They said that March would be the best

month as all animals would then be at low elevations and snow leopards would not be difficult to get.

At Rupa I took a photograph of the village, ascending to the roof of the younger *shikari* man's house for the purpose.

The names of the men were Charung Tanjan and Tikaram, the latter being a very intelligent youth. I took a great fancy to him and would gladly have him with me again. The temple at Rupa is of the usual type for this part of the world. A very gaudy gilt covered image of Buddha was the principal feature of it. All round the walls were paintings of Lamas and devils executed by an artist from Spiti. A man of the village had lately died at Pangi on his way up from Rampur and a number of red clad Lamas were holding a memorial service, so I gathered. At Chāmong 'Motee' was taken over in fair condition and four rupees disbursed as his keep for sixteen days. On this day the last of the burhel mutton was consumed; and glad I was, as the muscles of my jaws were getting enormous from the heavy exercise. Never was there tougher meat. When the animals are shot in autumn, after months of splendid grazing, the flesh is excellent; but in the spring, after a winter's diet of hardy shrubs and such like fare, the reverse is the case.

The return journey down the valley showed how much the aspect of the country had changed during the past fortnight. The poplar trees were now in leaf; snow bridges fast disappearing; hillsides green; and people busy in the fields. Near Suinam village a number of children from four to ten years of age were being made to weed the fields. They began early in life to learn to cultivate; to carry loads; and to tend the village stock. A small baby was brought to me to have its foot doctored. There had been a severe burn but the wound had sufficiently healed. After application of some ointment it was bandaged up, and the mother went away highly pleased with my safe assurance that all would be well with the little one's foot in a few day's time.

It was hot, toiling up the hill from the iron bridge to the Benang Pass. Nothing had yet been done to the 'road'. That portion of this road between Shiasu bridge and Poo—the mission station below the Shipki Pass—was said to be so bad as to be dangerous; it being not uncommon for laden animals to fall and be killed, with total loss of goods to the trader. This state of affairs struck one as somewhat strange in view of this being one of the trade routes between India and Tibet; and the fact that a British Trade Agency was established at Gartok so far back as the year 1904.

On arrival at Jangi on June 15, the unwelcome news was received that no extension of leave could be given as circumstances necessitated my transfer to another cantonment: so next day I set out for Charong, crossing the Sutlej by the 'Jhula' below Akpa village and proceeding to camp at the picturesque village of Rispa. The 'Jhula', or swing bridge, was a substantial affair of galvanized rope wire, the means of crossing being by a wooden seat hung on the main rope and pulled backwards and forwards from either side. The cooly Gangaram, a man of timorous nature,

was unashamedly terrified at having to be swung in mid-air over the roaring river! I had him tied in as a matter of wise precaution. Some of the baggage damsels who came from Rispa to take on the loads were very nice looking, and more extremely pretty women were seen up this valley; also many men, and women, who were the reverse of good looking.

The day's journey of June 17th took us from Rispa to Lamber *via* Tangi, the first mile or so being along the left bank of the Sutlej. Leaving the main river the path turned up the Tadoong Gad, to cross it a short distance up by a ford, and then wind up the hill sides, mainly through fir woods, to the village of Tangi, which occupies a fine commanding position on an open spur of the mountain. Looking backwards there were very fine views of the Kashang and Ashang Valleys and the snow covered mountains encircling them. After Tangi the path descended again to the bed of the nullah, and from there the remaining two miles to Lamber was mostly a scramble over the boulders of the bed of the stream.

At Lamber the people told me that the rest of the 'path' to Charong was of the same description and so I found it, being fully satisfied by evening with the exercise afforded by this tiresome march of thirteen to fourteen miles.

There was an earthquake near Lamber in April of that year, and a large piece of the hill side tumbled into the river. The occurrence took place at night and gave the villagers a great fright. Three men of Charong were on their way home and, bribed by a box of matches, sat for their photograph, which turned out to be a very good one. Three uglier countenances never faced a camera.

The gorge of the Tadoong stream from Lamber to the junction with the Kuno nullah is a most imposing one. Many of the mountains thereabouts are quite inaccessible, the precipices being immensely grand. Within six miles on either side the mountain ridge is over 20,000 feet, some of the peaks being as much as 21,800 feet in height; so it can be imagined how precipitous are the mountains, rising as they do from the river bed (10,000 feet or so) to those great elevations. At one spot there was a glorious waterfall, many hundreds of feet in height, thundering down the perpendicular cliffs. There was a large volume of water, all melted snow, rushing down the rocky stream: but it was merely a trickle compared to what it would become as the summer months progressed.

Coolies were changed near Kuno: the path from this place to Charong being quite good, camp was pitched by three in the afternoon. Late in the evening, standing at the tent door, I sighted two burhel ewes on the opposite side of the river valley.

IV

*To climb the trackless mountain all unseen
With the wild flock that never needs a fold!*

—BYRON.

The local *Shikari*—Durjan by name—was about to start off to Tibet on a trading expedition, but was induced to remain for three

days. Camp was moved on the morning of June 19th to about seven miles up stream from the village of Charong. On the way a good many burhel were seen, but none carried horns of much over 20 inches. At daybreak on the 20th, accompanied by Durjan and two coolies, I went some miles further up the bed of the river. Three lots of burhel were found—twenty-four animals in all—but there were no good heads among them. A lofty hill was climbed much of it being very steep, and a view of the entire opposite hillside obtained. There were many burhel scattered about, and it was interesting to watch them. At about 9.30, they all began to move upwards among the snow to lie up for the day. The younger rams were very playful, sportively butting at one another, and it was a pretty sight to watch their gambols.

Higher and higher we climbed to attain a ridge well within the snow line. From there a splendid view of the magnificent scenery was obtained. Nothing is more fascinating than to roam along the crest of the higher ranges in mighty mountains. You feel that the whole country below is yours: in it you are supreme. Also you feel what a very very insignificant creature you are! In the vast sea of mountains and valleys unfolded around you, you feel a wild, unfettered freedom; you follow the sheep trail, always a good path, leading you securely round the pinnacles: with strained excitement you walk slowly along, every step bringing new pictures in view: you look on a vast bewildering landscape of lofty ranges rich in colour, abounding in sculptured crags, deep ravines, and green basins—the desolation relieved by the shrill call of the ram chukor and the majestic sailing of the lammergeyer in the brilliant blue of the sky.

In this way we carefully wound our way along the slopes of the mountains for several miles, thoroughly searching four large valleys and, gradually descending, to finally arrive at a magnificent grazing ground of wide undulating grass downs. A few ewes were all we saw. We then made our way by downward sheep paths to the main nullah, and so to camp—a long dreary trudge over the slippery boulders of the stream. At one place the snow bridge, by which a crossing was imperative, was broken down, leaving a wide and gaping fissure over the torrent below. This had to be jumped and it was a relief to be safely across. Further on the snout of a small glacier was passed, the twenty foot thickness of ice showing a deep emerald green where it was sharp cut and overhanging the bank of the stream. Camp was gained just as the last rose tints of the setting sun faded from the fleecy clouds surmounting the perpetual snows. The elevation attained during this day, the most exhausting with exception of one strenuous day in Kashmir in 1923, which I have ever experienced, was well over 15,000 feet.

Perseverance is often rewarded. The alarm clock was set for an early start the following morning, the intention being to again look up the many animals seen the previous day, with the forlorn hope of finding a shootable head. Standing at the door of the tent just after day-break Durjan drew my attention to some animal far up the hillside above the camp. The field glasses

showed three rams of which one possessed a good head. The fine sweep of his horns was silhouetted against the sky as he stood on the brow of the hill. In a few minutes the animals disappeared from sight, and we at once started in pursuit.

The climb was not very steep and the distance about 1,200 yards. Arrived at the top all likely ground was searched in vain, so the *shikari* started back to pick up the tracks and find which way the animals had gone. I sat down near a bush with rifle and field glasses ready. Having proceeded only some sixty yards up the slope the *shikari* whistled and pointed. Two burhel were just topping a ridge about 120 yards distant. They were among rocks, and difficult to define. Quickly putting up the field glasses one of them was seen to be the big ram. He was gazing in my direction, and only his chest and part of one shoulder was visible. The shot, quickly taken, was fortunately a good one, the bullet striking the point of the shoulder and making its exit behind the shoulder on the opposite side. The ram fell dead in his tracks and his smaller companion fled clattering down the hill. I was delighted to find the horns taped 26 inches. A very handsome trophy indeed. The cooly with us carried the animal on his back up a steep slope on which I had to go, in places, on all fours! He was helped, certainly, by the *shikari*, who steadied him by holding on to a horn, but the performance was pretty good as the animal weighed 140 lbs. He carried him with ease down to the camp.

The bagging of this burhel was a great stroke of luck which I felt was deserved, and steps could now be retraced in hope of a snow leopard, said to be killing sheep near Kuno, and perhaps a snow bear up the Kashang Gad from Pangi. The *shikari* assured me that this 26 inch burhel was the biggest as to horns that he had ever seen shot. That the burhel of these parts *do* grow larger horns was certain, as one—a very old and much weather-worn head—which I saw at Ramrikcha temple near Charong measured 28 inches, and must have originally been quite an inch longer than that. The people told me the many horns at the temple were those of animals killed by snow avalanches, but I suspected that a number of animals were killed by being driven in the snow during the winter months.


The *poojari* at Ramrikcha temple had some scars on his face, the result of an attack by a snow leopard. It seemed that a few years previously a snow leopard got into a hut near the temple, where sheep were kept at night, and killed twenty-one of them. The next evening he was around the place again. The *poojari* saw that the murderer was after some goats so went out to put them into a hut. The leopard apparently resented this as he went for the priest and clawed his face! That same night the leopard visited Charong village, a mile away, and attacked and clawed a woman there. The villagers decided that the animal must be killed, so tracked it to where it was lying up among some bushes about half a mile off and, surrounding the place, slew it with stones and sticks without damage to themselves. A plains panther would not have died under those circumstances without leaving his mark on a few of his assailants. At Chakrata a panther—a small



A FINE BURHEL— $26\frac{1}{2}$ "



IN THE TADONG GUD
'River, Ravine and Splintered Precipice'



THE TIMOROUS GANGARAM CROSSING THE SUTLEJ



THREE HANDSOME MEN OF CHARONG

female—was brought to me for the Government reward, having been killed by a villager with a blow from a hatchet. The skull was cleft to the brain. It had attacked the man after being wounded by a companion with a gun.

On June 22nd camp was moved to Kuno, the *shikari* coming with me as he was eager to catch up the people with whom he was to go to Tibet. His way was up the Kuno nullah at head of which is the Sholarang pass,—six marches—all the way being over the rough bed of the stream. I did not envy him his journey. The snow fall at Charong is heavy and the people cannot leave the village during the winter. A few of the men go to the lower hills with goats and sheep, but most of them, and all the women and children, remain in the village. A great store of firewood has to be collected, and all grain has to be ground by hand as the frozen water mills cannot be worked. The principal occupation all those weary winter months is weaving cloth and knitting.

On leaving camp for the march to Kuno I went up the hill where the burhel was shot and was rewarded by the magnificent view in the clear morning air, also a photograph of Charong Peak 19,000 feet—which had been under cloud on the previous day. Towards evening it was evident that bad weather was coming on as it began to rain; mists were stealing along the higher slopes, while all the mountain tops were hidden in clouds.

At Kuno there was no further news of the snow leopard. Days were precious, so it was decided to make for Pangri without delay. There was rain during the night and a village dog seized the occasion to steal half a pound of butter, also the burhel head which was, fortunately, retrieved undamaged. A group of the women and children of Kuno was taken, as also a photograph of a 'zho' the half cow, half yak, of those parts. Camp was struck about nine o'clock, after the tents had dried, and the march to Lamber accomplished by the evening. Consequent on the recent rain and more rapid melting of the snows, the river was now more noisy and turbulent and had greatly increased in volume. The water was of a light coffee colour instead of the lovely green and blue hues seen on the upward journey.

During the early part of the next day light rain was falling for some miles of the way, everything getting thoroughly wet. All were glad to find sunshine and dry weather in the lower reaches of the valley. Many people from all parts of the country were met journeying to the Ramrikcha temple, at which an annual fair was to be held during the next few days. From Rispa to Rorung the journey was uneventful. Gangaram was once again in a great funk at having to cross the Sutlej, now far more formidable in appearance than on the sixteenth of the month. He tied himself in like a piece of luggage, and, seeing the state of his nerves, the merry damsel at the winch kept him dangling in mid-air over the raging flood. Shouting with glee she pulled him backwards and forwards in tantalizing fashion before winding him in to the safety of the landing platform. One cannot imagine a woman of the plains doing that!

The walk to Rorung was a very pleasant one through terraced fields, much of the way being under the welcome shade of walnut

and apricot trees. Extensive vineyards with prospect of many luscious bunches of grapes were seen near the village of Akpa. The view from Rorung was very fine, almost as grand as that from Pangi where we arrived on June 26th.

Six days could be spared for *shikar*, so on the following morning I went up the Kashang Gad, seeing splendid scenery on the way, and camped near the snow line at a place called 'Poki'. Simple names they have in this part of the world. Poki: Jangi: Pangi: Tangi! My *shikari* was a cunning looking fellow and I did not anticipate much sport. Next morning the previous nights tracks of a medium sized snow bear were found half a mile above camp. The tracks led over into the Ashang Gad, the animal being no doubt in search of food as wild carrots were not yet above ground in this valley. Week old tracks of a few burhel, and of a snow leopard, were also seen: but a prolonged reconnaissance made it fairly certain that, except perhaps for a stray bear, there was small hope of any *shikar*. The night was very cold, and the skies again quite clear. A second day's weary search for game proved it useless to remain any longer in this much talked of valley.

On June 30, a return was made to Pangi. The transport was somewhat varied, being made up of one bullock, two donkeys, two girls, one woman and one man. The man had the largest pair of hands I have ever seen on a human being.

At daybreak on July 1, I went up the valley north of Pangi to a bivouac at a camping place the people called Ting Ting. The elevation was about 13,000 feet. Three coolies, one of them a female lama—a nun I should call her—carried my kit. In the afternoon a walk to the snow line level enabled a large extent of country to be examined. Three burhel ewes were seen. The country was ideal for wild sheep, and the *shikari* said they were plentiful three years ago; but could not say why there were now so few. Doubtless he had himself been pretty busy reducing their numbers.

The hillsides were now becoming carpeted with flowers of many varieties. Purple iris, buttercups, daisies, pimpernels, forget-me-nots, and many kinds of small flowers of varied hues were all around me wherever I went. The clover was already ankle deep. I was sad at the thought of having to leave all this loveliness and wend my way back to the drab heat of the plains.

The vast panorama of mountains was very beautiful. Below, and on either side, lush grass land sloped down to the fir forests, which were parted by the silver streak of the rivulet hastening to add its waters to the mighty river far down the valley. The lofty range behind camp stood out boldly, its high turreted rocks and rough peaks forming fantastic shapes against the sky line. The camp fire burned brightly. Beyond the cavernous depths of the Sutlej cañyon uprose the mighty masses of many snow-capped giants. The sunset on the snows was magnificent, and just as the last lingering rose tint faded out of the sky, the great round orb of the full moon appeared over the edge of the mountain and flooded the scene with its soft luminance. The stars at that altitude appeared twice as large and brilliant as at lower elevations. All

was peaceful ; and the glamour of the wondrous scenery was upon me as my eyes closed in the dreamless slumber of a night in that perfect air. The memory of that evening of the first of July will not easily fade from my mind.

The dew that night was very heavy, so much so that blankets had to be dried in the sun until near eight o'clock. When we started down the hill the coolies gathered flowers and placed them in their hats, and the lady lama very prettily presented me with a flower for my own headgear ! I ascertained from her that any man, or woman, who wishes may become a lama. A married woman may also take the vow, but has to pay for the privilege. The nuns wear no ornaments, and their hair is cut short like a man's—not like a man of Kunawur—he wears his long ! Some of the nuns wear pyjama trousers but this one had the usual blanket dress. They are not exempt from 'begar', and among the lamas only the well-to-do men are excused this work. But now that the 'begar' system has been abolished perhaps the custom has been changed.

That same evening we got to Chini and found a great change in the appearance of the place. There was much vegetation : many flowers : wonderful blue and purple iris : huge white and yellow flowers, like daisies but three inches in diameter : splendid roses of pink and yellow in many shades ; nature was indeed busy putting forth her powers after the long winter sleep. The fruit was not yet ripe but cherries, gooseberries, asparagus, and several kinds of vegetables were brought to me from the Salvation Army garden and formed a very welcome change of diet.

Some remarks as to the country and its inhabitants may be here suitably inserted. Kunawar is the name of that portion of the State of Bashahr which lies beyond Wangtu. It is recorded that the people of Kunawar, alone among the neighbouring States, successfully resisted the Gurkha invasion of these hills. I saw nothing to enable me to suppose that the men of Kunawar were of such a warlike disposition. After passing Chini the influence of Buddhism becomes more and more marked : and Hinduism fades off gradually into the Buddhism of Tibet. Lamas in red and in yellow robes are to be seen, and the main walls and lamaist gateways are commonly met with.

The men of Kunawar wear caps of two kinds : a round one of wool, like a tam-o-shanter without bob on the top ; or a round cap, pork-pie shape, with edge turned up for about three parts of the way round. These are sometimes trimmed, as to the turned up part, with red cloth. Puggies are not worn. Shoes are mostly of woven wool with soles of rough leather. The goat's hair and grass shoes of the lower country are not seen. Clothes are of the excellent all-wool homespun material. It is made in lengths of about ten yards with a width of 21 inches. Such a piece, to a casual purchaser, was about Rs. 16. After weaving, the cloth is put into water and then well stamped on and kneaded by the women. They can be seen at this work practically every day at every village. The dress of the women is a blanket, worn as a coat and skirt, the spare stuff being folded behind and falling from the waist. This one piece costume is held up by a rope of goat's hair passed round

the waist. In some cases a woollen coat is also worn as a bodice. This style of dress has the appearance of a coat and skirt, and is well suited to the climate. The men's coats are very loosely made, with large sleeves, and are long, as a rule down to or below the knees. Trousers are of inferior, and more coarse, material than the coats. All the people carry a blanket and a goat's hair rope at all times. When not in use the blanket is rolled up round the waist. Much of the population emigrate in the winter to the lower valleys, taking the sheep and goats with them. The custom of the country is that the inferior when meeting with a superior always takes the downhill side of the path; and a low caste man, when speaking to a superior, has to cover his mouth with his hand to prevent his breath reaching the face of the man of higher status than himself!

The weather was now much more warm, so the fourteen miles to Urni on July 5th seemed long and tiring. 'Motee' was requisitioned to carry me some miles of the way—the first time he had been ridden since the march from Darangati to Sarahan—there being now no need to keep in training for the strenuous mountain work. When passing Rogi news was brought of a villager having that morning fallen off the 'jhula' by which the crossing of the Sutlej was effected. He was instantly swept away in the roaring flood, and can have lived but a few brief moments in those icy waters. Gangaram's reception of the news seemed to indicate that he felt he had not been quite so foolish as was thought at the time of his timorous crossing on the way to and from Rispa.

The valley opposite Rogi, access to which can be had either by way of Kilba or by this 'jhula' below Rogi, is said to provide lovely scenery, superior even to that of the Chini Valley. It is possible to travel by this route and so gain the valley of the Tons River: but it is a journey concerning which enquiry should be made. The pass to be crossed at the head of the valley would have an elevation of about 15,000 feet.

At Urni the house flies were very bad. They nearly ate me and my tea, and even after dark did not let me have dinner in peace. Flies are always a great plague at staging bungalow and near villages in the hills during the summer months. When crossing the river at Wangtu the increase in the volume of water was very marked. The noise of the rushing torrent was tremendous, it being difficult to make oneself heard. The valley was much changed in appearance, all being now green and luxuriant where formerly it was bare and rocky. Fortunately the day was cloudy or it would have been exceedingly hot in that narrow gorge. Many flowers were seen, among them a large white flower growing on a creeper. It had a most delicate, exotic, perfume.

The Forest Bungalow at Nichar was very shady and cool. It is beautifully situated, being enclosed by stately deodar trees, with an unhampered view of the rugged precipitous hills on the far side of the Valley.

The views on the way to Taranda and Sarahan were just wonderful beyond description. Everything was clear and fresh. Wild flowers, ferns, and bracken were in luxuriant profusion. Jewelled butterflies flitted in sun and shade and the distant

hill sides were shrouded in a lovely blue and purple haze. The road was now thronged by mules, ponies, and bullocks taking food supplies to the many hundreds of sawmen and wood cutters working in the forest. All the forests of Bashahr and Kunawar are under long lease to the Imperial Forest Department. A wise arrangement. Many of the animals met with were bleeding at the nostrils owing to the leeches which infest the lower hills at this season of the year. For the most part the beasts looked fat and well fed. The men with them were Bunjaras, but not recognizable as being of the same race as the much finer looking gipsies of the Deccan : those nomadic people whose bullocks formed the bulk of the transport for the armies of the campaigns of the Duke of Wellington.

Mules conveying grain carry two maunds and payment was made at Rs. 4-8-0 a maund, whereas the payment for the slower travelling bullocks, carrying the same weight, the same distance, was Rs. 3-10-0 a maund. A number of sheep and goats was also met with, each animal carrying his little sack of grain : some of them were going to Rupa and other villages of the Thamin Valley. There is a very busy grain trade to the higher valleys of this country, and the Hindustan-Tibet road, so far as it goes, must be of inestimable benefit to the people. The herds which take grains to Tibet return with wool and other produce of far-off Tibetan villages.

The sloping roofs of houses of Sarahan, Taranda and other villages are of wooden planks ; those of Gaora and the lower country are mostly of stone slabs. Cultivation is now more extensive and the valley broadens out, showing that one is leaving the higher mountains. Gaora Bungalow is at an elevation of 6,025 feet and commands a fine extensive view. The roar of the Sutlej could be but faintly heard. The descent from the comparatively cool temperature at Gaora to the heat of the valley at Rampur (3,280 feet) was somewhat unpleasant—even at that date, July 10. The icy flood of the Sutlej however made an appreciable difference when one was on its banks.

The P.W.D. Rest House is about three-quarters of a mile east of the town and is on the bank of the river, which runs at that point between perpendicular rocks of marble like formation. Passing through the wide clean streets of the town a number of public buildings are seen ; all well built and of pleasing design. The Guest House where I stayed is some little distance down stream from the town, and not far from the river bank, the rush of the waters being pleasant to the ear.

Rampur is the first place up the valley of the Sutlej where Lamaist buildings are to be seen. That at Rampur is a quite recent structure. It contains modern frescoes and a huge prayer wheel. In one of the Raja's garden houses are some lamaist frescoes. One fresco represents the treaty between Tibet and Bashahr concluded about A.D. 1650, when Bashahr was supported by the Moghul Emperor. In the middle of the picture is a figure apparently meant to represent the Moghul Emperor surrounded by his soldiers. A party of Bashahr people, distinguished by their

round black hats, are placed in front of the Moghul, while the embassy from Tibet is on the right side of the painting. The Tibetans were beaten by the Moghul Army and had to cede the Sutlej Valley down to the Wangtu bridge to the Bashahr State.

I was very glad to leave the steamy heat of the Rampur Valley and get to Bahli. An early start was made on account of the heat, and the three or four miles along the river accomplished before the rays of the sun got to the valley. The transport consisted of five mules and four coolies. One of the mules fell down the hill side but the damage to the load was fortunately not great, and the mule was unhurt. The gradient of the path was good, so the rise of 4,800 feet in seven miles was not felt. It was cloudy most of the day but the view of the snows over towards Bhabah nullah was very fine.

The Soongri and Bahli villagers were just then visited by a severe influenza epidemic so there was some difficulty in obtaining coolies for the march to Roru. Enhanced cooly rates had just been promulgated so the last two marches in Bashahr were slightly more expensive. The valley below Soongri was looking exquisite in its green summer garb. Each range of hills seen on the return journey at these lower elevations had a different shade of colour. As range after range recedes in the distance the colour changes from the bright green of the nearest to the dark purple of the distant hills. Those wonderful landscapes! The inhabitants of those beautiful hills should indeed be happy: doubtless they are, their faces show that, but their life is a hard one in many ways and counterbalances to some extent their climatic advantages.

Nearing Roru there was heavy rain, but the baggage did not get wet. I rode 'Motee' to keep the saddle dry. The Pabar River was in heavy flood; and stranded against a rock in mid-stream was the carcass of a huge buffalo, his very large horns conspicuous against the flood. Two king vultures were busy picking the bones of the unfortunate beast which had most probably fallen accidentally into the torrent and been swept away.

Between Arakot and Tewni the people complained to me of the damage done by sambur to their crops along the valley. There are many sambur in those forests. Tewni was hot: how hot I cannot say, as my thermometer had been left at Rampur. I managed to catch a few labeo and rohu, up to three pounds in weight, but no mahseer. The Tons is at all seasons an unsatisfactory mahseer river.

On July 19 we got to Kathyan—'Motee' and I! In former days I had always hoped for a panther at this place but had no luck. An entry in the visitors book under date October 22, 1918, recorded that an official on tour had shot a panther—7 feet 3 inches in length—from the west bedroom window; so perhaps 'Judy' and two other dogs taken by panthers close to the bungalow in June 1914, were avenged. It was near here, in the village of Chajjar, that I took a photograph of a dog which had been three times grabbed by a panther and survived the adventures. This was due to his heavy spiked collar. I now learnt that he had recently died a natural death. Between Kathyan and Mundali the Sianas

arranged two beats for bear, without result. This valley always holds bear in September when the crops are ripening.

That night at Mandali there was a heavy thunderstorm. The clouds grew black, lightning flashed along the crest of the hills, peals of thunder reverberated among the woods, small rivulets soon became rushing torrents. In the morning it was fine, but there was a thick mist almost all the way along the ridge. I got to Chakrata on the afternoon of July 21 in time to avoid a wetting. The coolies with the kit got caught in the evening storm: all the baggage was thoroughly soaked, and this the last march and the only occasion on which the baggage had got wet!

A few remarks as to supplies and arrangements may be welcomed. All usual supplies of the country, including Kerosine Oil, can be obtained at Roru, Sarahan, Nichar, Chini, and Rampur. At Chakrata *all* stores for the trip can be purchased; so nothing need be taken from the plains the same, of course, being the case if Simla is the place of departure.

A few tinned stores can be got from the Bungalow *khansamah* at Sarahan, which is the only place beyond Fagu, so far as I know, where there is a *khansamah*. Nearly all the Bungalows mentioned have cooking utensils and table kit which can be utilized by the traveller. The Divisional Forest Officer (Kotgarh P. O., Bashahr State) should be written to for permission to occupy the Rest Houses of the Forest Department, and the Executive Engineer, Hindustan-Tibet Road, Simla, should be addressed as to all other Bungalows on the road. 'Begar'—the obligatory carriage of baggage by the people of the country—has, I believe, been now abolished; but no doubt the necessary transport can be obtained. Pack mules could travel the whole distance, and could be hired at Chakrata or Simla. The Superintendent Simla Hill States should be written to as regards the proposed journey. The services of a State Tahsil chaprassi, if obtained, would smooth away many of the difficulties of travel.

And now comes the parting of the ways. Those among our readers who may be induced, by what I have so inadequately attempted to describe, to take a similar holiday amidst the grand scenery of the mighty Himalayas, will ever carry with them the memory of what they have seen: and, by the many photographs which they will doubtless secure will have such a record as will enable them, whenever they wish, to again 'take to the open road' and travel in vivid memory with the everchanging and magnificent scenery which was before them.

ON THE DOWN PLUMAGES OF SOME INDIAN BIRDS

BY

CLAUD B. TICEHURST, M.A., M.B.O.U. (Late Capt. R.A.M.C.)

(With a plate)

INTRODUCTION

In 1908 I described (*British Birds' Magazine*) the down plumages of some of the commoner British nidicolous birds, a study which up to that time had not been touched. During my stay in India I collected a fair number of notes on the down plumages of Indian birds, and these supplemented by notes and specimens sent me by Mr. H. Whistler, Mr. A. E. Jones, Col. Kinloch, Mr. Browne and others, form the subject of this paper. It is hoped that as the result of this preliminary paper and the appeal I have made through this Journal that much more material will be sent, so that a further instalment can be written on this neglected subject.

The young of birds when first hatched are either clothed with down and leave the nest almost at once (Nidifugous), or the down is comparatively scanty, until they can fly (Nidicolous). If nidifugous chicks have hitherto received but scant attention, nidicolous chicks have been entirely neglected by all naturalists.

Of how much value an exact knowledge of down plumages may be, cannot be at present said until we know these plumages in a good many genera and species, but it may be pointed out that the positions of some genera are so obscure that anything which may throw light on this subject would be welcome, and it is just at this age in a bird's life-history when lost affinities might be to well developed *rami* and close-set *radii*. So too the hair-like tips to down, revealed. Take for instance the *Crateropodidæ*—the ornithological 'waste paper basket' into which most genera that lacked an obvious niche were relegated—with its six sub-families and seventy-six genera (F.B.I. Ed. I, vol. i). Since the 'Fauna' was written, several genera have been removed entirely from this family on one ground or another, and I think it quite possible if one could but examine chicks of each of these seventy-six genera, one would find that some were in the wrong sub-family, or even did not belong to this family at all.

CHARACTER OF DOWN

In the newly hatched chick, down may be of two kinds:—

(1) *Prepennæ*, i.e., preceding and replaced by true feathers.

(2) *Preplumulæ*, i.e., preceding and replaced by adult 'under down'.

The two kinds are so similar in appearance that they can only be differentiated by examination of what is replacing them. (Fig. i *a* and *b*). Passerine and many other orders have only *prepennæ*; Cormorants have only *preplumulæ*; Hawks have both, the latter making up the bulk of the down.

Though the down of various birds varies so much in appearance, (compare a Pigeon and a Wader), it is only a matter of degree of development of the constituent elements. These consist of (1) *calamus*, very short, (2) *rachis*, ill-defined or absent, (3) *rami* long, slender, variable in number, (4) *radii*, on each *ramus*, well developed to practically absent (Fig. i *c*). Thus the hair like down of Pigeons is due to few *rami* without *radii*, (Fig. i *d*) and the softness and 'fluffiness' of the down of waders and compactness in sandgrouse is due to well developed *rami* and close-set *radii*. So too the hair-like tips to down, as in Herons, etc., is due to lack of *radii* at the extremities of the *rami*. The weak loose-looking down of *Passeres* is due to the sparseness of the down tufts composed of rather long slender *rami* with comparative short *radii* which do not reach to the tips of the *rami*.

Fig i






Fig ii



Down Tracks of a Passerine

- | | |
|------------------|------------|
| 1. Superciliary. | 5. Spinal. |
| 2. Supraorbital. | 6. Ulnar |
| 3. Occipital. | 7. Femoral |
| 4. Humeral | 8. Crural. |
| 9. Uropygial. | |
- (Ventral not shown)

Fig iii



Downy Chick (Dorsal view) of Platyoche senegalensis (semi-diagrammatic) showing pattern of down.

DISTRIBUTION OF DOWN

In *Passeres*, when down is present, it is always distributed in definite tracks, which of course, correspond to subsequent feather tracks, though each feather track is not necessarily preceded by down. In *British Birds' Mag.*, vol. ii, pp. 186-8, I named these tracks according to their anatomical positions, and this nomenclature, I see no reason to alter (Fig. ii).

- | | |
|--|------------------|
| 1. Outer-supraorbital or superciliary. | 6. Ulnar. |
| 2. Inner supraorbital or supraorbital. | 7. Femoral. |
| 3. Occipital. | 8. Ventral. |
| 4. Humeral. | 9. Crural. |
| 5. Spinal. | (10. Uropygial.) |

In many Nidifugous birds, down appears to cover the whole body; in certain other birds (Hérons, Hawks, Pigeons) down does not cover the entire body, but the tracks are not so simple as in *Passeres*, and their positions are not yet worked out.

It is noteworthy that, of these tracks, No. 1 is usually absent, and that 2, 3, 4, 5 usually present, 6 and 7 fairly often, 8 and 9 less often, and No. 10 only in one species.

Tongue Spots.—The meaning of tongue spots in newly hatched chicks is not clear; usually they consist of two oval spots at the base of the tongue, one on each side, and sometimes also at the tip. They appear to be well marked in Warblers and Larks only, and a trace in a few other species. They disappear as soon as the young are fledged.

REMARKS

1-3. *Corvidæ*. It is curious to find that there is no down in the Jungle Crow, British members of the genus *Corvus*, viz., *corax*, *corone*, *frugilegus*, *monedula*, have down present in varying amount; in *Garrulus glandarius* it is absent as also in *Pica pica*.

4-6. *Paridæ*. The distribution of down in *Parus major* is the same as in these two Indian species. As in *Æg. concinna* so in *Æg. caudata* down is absent.

8-14. *Turdoinæ*. It will be noted that the distribution in the two *Trochalopterum* is the same and in *Garrulax*, and *Pomatorhinus* is only slightly different. The total absence in *Argya*, *Pyctorhis* and *Stachyridopsis* is noteworthy—but specimens from more species and more genera are required before one can begin to generalize.

15. The sole *Brachypterygine* representative resembles *Trochalopterum* in down distribution, but here again more genera and species are required.

16-20. *Brachypodinæ*. Down seems to be absent in this group but one would like to see examples of more genera and species.

21-27. *Phylloscopus* and *Seicercus* in many ways are rather closely allied and also this is born out by the presence of, though reduced, down tracts. In species of *Sylvia*, *Acrocephalus*, *Hippolais*, *Cisticola*, *Prinia*, *Fanklinia* I have examined down is entirely absent. It is well marked however in *Locustella naevia*.

28-30. Down is absent also in the European *L. collurio* and *senator*.

32-34. Evidently quite a *sturnine* character are the large flanges of the gape. Much the same distribution of down also is seen in these genera. The crural and ventral tracks when only slightly marked as in *Acridotheres tristis* may be the subject of variation either individual or accidental; thus in other specimens of this species I have found no ventral or crural tracks but these were a nestful of chicks of varying ages and I think probably in this case loss of down tracks was accidental.

37-41.—The Fly-Catchers are split by some authors into many genera and lumped by others in a few genera; it is highly desirable therefore for more species of all the Indian genera to be examined to see if the down plumages can throw any light on how the Fly-Catchers should be grouped.

42-43. The similarity in the two *Saxicola* will be noted and the down distribution is the same in the English Stonechat and Whinchat.

ABBREVIATIONS.—Sh. = Short; F.L. = fairly long; L. = long; V.L. = very long; V.Sc. = very scanty; Sc. = scanty; F.P. = fairly plentiful; P. = plentiful; V.P. = very plentiful; X. = track present.

	Length	Amount	Colour	Superciliary	Supraorbital	Occipital	Humeral	Spinal	Ulnar	Femoral	Ventral	Cruial	Tongue Spots	Remarks
1. <i>Corvus c. intermedius</i> ...														
2. <i>Garrulus lanceolatus</i> ...														
3. <i>Nucifraga hemispila</i> ...	Sh.	Sc.	White		Absent								Nil	
4. <i>Parus c. kaschmiriensis</i> ...	"	"	Pale grey		X	X	X	X	X	X			"	
5. <i>Parus melanolophus</i> ...	"	"	Dark grey		X	X	X	X	X				"	
6. <i>Aegithaliscus c. iredalei</i> ...													"	
7. <i>Sitta c. castaneiventris</i> ...	Sh.	V.Sc.	Grey		Absent								"	
8. <i>Garrulax albogularis</i> ...	F.L.	F.P.	White		X	X	X	X	X	X			"	One tuft only.
9. <i>Trochalopteron l. lineatum</i> .	"	"	Dark grey		X	X	X	X	X	X			"	
10. <i>Trochalopteron e. erythrocephalum</i> .	"	"	Buff grey		X	X	X	X	X	X			"	
11. <i>Argya c. caudata</i> ...													"	
12. <i>Pomatorhinus e. erythro-</i> <i>genus.</i>	V.L.	F.P.	Dark grey		Absent	X	X	X	X	X	*		"	* Slight. Bill short and straight.
13. <i>Pycnorhis sinensis</i> ...													"	
14. <i>Stachyridopsis pyrrhops</i> ...													X	
15. <i>Leioptila c. pallida</i> ...													Nil	
16. " <i>p. psaroides</i> ...	L.	F.P.	Pale grey-brown		X	X	X	X	X	X			"	
17. <i>Pycnonotus h. hemorrhous</i> .					Absent								"	
18. <i>Pycnonotus h. intermedius</i>													"	
19. <i>Pycnonotus l. leucogenys</i> ...													"	
20. <i>Olocompsa e. fuscicaudata</i> .													"	
21. <i>Cisticola j. cursitans</i> ...													"	
22. <i>Franklinia buchanani</i> ...													"	

A faint trace only.

	Length	Amount	Colour	Superciliary	Supraorbital	Occipital	Humeral	Spinal	Ulnar	Femoral	Ventral	Cutural	Tongue spots	Remarks
55. <i>Monticola solitarius</i> ...	F.L.	F.P.	Dark grey		XXXX	XXXX	XXXX	XXXX	X	XX			Nil.	
56. <i>Monticola cinclorhyncha</i> ...	V.L.	P.	Rufous-buff										"	
57. <i>Myophonus temminckii</i> ...	F.L.	F.P.	Dark grey-brown	X	XXXX	XXXX	XXXX	XXXX	X	XX			X	
58. <i>Laiscopus c. whymperi</i> ...			Black										"	
59. <i>Prunella rubeculoides</i> ...	"	"	Grey-brown		XXX	XXX	XXX	XXX	XXX	XX	X		Nil.	Spec. rather old. Bill
60. <i>Ploceus p. philippinus</i> ...	"	"	White										"	bright pink, mouth
61. <i>Ploceus m. flaviceps</i> ...	"	"	"										"	bright orange,
62. <i>Uroloncha malabarica</i> ...	Sh.	V.Sc	"										"	flanges white.
63. <i>Uroloncha p. punctulata</i> ...	"	Sc.	"										"	
64. <i>Serinus pusillus</i> ...	"	F.P.	Pale grey		X	X	XX	XX	X	X	*		Nil.	
65. <i>Gymnorhis x. xanthocolis</i>													"	* Slight.
66. <i>Passer d. indicus</i> ...													"	
67. <i>Passer pyrrhonotus</i> ...													"	
68. <i>Montifringilla n. adamsi</i> ...	F.L.	Sc.	Grey-white		XXX	XXX	XXX	XXX	XX	X		X	"	* Slight.
69. <i>Emberiza c. stracheyi</i> ...	"	F.P.	Dark grey		XXX	XXX	XXX	XXX					"	
70. <i>Riparia chinensis</i> ...	"	"	White										"	
71. <i>Hirundo d. erythropygia</i> ...	"	Sc.	Pale grey		XXX	XXX	XXX	XXX					"	
72. <i>Hirundo r. javanica</i> ...	Sh.	P.	Pale brown		XXX	XXX	XXX	XXX		X	X		"	
73. <i>Hirundo s. filifera</i> ...	"	Sc.	Dark grey		XXX	XXX	XXX	XXX		X	X		"	
74. <i>Motacilla c. melanope</i> ...	L.	P.	Dirty white		XXX	XXX	XXX	XXX	XX	X			"	
75. <i>Motacilla c. calcarata</i> ...	F.L.	F.P.	Yellow white		XXX	XXX	XXX	XXX	XX				Tree	
76. <i>Anthus s. jerdoni</i> ...	L.	P.	Pale grey		XXX	XXX	XXX	XXX	XX	X			Nil.	* Scanty.
77. <i>Anthus roseatus</i> ...	F.L.	F.P.	Grey	XXX	XXX	XXX	XXX	XXX	XX	X	X	X	"	* Scanty.
78. <i>Anthus rufulus</i> ...	"	P.	Pale grey		XXX	XXX	XXX	XXX	XX	X	X	X	"	

45-50. The uniformity of down distribution in this group is interesting; the presence or absence of the superciliary track is a feature on which not too much stress can be laid as I know in some species it varies in individuals. *Ph. phænicurus* has the same distribution as that of *phænicuroides* and *Erithacus rubecula* is the same as in *Cyanosylvia suecica*—a near ally. The colour of most passerine down is 'neutral'—a grey, brown, black or white—a distinct red-brown as in *Rhyacornis* is a most unusual colour.

51-52. Is it certain that *Thamnobia* and *Copsychus* are *Ruticilline* or even belong to the *Turdidæ*? If so they are most aberrant. They are the only species examined among this family which lack down; their juvenile plumage has not the well marked spotted appearance of Redstarts or Thrushes. They are most remarkable in having when hatched a black skin, a condition I have not seen in any other of the *Turdidæ*; moreover, *Copsychus* at all events, has a complete post-juvenile moult which none of the *Turdidæ*, so far as I know, have.

53-54. *T. merula*, *philomelos*, *pilaris* and *viscivorus* all have the same down distribution as these two Thrushes. The absence of the ulnar track in *Monticola* may be a characteristic as opposed to *Turdus*.

58-59. The down in *P. modularis* is the same as in these. The mouth colours in *L. collaris* is remarkable.

60-63. The great reduction in tracks in *Uroloncha* (*Viduinæ*) is noteworthy as compared with the *Ploceinæ*; as also are the mouth colours of *Uroloncha*. The palate and floor of the mouth is yellow (*malabarica*) whitish (*punctulata*) outlined with black, and the base and tip of the tongue is also black. Similar mouths are found in African *Viduinæ* (Cf. Ibis 1916, p. 275).

65-67. The total absence of down in *Passer* and *Gymnorhis* is curious as other finches are well supplied with down. The Sparrows might well stand as a sub-family *Passerinæ* since they have also other peculiarities.

69. Buntings would seem always to have 'full' track; the same distribution is found in *E. schæinclus* and *citrinella*.

74-75. In the Wagtails the superciliary, femoral and crural tracks appear to be variable in the different species and in individuals; thus. *M. yarrelli* has these tracks present in addition to the others. In a nest-full of chicks of *M. flava thunbergi* (Norway) which I examined, the superciliary was absent and the femoral and crural were present, though scanty, in some but not in others, the distribution otherwise was as in *calcarata*.

76-78. The distribution in *Anthus* is remarkably uniform; the English *Anthus trivialis*, *pratensis* and *obscurus* have also the same tracks present.

79-85. Here again in the *Alaudidæ* there is a remarkable uniformity; it is interesting to note that *Alæmon*, which in many ways has diverged from the 'typical' Lark, retains the typical Alaudine down distribution, which surely suggests that this character is an archaic one and should be of value; the long legs and bill and aberrant nesting habits are adaptations. The down in *Alæmon* is remarkably long and plentiful, and the chicks are said to leave the nest and run before they can fly, the only instance I can recall of a nidicolous bird tending to become nidifugous. It would be interesting to know on what day the young do leave the nest.

The down in *Alauda gulgula* (and in *arvensis*) is grizzled, that is, of two colours; in all the other species of genera examined it is unicolorous.

90-103. Heelpads, tuberculated or spiny pads under the tarso-metatarsal joint, must I think be associated with 'hole-breeding', and help to balance the body on the metatarsus and feet; yet it is absent in *Merops* and *Upupa*. These pads disappear after the young are fledged. Note the presence of slight pads in *Temenuchus pagodarum* which frequently nests in old Wood-pecker holes, and the absence in the other Mynahs examined.

(For further notes on 'heelpads', see Chasen.)

104. *Caprimulgus e. unwini*. General colour pale isabelline, paler than in *europæus*; dark spots on head, wings and thighs black not brown as in *europæus*. Almost completely clothed with down.

105. *Caprimulgus mahrattensis*. General colour whitish-isabelline and so rather paler than *unwini*; dark marks less pronounced, but in the same positions, than in *unwini*.

106. *Glaucidium cuculoides*. Clothed with short white down all over when first hatched.

107. *Bubo coromandus*. No tracks. Almost entirely covered with short, pure white down when first hatched.

108. *Otogyps calvus*. Crown from level of eyes and nape clothed with short, close grey-white down, centre of chin and throat sparsely tufted same; crop with thicker buff tinged down; sides of breast and flanks longer pale buff down; rest of upper parts, grey-white. Bare skin of face and upper parts of head and cere pale bluish-lead colour in contrast to skin of throat, chin and neck, which is pinkish-lead. A bare ring $1\frac{1}{4}$ in. wide round neck from nape bare with longitudinal flaps of loose skin, a prominent one on each side being probably the origin of the adult wattles. (*H. Whistler*).

109. *Pseudogyps bengalensis*. Down covers whole body except a track from base of neck broadening to vent. Down of head and neck shorter than the rest. Mouse-grey in colour except darker patches to the uropygium and along ulnar border. Cere and base patch round eye blue-lead.

110. *Milvus m. govinda*. Plentiful. Long. No tongue spots; cere and gape livid grey; general colour whitish, tinged with buff; eyelids, dark mark in front of and behind eye, along radius of wing and down both sides of spine coalescing in the middle, sooty brown.

111. *Falco jugger*. Fairly long, dull white. Whole body covered except in definite places (=Apterion). Whole head covered, and down continuous thence down hind neck to spinal track, which spreads out over back of thighs and is continuous with lateral ventral tracks. Between the spinal and humeral tracks, an apterion with sparse short tufts of down. From the chin run two ventral neck tracks, separated by apteria from the dorsal neck track and from each other, which at the top of the sternum, divide into lateral and median sternal tracks, the apteria between them having scanty, short tufts; apterion in centre of abdomen, rest covered with sparse down continuous with dorsal track laterally. Upper surface of wing except *manus*, clothed with long down, under surface, four rows of short down. Thumb with distinct claw.

112. *Butastur teesa*. Prepennæ and preplumulæ both present.

Prepennæ, fairly long and plentiful over the whole body; white above, tinged with isabelline at tips; white below.

Preplumulæ; short, tufted, white.

Bill bluish, black at tip; cere and legs yellow, claws slate; iris dark brown.

113. *Aquila rapax vindhiana*. Prepennæ and preplumulæ present—white, sparse on cere and round orbit otherwise completely covering body. Preplumulæ makes up the bulk of down.

Bill, dull black, cere gape and legs yellow. Iris bluish-brown.

114. *Crocopus*. Long, plentiful, hair-like. Whitish.

115. *Streptopelia o. meena*. Long, plentiful, hair-like, yellowish-white. Whole body covered except sides of head, centre of abdomen and breast, chin and upper throat. Distribution and character is as in *Columbæ*.

116. *Streptopelia suratensis*. Long, plentiful, hair-like, rather darker than in *meena*.

117. *Pterocles senegalensis*. General colour above golden brown with black tips, and creamy-white. A more definite pattern than in *alchata*. A whitish line from base of bill passes over the head as a coronal streak (broken on forehead) to hind neck and then to middle of the back where it divides, making the top of a 'figure of eight'. From base of bill another white streak passes in front of eyes, and is lost in the ear coverts. Well marked 'figure of eight' on lower back outlined by black and enclosing brown down. Rest of crown pale ginger brown dotted with black; underparts buffish white, darkest on the pectoral region; bill and bare space round eyes, lead-blue; iris hazel; feet dusky-flesh, nails white. (See Fig. iii).

118. *Pterocles senegallus*. General colour above very pale sandy yellow, much paler than in *senegalensis* or *alchata*, with black tips to the down tufts, which on the head are not well marked except above the eyes; area round eyes pale ginger-brown. Underparts very pale isabelline; down pattern less variegated than in *senegalensis* and *alchata*.

119. *Pterocles alchatus caudacutus*. General colour above mottled very pale buffy-white, pale-ginger with black tips. No definite pattern, but supercilia and coronal streak whitish. The brown down here and there has long black, hair-like tips, while on the thighs, the whitish down has similar white hairs. Underparts, whitish to buffish-white.

120. *Gallus g. gallus*. From base of bill through eyes and down sides of neck, a black line; over this a broad line of pale buffish-brown; from base of comb down crown, neck, and dorsum to tail, a broad black line outlined on dorsum by a creamy streak on each side; wings chestnut; breast and thighs pale buffish-brown; rest of underparts creamy-white.

121. *Tragopan satyra*. Side of head and supercilium dirty buffish-white, dark tipped on ear-coverts; crown from bill ferruginous; upper parts ferruginous-brown with darker tips and paler bases, but no distinct pattern; tail ferruginous; chin and throat buffish-white; rest of underparts same, with faint dark tips and tinged with rust on sides and on thighs.

122. *Perdicula asiatica*. From base of bill over crown, down centre of back to tail, a broad chestnut band edged on each side with dark brown; outside this from base of bill a broad ochraceous supra-orbital stripe reaching nape and surmounting a narrow dark brown superciliary stripe; ear-coverts and short moustachial streak dark brown. Rest of upper parts dappled rusty brown and light; underparts ochraceous-grey.

123. *Francolinus francolinus*. Upper parts pale creamy-buff, from base of bill down centre of crown a broad chocolate-brown line, a short narrow line behind eyes, and a long, narrow one over the eyes and one from base of bill below the eyes, brown; three broad parallel brown lines on dorsum, and a broken one over thighs; underparts cream white; wings creamy-buff, mottled with brown.

N.B.—The description is taken from a chick of *arabistanicus*, the markings on the hill race (*melanonotus*) are similar, but the whole coloration is so much darker that one would hardly suppose the two to belong to the same species. It would be very interesting to see what the chick of the Plain bird (*asiæ*) is like; likewise the chicks of the various races of *pondicerianus*.

124. *Alectoris græca chukar*. A broad band from bill over crown to nape chestnut outlined in dark brown; from base of bill, a broad creamy white superciliary streak surmounting a narrow dark brown one; on the dorsum a broad median chestnut band and on each side of it a narrow creamy white streak; wings and back of thighs girdled brown and cream; underparts creamy grey, a short dark brown moustachial streak from base of bill.

125. *Tetraogallus himalayensis*. Upper parts stone-grey richly variegated, with black tips and subterminal pale-buff markings, making no definite pattern. Head, very pale creamy buff, tipped with black, and having four black lines from base of bill (1) along the side of the crown, (2) over the eye, (3) beneath the eye and (4) as moustachial streak, all more or less blending behind the eye and ear coverts. Underparts white, greyish-white on breast and flanks. Feet and legs, horny-yellow, bill black, iris olive-brown.

126. *Esacus recurvirostris*. Covered with close 'furry' down. Narrow frontal band and line to eye, under eye and thence to occiput black; broken black lines on each side of spine, meeting at level of thighs and sending transverse bar to base of wings, black; short black line over thighs; rest of upper parts 'pepper and salt' colour, finely dotted with black. Underparts white.

127. *Cursorius coromandelicus*. Whole head, wings and upper-parts mottled ginger, creamy-white and black with no definite pattern. Underparts, creamy-white, pectoral region ginger; legs greyish,

128. *Glareola p. pratincola*. Head, wings and back grizzled dark brown and pale buff with median dark line down crown; underparts pale buff.

129. *Glareola lactea*. Whole of upper parts finely mottled greyish-white, dusky, and pale buff, narrow black line behind eyes; underparts white.

130. *Charadrius a. alexandrinus*. Forehead and carpus white; black line on forearm and behind each eye; white hind-collar; crown and rest of upperparts mottled black and buffish-white; underparts white.

131. *Charadrius dubius jerdoni*. Forehead, carpus, edge of ulnar and collar of hind neck white; circle round crown from eye to eye and above white collar, black; lines on thighs and forearm, tail, black; rest of upperparts grizzled black, white and ginger points.

132. *Charadrius mongolus atrifrons*. Upper parts deeper buff than in *alexandrinus*, mottled with black on crown; wings and whole of back, tips of wings and collar of neck grey-buff; narrow, short black lines on lores, behind eyes and on the ear coverts. Underparts white, grey-bluff at the sides; legs and feet plumbeous.

133. *Himantopus himantopus*. Head and upper parts mottled pale buff and black; black spinal line; black line from base of bill to eye, and on ulnar. Round eye and whole of underparts pale buff.

134. *Dromas ardeola*. Entirely covered with long, very pale smoke-grey down, darker on forehead and round eyes. Legs slate-grey.

(N.B.—The character of the down is more *Larine* than *Limicoline*, and lacks the mottling and lines which the latter nearly always shew.)

135. *Ibidorhynchus struthersi*. Upper parts mottled white, buff and black, underparts white. Bill straight. (N.B.—This is such a very poor specimen that details of pattern, etc., cannot be made out.)

136. *Lobivanellus indicus*. Upper parts grey brown mottled with black; behind eyes and round occiput a broad black collar, under this from below eyes, a broad white collar. Pectoral band black; chin and rest of underparts white, tinged with buff on belly and flanks.

137. *Scolopax rusticola*. From base of bill to eye, over and behind it, a narrow, dark brown line, from centre of base of bill to centre of crown, a dark brown line; between the two lines, creamy-yellow; rest of crown pale chestnut; side of head variegated, chestnut and creamy-yellow, as are the wings and rest of upper parts, shewing a broken spinal brown streak; underparts creamy-yellow tinged with ferruginous on throat and thighs.

138. *Larus gelastes*. Head, sides of head, upper parts, wings and thighs, evenly mottled with about equal areas of black and white alternating; underparts more white but black spots across lower throat and on sides of breast and flanks.

139. *Sterna caspia*. Like *nilotica*, but much fewer spots on crown and dorsum, and none behind the eyes.

140. *Sterna nilotica*. Upper parts dirty grey-buff with black spots on crown, behind eyes and on the dorsum where they tend to form stripes; carpus white, throat dusky to blackish, rest of underparts, nearly white.

141. *Sterna b. bengalensis*. Upper parts almost white with black spots on crown and ulnar, and tending to run into lines on the back. Underparts pure white.

142. *Sterna albifrons saundersi*. General colour more buffish than *melanogaster*, and more mottled with black; warmer buff than *a. albifrons*; underparts nearly white. Legs flesh.

143. *Sterna melanogaster*. Upper parts cream; broken line of black from base of bill on each side of crown and down dorsum; rest of upper parts spotted with black except wings; underparts cream-coloured.

144. *Egretta garzetta*. Long, plentiful, nearly white.

145. *Bubulcus coromandus*. Whole crown white, down 20 mm. long; short white down on ear-coverts and sides of neck; dorsum grey-white, underparts white. The down is distributed in definite tracks, but further specimens are required to make out the details. Down longer in some tracks than others. Iris, pale straw, facial skin, yellowish-green.

146. *Ardeola grayi*. Less long than in *garzetta*, nape and dorsum ruddy-buff, crown white or buffish-white, rest white.

147. *Megalornis antigone*. Entirely covered. Head, neck and throat pale isabelline; lower neck sides of breast and belly, vent and thighs darker isabelline; centre of breast and belly white from base of hind neck two broad lines of dark brown, enclosing a paler brown median stripe, run to tail; large buffish-white spot opposite base of both wings; wings cinnamon-brown, tip buffish-white.

148. *Megalornis leucogeranus*. Entirely covered. Underparts dirty buff, belly white. Head and neck dark isabelline; creamy streak on each scapular, rest of upper parts pale chestnut, darker on mid-dorsum (captivity bred).

In a former number of the Journal I made an appeal to members to supply me with material and I again appeal to them. My opportunities of getting such myself are *nil*,—I must rely on those still in India. I hope I have shewn that, apart from the fact that the down plumages of many Indian birds are still quite unknown, there may be learnt from them some information of important character and taxonomic value.

In order to help in this research members *need not be skilled ornithologists*; all that is required is that a chick fairly recently hatched (or even an egg on hatching point) shall be *labelled with the name of the species, wrapt in linen, and put into spirit or weak formalin*; at the end of the nesting season the bottle or tin may be sent to The Honorary Secretary, Bombay Natural History Society, 6 Apollo Street, Bombay, who has kindly offered to see to the despatching to England.

BOMBAY NATURAL HISTORY SOCIETY'S
MAMMAL SURVEY OF INDIA, BURMA AND CEYLON

REPORT No. 40

BY MRS. HELEN M. LINDSAY

(*With Field Notes by the Collectors*)

COLLECTION	No. 40.
LOCALITY	Gwalior State.
DATE	June-December 1922.
COLLECTED BY	Messrs. C. Primrose and J. R. O'Brien.

This collection is the continuation of that commenced in July 1914 by Major E. W. Mayor to represent the fauna of the Gwalior State in Central India. This was interrupted by the outbreak of war, but the few specimens then obtained were described in Report No. 21.

Report No. 7, also deals with specimens from Central India and describes in detail the nature of the country in that area. Gwalior State in the main shows the same features as those described for the districts of Damoh and Saugor in that report.

Gwalior State extends from 22° 10'–27° N. Latitude and from 74° 38'–79° 8' Longitude. It forms three natural divisions :—

1. PLAIN represented in this collection by the localities Sheopur, Sabalgarh and Morar visited by Mr. Primrose and Guna visited by Mr. O'Brien. Its rainfall is 40 inches. The hot season is oppressive and the cold in winter severe.

2. PLATEAU represented by Agar Malwa whence at 1,500' specimens were obtained by Mr. O'Brien. Its rainfall is 30 inches. The climate is comparatively equable, being free from extremes.

3. HILLS. This area is in the extreme south-west of the State but does not seem to have been visited by the collectors. Anjhera Zilla is its centre. It is covered with thick jungle.

The geological formation of Gwalior State resembles that described for Damoh and Saugor districts in Report No. 7.

Gwalior proper is largely occupied by the Vindhyan series with four principal ranges capped by massive beds of sandstone which form the chief mineral product of the State.

In the south sections the formation is also sandstone, but with fossil beds in a series of calcareous strata along with extensive outcrops of gneissose rocks. The rest of the area is Deccan trap. The best soil is found in the Malwa part of the State, but the part round Bhind is covered with alluvial soil of fair fertility. Guna is the chief centre for the sale of grain. The principal crop of the State is Jowar (Great Millet).

In the north the vegetation in waste tracts consists largely of deciduous trees and shrubs, many of which flower when leafless during the hot season. Further south the low hills are covered with jungle, and Bamboos abound in places. The typical forest of the Central Indian Highlands is found in the extreme south.

As noted in Report No. 7, the fauna is of the same general type as that of Nimar and the Berars. But in the Gwalior collection there are no deer, monkeys or otters and only one species of Hare and one Antelope were obtained. Again the absence of Tree Shrews (*Anathana*) is to be recorded.

The present collection numbers 898 specimens belonging to 27 genera.

(1) *PTEROPUS GIGANTEUS*, Bruenn
The Common Flying Fox
(Synonymy in No. 2.)

Morar, ♂ 2; Bhind, ♂ 2; Sabalgarh, ♂ 8, ♀ 4; Sheopore, ♂ 1, ♀ 1; Guna, 1,564', ♂ 1, ♀ 4; Chachora, 1,681', ♂ 7; Agar Malwa, ♂ 2, ♀ 1.

- (2) HIPPOSIDEROS FULVUS PALLIDUS, K. And.
The Bicoloured Leaf-nosed Bat
 (Synonymy in No. 2.)

Sheopore, ♂ 1, ♀ 1.

- (3) LYRODERMA LYRA, Geoff.
The Indian Vampire Bat
 (Synonymy in No. 1.)

Bhind, ♂ 6, ♀ 6; Sheopore, ♀ 1; Chachora Fort, 1,681', ♂ 30, ♀ 16; Agar Malwa, ♂ 3, ♀ 1.

- (4) PIPISTRELLUS MIMUS, Wroug.
The Southern Dwarf Pipistrelle
 (Synonymy in No. 1.)

Guna, 1,564', ♂ 1.

- (5) PIPISTRELLUS BABU, Thos.
The Babu Pipistrelle
 (Synonymy in No. 26.)

Sabalgarh, ♀ 1.

- (6) SCOTOPHILUS KUHLI, Leach.
The Common Yellow Bat
 (Synonymy in No. 1.)

Guna, 1,564', ♂ 9, ♀ 2; Agar Malwa, ♂ 3.

- (7) SCOTOPHILUS WROUGHTONI, Thos.
Wroughton's Bat
 (Synonymy in No. 1.)

Sabalgarh, ♀ 1.

- (8) TAPHOZOUS LONGIMANUS, Hardw.
The Long-armed Sheath-tailed Bat
 (Synonymy in No. 26.)

Binaganj, 1376', ♀ 1.

This is a very large specimen H.B. 104 mm., T. 26, F.A. 61.

- (9) LIPONYCTERIS KACHHENSIS, Dobs.
The Cutch Sheath-tailed Bat
 (Synonymy in No. 3.)

Morar, ♂ 8; Sabalgarh, ♀ 1.

- (10) RHINOPOMA HARDWICKEI, Gray.
The Lesser Indian Mouse-tailed Bat
 (Synonymy in No. 3.)

Morar, ♂ 1, ♀ 2.

- (11) RHINOPOMA KINNEARI, Wroug.
The Greater Indian Mouse-tailed Bat
 (Synonymy in No. 1.)

Morar, ♂ 1, ♀ 4.

- (12) PACHYURA, sp.
Shrews

Morar, ♂ 9, ♀ 5; Bind, ♂ 3; Guna, ♂ 5, ♀ 29; Rothiai, ♂ 6, ♀ 1; Binaganj, ♂ 9, ♀ 4.

Awaiting further study.

- (13) FELIS AFFINIS, Gray and Hardw.

The Jungle Cat

(Synonymy in No. 1.)

Morar, ♀ 1; Binaganj, ♀ 1; Rothiai, ♂, 1; Agar Malwa, ♀ 2.

- (14) PARADOXURUS NIGER, Desm.

The Indian Toddy Cat

(Synonymy in No. 5.)

Guna, ♂ 1, ♀ 1; Binaganj, ♂ 1, ♀ 3; Chachora Fort, ♀ 1; Agar Malwa, ♀ 1.

- (15) HERPESTES EDWARDSI FERRUGINEUS, Blanf.

Blanford's Indian Mongoose.

(Synonymy in No. 24.)

Morar, ♂ 1; Sabalgarh, ♀ 1; Sheopore, ♀ 1.

- (16) HERPESTES EDWARDSI, Geoff.

The Common Bengal Mongoose

(Synonymy in No. 1, under M. Mungo.)

Morar, ♀ 1; Guna, ♂ 1, ♀ 2; Chachora Fort, ♀ 1; Agar Malwa, ♂ 4, ♀ 3.

- (17) CANIS INDICUS KOLA, Wrough.

The Common Jackal

(Synonymy in No. 1.)

Binaganj, ♀ 2; Agar Malwa, ♂ 2, ♀ 2.

- (18) VULPES BENGALENSIS, Shaw.

The Indian Fox

(Synonymy in No. 1.)

Morar, ♂ 1; Binaganj, ♂ 2, ♀ 1; Agar Malwa, ♂ 2, ♀ 1.

- (19) FUNAMBULUS PENNATI, Wrough.

The Common Five-striped Squirrel

(Synonymy in No. 1.)

Morar, ♂ 32 (1 juv), ♀ 13; Bhind, ♂ 10, ♀ 2; Guna, ♂ 3, ♀ 4; Rothiai, ♂ 3, ♀ 3; Khatia, ♂ 2, ♀ 3; Binaganj, ♂ 7, ♀ 5; Agar Malwa, ♂ 13, ♀ 4.

- (20) TATERA INDICA, Hardw.

The Indian Gerbil

(Synonymy in No. 1.)

Morar, ♂ 19, ♀ 26; Bhind, ♂ 6, ♀ 2; Guna, ♂ 3, ♀ 15 (9 juv.) Rothiai. ♀ 1; Khatia, ♂ 5, juv. ♀ 3; Binaganj, ♂ 4, ♀ 6; Agar Malwa, ♂ 38, ♀ 20.

- (21) GUNOMYS BENGALENSIS, Gray and Hardw.

The Bengal Mole-Rat

(Synonymy in No. 15.)

Morar, ♂ 2, ♀ 2; Guna, ♂ 2, ♀ 4 (2 juv.); Rothiai, ♀ 7 (4 juv.); Agar Malwa, ♂ 6; Binaganj, ♂ 6, (4 juv.), ♀ 1 juv.

- (22) RATTUS RATTUS RUFESCENS, Gray.

The Common Indian Rat.

(Synonymy in No. 1.)

Some of the specimens are coloured like *R. rufescens*; Others like *R. alexandrinus*, but all are parasitic.

Morar, ♂ 1, ♀ 1; Bhind, ♂ 4, ♀ 2; Sabalgarh, ♀ 2; Guna, ♂ 3, ♀ 6; Binaganj, ♂ 9, ♀ 6 (2 juv.); Agar Malwa, ♂ 16, ♀ 2.

(23) MILLARDIA MELTADA, Gray.

The Soft-furred Field Rat

(Synonymy in No. 1.)

Morar, ♂ 1; Bhind, ♂ 2; Guna, ♂ 7, ♀ 15; Khatia, ♀ 1; Rothiai, ♀ 1;
Binaganj, ♂ 3, ♀ 2; Agar Malwa, ♂ 10, ♀ 4.

(24) LEGGADILLA SADHU, Wrough.

The Ashy Spinpy Mouse

(Synonymy in No. 3.)

Bhind, ♂ 4, ♀ 4; In al. 6; Agar Malwa, ♂ 3, ♀ 2.

(25) MUS HOMOURUS, Hodge.

The Himalayan House Mouse

(Synonymy in No. 15.)

Guna, ♂ 3, ♀ 8; Rothioi, ♂ 3, ♀ 3; Binaganj, ♂ 3, ♀ 2.
The colour of the belly is whitish.

(26) MUS URBANUS, Hodgs.

The Nepal House Mouse

(Synonymy in No. 15)

Binaganj, ♂ 30, ♀ 13; Bhind, ♀ 1, In al. 2; Sabalgarh, ♂ 2.
Sheopore, ♀ 1; Agar Malwa, ♂ 18, ♀ 10; Chachora Fort, ♂ 9 ♀ 1.
Belly without sharp change of colour.

(27) LEGGADA BOODUGA, Gray.

The Southern Field Mouse

(Synonymy in No. 1.)

Morar, ♂ 8, ♀ 5; Bhind, In al. 3; Agar Malwa, ♂ 5.

(28) VANDELEURIA OLERACEA, Benn.

The Deccan Tree Mouse

(Synonymy in No. 2.)

Morar, ♂ 1, ♀ 1; Guna, ♀ 4, Rothiai, ♂ 1, ♀ 1; Khatia, ♂ 2, ♀ 1.
Binaganj, ♂ 3; Agar Malwa, ♂ 11, ♀ 3.

(29) GOLUNDA ELLIOTI, Gray.

The Indian Bush Rat

(Synonymy in No. 1.)

Morar, ♂ 2, ♀ 3; Binaganj, ♂ 5, ♀ 4; Agar Malwa, ♂ 26, ♀ 26.

(30) LEPUS RUFICAUDATUS, Geoff.

The Common Bengal Hare

(Synonymy in No. 15.)

Morar, ♂ 3, ♀ 1; Guna, ♀ 1; Binaganj, ♂ 1, and 5 juv.

(31) ANTILOPE CERVICAPRA, L.

The Black Buck

(Synonymy in No. 1.)

Guna, 1,564', ♀ 1.

(32) ACANTHION LEUCURUS, Sykes.

The Common Indian Porcupine

(Synonymy in No. 1.)

Agar Malwa, ♂ 2, ♀ 2.

REPORT No. 41, ASSAM AND MISHMI HILLS

BY MARTIN A. C. HINTON AND HELEN M. LINDSAY

(Published by permission of the Trustees of the British Museum)

COLLECTION	No. 41.
LOCALITY	Assam and Mishmi Hills.
DATE	October 1919–July 1920.
COLLECTED BY	Mr. H. W. Wells.

The collection made by Mr. Wells in Assam and the Mishmi Hills aids considerably in the completion of the survey of the mammals of N. E. India, filling, as it does, the gap between the collections described in Reports Nos. 23, 26, 27, 28, 37, dealing with Sikkim, Darjeeling, Bhutan, Duars, Kalimpong and Nepal respectively on the west side, and Report No. 38 describing collection of specimens from the Naga Hills on the east.

The area visited by Mr. Wells lies between 25° N. and 29° N. and 90° E. and 97° E, thus practically embracing the whole province of Assam—53,000 sq. miles—excluding the small part comprising Manipur and the Lushai Hills—8,500 sq. miles. According to the *Gazetteer*, from which much of the information about the country has been taken, Assam falls into three natural divisions:—

(1) **SURMA VALLEY.**—A flat plain about 125 miles long by 60 miles wide shut in on three sides by ranges of hills. The western end of the valley is very low, only about 22·7 ft. above sea-level, and in the rains the West Sylhet part is completely under water. Cachar and East Sylhet are higher. Here 'blue hills bound the view on almost every side, the villages are buried in 'groves of slender palms, feathery bamboos and broad-leaved plantains, and 'even in the dry season the country looks fresh and green.' Little or no forest exists in Sylhet, but there are extensive Reserves in the South and East Cachar District.

(2) **BRAHMAPUTRA VALLEY.**—An alluvial plain about 450 miles in length with an average breadth of about 50 miles, this is also shut in by hills on every side except the west. As a whole, it is a plain of fairly uniform breadth except in the centre where the Mikir Hills project from the main mass of the Assam Range. Outcrops of gneissic rock above the alluvium between Tezpur and Dhubri, and ranges of low hills in the central part of Goalpara District break the even level of the plain. On either side, the Brahmaputra is edged by stretches of marsh land, beyond which comes a belt of rice fields and fruit trees. Beyond that belt near the hills, grassy plains and forests are interspersed with prosperous tea gardens. There are still large areas not cultivated, consisting of huge wastes where heavy forest alternates with Savannah land, covered with a dense growth of reeds and elephant grass, 14'–16' high. In these wastes are found buffalo, rhinoceros, and swamp and hog deer.

(3) **THE ASSAM RANGE.**—Mountains separating these two valleys and projecting at right angles from the Burmese system. It lies almost due east and west. At its western end it attains a height of more than 4,600 ft. in the peak of Nokrek, above the station of Tura, the head-quarters of the Garo Hills District. Here the hills are broken up into sharply serrated ridges and deep valleys, all alike covered with forest. Near Shillong in the Khasi Hills, 6,450 ft. is the highest peak in a tableland about 6,000 ft. high with wide stretches of oak and pine. The highest point in the province of Assam proper is Japvo, 9,890 ft. in the Naga Hills, where the hills are all of the serrated type, the sides clothed with forest or fallowing jhums.

The MISHMI HILLS shutting in the eastern end of the Brahmaputra Valley seem to be a continuation of the Burmese axials. There are some peaks of 1,500 ft. but little is known about this tract of the country.

The basis of the whole Assam Range is a gneissic rock. Sandstones and conglomerates of the Cretaceous System are superimposed upon the gneiss, and are themselves in turn overlaid by limestone and sandstone of the Nummulitic age, especially at the western end of the range. Thus the late Mrs. Jackson in her note on the Garo Hills speaks of the fossil beds and the limestone caves to be found there. It appears, to quote the *Gazetteer*, little is known of the eastern extremity of the Assam Range. Apparently Upper Tertiary sandstones

'are succeeded by a series of hard sandstones, slates and shales with quartzose beds, while still further east serpentine dykes, indentical in composition with those of Burma, run north and south.' The hills containing the coal measures of Sibsagar and Lakhimpur consist of an enormous thickness of sandstones, the upper series of which are topped with conglomerates and clay.

The climate of Assam is characterized by coolness and extreme humidity. In the Rains especially, it does not stimulate to prolonged physical exertion. Mr. A. Primrose in his note on Assam states that except for four months of the year the climate is decidedly bearable to Europeans, but these must be of naturally strong constitution and thoroughly acclimatized, for all the year round malaria is prevalent. There are two distinct seasons:—cold from November to February; rains from March to October. Rainfall is always abundant, so that floods rather than droughts occur, the average amount annually precipitated varying from 85" at Sibsagar, 115" at Goalpara and 124" at Silchar to 458" at Cherrapunji. It has been noted that since the earthquake of 1897 the floods in Lower Assam are of greater intensity and duration. Earthquakes are very frequent all over the province.

As the *Gazetteer* points out, the character of the surface and the climate of Assam are ideal for forest growth and the area of wooded country is very extensive. But the method of cultivation employed by most of the hill tribes does not help in the growth of really great forest. This method is called the JHUM system and is thus described in the *Gazetteer*:—

'A patch of land is cleared with axe and fire, the soil is hoed up and the seeds of hill rice, chillies, cotton, millets, gourds and other vegetables are dibbled in among the ashes. The same plot is seldom cropped for more than 2 or 3 years in succession. After this time the weeds spring up in great luxuriance and further cultivation would destroy the roots of Ikra or Bamboo jungle, upon which the land depends for its fertility. JHUMS are left fallow for as long a time as possible, four years at the very least. This system of cultivation and forest fires have denuded the interior of the hills where the people chiefly live.'

For a Field Collector Assam presents many difficulties.

On the one hand, in the hill areas of the frontier, such as Mishmi Hills, the Abor, or Aka country, the aboriginal tribes resent interference and intrusion, and, as Mr. A. Primrose says in his Note, the greatest tact, patience, and firmness are required in any dealings with them. But he has found that though they deeply feel harshness or what they consider injustice, they do respond to kindly treatment, and, away from outside influence, are a 'very cheerful, simple, honest and likeable lot.' On the other hand the habits of the imported coolies in the cultivated areas and tea districts have practically exterminated all the mammalia around them. The Mundas, Oraons, and especially the Santhals eat any sort of animal, and employ fires, traps and poison as well as bows and arrows in their pursuit of creatures for food. Mr. Primrose remarks on the fact that few animals are ever met with in the forests, in which the silence is intense, just as in the forest of the Sunderbans area. He found better results to be got on the outskirts of the forests or along the streams running into them.

The question of transport is another difficulty. Mr. Wells found roads were for the most part bad, when they did exist at all, bullock carts were hard to obtain, while boat hire was excessive, and heavy mists caused much delay. But for a collector, the climate is the biggest drawback. In his diary, Mr. Wells frequently mentions the loss of his traps by heavy rain washing them away or by debris brought down by the storm covering them up, also the loss of time caused by inundation of the jungle paths. The climate not only affects one's health and working capacity, but it also causes endless trouble in the preparation of specimens. Even with a fire it was found impossible to get the skins of his specimens dry enough to ensure their preservation, and many of these were lost owing to the excessive humidity.

In his diaries Mr. Wells records his thanks for kind assistance to the following:—Mr. Bishop of Godagari; Mr. Brown of Hokenguri; Mr. Garrett, the Forest Officer; Mr. Milroy, whom he accompanied on keddah; Mr. Middleton of Sonepur; Mr. Hinde of Rangiya; Dr. Grant of Margherita; Mr. James, the missionary at Cherrapunji; The late Mrs. Jackson gave invaluable help to Mr. Wells in the Garo Hills area.

The areas visited between October 1919 and July 1921 may best be divided into the following centres:—

1. MARGHERITA—Ledo, Namlang River, Powai, Doom Dooma, Digboi, Bara Hapjan, Kharjan.
2. SADIYA—Dibong River, Lohit Valley.
3. GOLAGHAT—Mikir Hills, Kaziranga Reserve, Kumta.
4. GARO HILLS—Tura, Rangapari, Dura Bandar, Duragiri.
5. SHILLONG AND KHASI HILLS—Mowphlong, Cherrapunji, Laitkynsao, Laiterai, Nongprieng, Nongpoh.
6. JAINTIA HILLS—Jowai, Jarain, Syndai, Shanpung, Konshnong, Hot-springs.
7. S. KAMRUP—Kulsi, Rajapara.
8. CACHAR—Lamsakhang, Langting, Dirangmukh, Hathikhali, Lanka.
9. N. KAMRUP—Angarakhata, Matanga River.
10. MISHMI HILLS—Dening, Tiki, Dreyi, Tejir.

The collection numbers 2097 specimens in 98 species belonging to 70 genera. It includes three new species, viz. :—

Soriculus radulus, *Leggada jacksoniæ*, *Leggada nagarum*; also ten new sub-species, viz. :—*Pithecus pileatus saturatus*, *Pithecus pileatus tenebricus*, *Scotamanes ornatus imbreensis*, *Tupaia belangeri assamensis*, *Tupaia belangeri versuræ*, *Dacnomys wroughtoni*, *Rattus listeri garorum*, *Callosciurus erythræus wellsi*, *Dremonys lokriah garorum*, *Rattus rattus wellsi*. In character it closely resembles that from Nepal described in No. 37, but here there are no bears, no Tatera, no Funambulus represented. Part of the material collected was examined by the late Mr. Wroughton, and his notes, which are in each case acknowledged, have been incorporated in this Report as well as those from the diaries of the Field Collector, Mr. Wells.

(1) HYLOBATES HOOLOCK, Harl.

The Hoolock

(Synonymy in No. 14)

Naga Hills—Margherita, (1,200') ♂1 (juv.), ♀1.

Lakhimpur—Bara Hapjan (300') ♂1, ♀1; Golaghat ♂1.

Mishmi Hills—Chikorpani, ♂1 skeleton.

These creatures are fairly common, but keep mainly to the hills, occasionally coming down to the low country probably in passing from one range to another. The range of this gibbon in the Mishmi Hills seems rather obscure. It has been reputed to range for 100 miles north of Sadiya but the Mishmis do not know of it beyond the Digaun River. The Mishmis say there are only three parties of these gibbons, each about 10 or 12 in number, in the Dening District.

Vernacular name—HUELENG (Khasi)

(2) MACACA ASSAMENSIS, Mc Cl.

The Assam Macaque

(For synonymy see Hinton and Wroughton, J.B.N.H.S., xxvii., p. 668.)

Mishmi Hills—Tegang River (2,000'), ♂2.

Vernacular name—TA'MIM (Mishi).

(3) MACACA MULATTA, Zimm.

The Rhesus

(For synonymy see Hinton and Wroughton, J.B.N.H.S., p xxxii, p. 668.)

Naga Hills—Margherita (200'), ♂1 (juv.)

Khasi Hills—Nongpoh (1,200'), ♀1 (juv.)

Jaintia Hills—Hot Springs (2,400'), ♀1.

S. Kamrup—Rajapara (600') ♂2; Kulsi (750'), ♀2 juv.

Cachar Hills—Lamsakhang (200'), ♀1.

Mishmi Hills—Dening (2,243'), ♀3.

Sibsagar—Golaghat (300'), ♂1.

These are very common near Margherita and Rajapara.

Vernacular name—SHIRIT (Khasi)

(4) PITHECUS PILEATUS, Blyth.

The Capped Langur

(Synonymy in No. 20)

Garó Hills—Tura: (1,400'), ♂ 1, flat skins 3, noskulls.

Khasi Hills—Laitkynsao (2,000'), ♂ 1 juv.

Jaintia Hills—Konshnong (3,000'), ♂ 1 juv., ♀ 2 (1 juv.).

Sibsagar—Golaghat (400'), ♀ 2; Raheng, ♀ 1.

This species is very common round Tura.

Vernacular name—INGAW (Khasi)

Mr. Wells notes that these monkeys are generally found in parties of 5-8, and the cry is a sharp chak-chak repeated at short intervals. The female I shot weighed 18.6 lbs. and measured H. and B. 560' ' ' T. 930' ' ' , H.F. 175' ' ' , Ear 30' ' ' . The young one is much lighter in colour with the tuft at the end of the tail a brownish white and distinctly waved hair.

(5) PITHECUS PILEATUS SATURATUS, Hint. subsp. nov.

The Capped Langur(1921 *Pithecus pileatus saturatus*, Hinton, J. B. N. H. S., xxix, p. 81)

Lakhimpur—Bara Hapjan (200'), ♂ 3.

Sibsagar—Golaghat (400'), ♂ 3.

This sub-species is met with in the low ground, but does not reach the cool jungles at high altitudes inhabited by *Pithecus pileatus pileatus*.

(6) PITHECUS PILEATUS TENEBRICUS, Hint, subsp. nov.

The Capped Langur(1921 *Pithecus pileatus tenebricus* Hinton, J. B. N. H. S., xxix, p. 81)

North Kamrup—Angarakhata (300'), ♂ 3, ♀ 2.

The geographical variation and distribution of the species are fully discussed in the paper cited above.

(7) PTEROPUS GIGANTEUS, Brünn.

The Common Flying Fox

(Synonymy in No. 2)

Sibsagar—Golaghat (300'), ♂ 3, ♀ 1; Doom Dooma, ♂ 4.

(8) CYNOPTERUS SPHINX, Vahl.

The Short-nosed Fruit Bat

(Synonymy in No. 4)

Sibsagar—Golaghat (300'), ♂ 1.

Naga Hills—Margherita (400'), ♂ 1

(9) RHINOLOPHUS AFFINIS HIMALAYANUS K. And.

*The Himalayan Horseshoe Bat*1897, *Rhinolophus affinis*. Blanford Mamm. No. 150 (partim).1905, *Rhinolophus affinis himalayanus*. K. Anderson, P.Z.S., p. 103.

Khasi Hills—Cherrapunji (5,500'), ♂ 2, ♀ 8; Laitkynsao (4,000'), ♂ 8, ♀ 15.

Jaintia Hills—Shangpung (4,000'), ♂ 8, ♀ 15; Syndai (2,500'), ♂ 1 Konshnong (3,000'), ♂ 1, ♀ 2.

The caves at Laiterai, 5 miles from Cherrapunji, where these bats live, were, 200' up the sheer face of a cliff and without a rope could not be got at from any point. The Khasis say that in the months of February, March and April the bats issue every evening from the caves in myriads. From May onwards they are said to hibernate.

(10) RHINOLOPHUS BLYTHI SZECHUANUS, K. And.

*The Szechuan Horseshoe Bat*1918, *Rhinolophus blythi szechuanus* K. Anderson, A. M. N. H., 9, p. 376.

Sibsagar—Golaghat (300'), ? 1,

(11) RHINOLOPHUS FERUM—EQUINUM TRAGATUS Hodgs.

Hodgson's Horseshoe Bat

(Synonymy in No. 23, also No. 36).

Mishmi Hills—Dening (2,240'), ♂ 1.

Vernacular name,—K^h'PWINA (Mishmi).

(12) RHINOLOPHUS LEPIDUS Blyth

The Little Indian Horseshoe Bat

(Synonymy in No. 6)

Jaintia Hills—Syndai (2,500'), ♂ 1; Konshnong (3,000'), ♂ 1, ♀ 1, Shangpung (4,000'), ♂ 1, ♀ 1.

(13) RHINOLOPHUS PEARSONI, Horsf.

Pearson's Horseshoe Bat

(Synonymy in No. 15)

Jaintia Hills—Konshnong (3,000'), ♂ 8, ♀ 7.

Khasi Hills—Cherrapunji (4,500'), ♀ 1.

(14) RHINOLOPHUS PERNIGER, Hodgs.

The Himalayan Horseshoe Bat

(Synonymy in No. 14)

Jaintia Hills—Konshnong (3,000'), ♂ 3, ♀ 1.

(15) RHINOLOPHUS YUNANENSIS, Dobs.

The Yunan Horseshoe Bat

1872. *Rhinolophus yunanensis* Dobson, J. A. S. B., xli, pt. 2, p. 336.

1897. *Rhinolophus pearsoni* Blanford, No. 149 partim.

Mishmi Hills—Dening (2,240'), ♂ 1.

It is preferable to refer this specimen definitely to *yunanensis*, since the measurements are distinctly greater than for *pearsoni*. Mr. Wells noted it had a very strong smell.

(16) HIPPOSIDEROS ARMIGER Hodgs.

The Great Himalayan Leaf-nosed Bat

(Synonymy in No. 14)

Khasi Hills—Laitkynsao (2,000'), ♂ 2, ♀ 1.

Garro Hills—Dura Bandar (1,500'), ♂ 2, ♀ 7.

Jaintia Hills—Konshnong (3,000'), ♂ 15, ♀ 10.

„ Shangpung (4,000'), ♂ 4, ♀ 3.

S. Kamrup—Rajapur (600'), ♂ 3, ♀ 3.

Only the ♂s have the strong smell, and are very fat at this time of the year—August, 1920. H.W.W.

(17) HIPPOSIDEROS CINERACEUS, Blyth.

Blyth's Leaf-nosed Bat

(Synonymy in No. 37)

Khasi Hills—Cherrapunji (4,800'), ♂ 2, ♀ 1.

Jaintia Hills—Shanpung (4,000'), ♂ 1.

Lakhimpur—Golaghat (300') in al. 1.

Naga Hills—Margherita (200') in al. 1

Mishmi Hills—Tiki (1,440'), ♂ 1

(18) HIPPOSIDEROS GENTILIS GENTILIS K. And.

*The Bi-coloured Leaf-nosed Bat.*1918. *Hipposideros gentilis gentilis*. K. Andersen, A.M.N.H., 9, 11, p. 380.

Naga Hills—Margherita (200'), ♂ 1, in al. 2.

Sibsagar—Golaghat (300'), in al. 2.

Garo Hills—Dura Bandar (1,500'), ♂ 3.

Khasi Hills—Cherrapunji (5,300'), ♂ 10, ♀ 9, in al. 8.

,, Laitkynsao (2,000'), ♂ 6, ♀ 6.

Kamrup—Gauhati, ♀ 1; Rajapur, ♂ 1.

Mishmi Hills—Tiki (1,440'), ♂ 2; Dreyi (6000'), ♂ 1.

'Dr. Anderson was at work on the Rhinolophidæ but most unfortunately was prevented from finishing the classification beyond a fragmentary paper from which I have taken this name which only a short time ago we should have listed as *fulvus*.'—R.C.W.

(19) HIPPOSIDEROS LEPTOPHYLLUS, Dobs.

*The Khasi Leaf-nosed Bat*1874. *Phyllorhina leptophylla*. Dobson, J.A.S.B., p. 234.1891. *Hipposideros leptophyllus*. Blandford, Mamm. No. 160.

Khasi Hills—Laitkynsao (2,000'), ♂ 20, ♀ 3, in al. 3.

,, Nongpoh (1200'), ♀ 2.

S. Kamrup—Rajapur (600'), ♀ 1.

(20) COELOPS FRITHI, Blyth.

*The Tailless Leaf-nosed Bat*1848. *Coelops frithi*, Blyth. J.A.S.B., xvii, p. 251.1891. *Coelops frithi*, Blandford, Mamm. No. 168.

Khasi Hills—Cherrapunji (4,500'), ♂ 1.

(21) MEGADERMA SPASMA MAJUS, K. And.

*The Chindwin Megaderm*1891. *Megaderm spasma*, Blandford, Mamm. No. 170.1918. *Megaderma spasma majus*, K. Andersen, A.M.N.H., 9, 11, p. 383.

Kamrup—Kulsi (150'), ♀ 1; Rajapur (600'), ♂ 2, ♀ 3.

This sub-species was originally reported under the name of *M. s. trifolium* which in 1918 was sub-divided by Dr. K. Andersen into *trifolium*, *majus*, *medium* and *minus*.

(22) LYRODERMA LYRA, Geoff.

The Indian Vampire Bat

(Synonymy in No. 1)

S. Kamrup—Polasbari, (600'), ♀ 1.

N. Kamrup—Angarakhata (300'), ♂ 1, ♀ 2.

(23) BARBASTELLA DARJELINGENSIS, Hodgs.

The Eastern Barbastel

(Synonymy in No. 26)

Jaintia Hills—Shangpung (4,000'), ♂ 1.

(24) PIPISTRELLUS BABU, THOS.

The Babu Pipistrel

(Synonymy in No. 26)

S. Kamrup—Rajapura (600'), ♂ 3, ♀ 9.

(25) PIPISTRELLUS COROMANDRA Gray.

The Coromandel Pipistrel

(Synonymy in No. 5)

Jaintia Hills—Konshnong (3,000'), ♂ 1, ♀ 2.

Lakhimpur—Sadiya (600'), ♂ 2, ♀ 6.

(26) *PIPISTRELLUS MIMUS*, Wroughton.

The Southern Dwarf Pipistrel

(Synonymy in No. 1)

Sibsagar—Golaghat (350'), ♀ 6, in al ♂ 3.

Khasi Hills—Laitkynsao (2,500'), ♂ 1, in al 1.

N. Kamrup—Angarakhata (300'), ♂ 2.

S. Kamrup—Rajapura (600'), ♂ 4, ♀ 1.

'I named this Bat from the seacoast near Bombay, yet almost every Survey Collection from India proper, i.e., excluding Burma has obtained it.'

(27) *SCOTOPHILUS KUHLI*, Leach.

The Common Yellow Bat

(Synonymy in No. 1)

Naga Hills—Margherita (200'), ♂ 1.

S. Kamrup—Rajapura (600'), ♂ 5, ♀ 2.

Four specimens caught at Rajapura on November 14, 1920, were apparently hibernating. Colour is variable in both sexes, bright chestnut in three specimens, yellowish brown in the fourth. These bats had a heavy coating of fat and were very strong smelling.

(28) *SCOTOPHILUS WROUGHTONI*, Thos.

Wroughton's Bat

(Synonymy in No. 1)

Garro Hills—♂ 1 in al.

(29) *SCOTOMANES ORNATUS IMBRENSIS*, Thos. subsp. nov.

The Assam Harlequin Scotophil

1891. *Nycticejus ornatus*. Blanford Mamm. No. 197.

1922. *Scotomanes ornatus imbrencis*. Thomas, J.B.N.H.S., xxvii, p. 772.

Garro Hills—Duragiri (3,500'), ♀ 1, in al 1.

Jaintia Hills—Konshnong (3,000'), ♂ 1; Jowad (4,500') ? 1.

Mishmi Hills—Dening (2,240'), ♂ 1, ♀ 3.

Mr. Wells in his Diary for June, 1921, says that 'this species was fairly common over open ground. These bats have only just come out of hibernation as there were none about when we were last at Dening (March, 1921).'

Vernacular name for Bats—MAHTRA or TOBOK (Garro Hills), LYMBIT (Khasi)

(30) *MYOTIS SILIGORENSIS*, Tones.

The Darjeeling Bat

(Synonymy in No. 37)

Jaintia Hills—Konshnong (3,000'), ♂ 1, ♀ 1.

(31) *LEUCONOE* sp.

Jaintia Hills—Konshnong (3,000').

'The specimen is too young for identification.' R.C.W.

(32) *MURINA CYCLOTIS*, Dobs.

The round-eared Tube-nosed Bat

(Synonymy in No. 23)

Jaintia Hills—Konshnong (3,000'), ♂ 1, ♀ 1.

(33) *MURINA TUBINARIS*, Scully.

Scully's Tube-nosed Bat

(Synonymy in No. 25)

Jaintia Hills—Shangpung (4,000'), ♀ 1.

Mishmi Hills—Dening (2,240'), ♂ 2, ♀ 7.

These bats apparently breed in April and May as all five females taken at Dening in May 1921 were pregnant—three containing each one, and two with each two young.

(34) KERIVOULA HARDWICKEI, Horsf.

Hardwicke's Bat

(Synonymy in No. 26)

Jaintia Hills—Konshnong (3,000'), ♂ 6, ♀ 4.

Shangpung (4,000'), ♂ 6, ♀ 4.

S. Kamrup—Rajapura (600'), ♀ 1 unsexed 1.

(35) TUPAIA BELANGERI ASSAMENSIS, Wrought, subsp. nov.

The Assam Tree Shrew

1921. *Tupaia belangeri assamensis*. Wroughton, J.B.N.H.S., xxvii, p. 599

Lohit Valley (800'), ♂ 2.

Garó Hills—Tura ♂ 4, ♀ 2; Dura Bandar, ♀ 1; Duragiri, ♀ 2.

Khasi Hills—Laitkynsao (2,000'), ♀ 2.

'The forms *chinensis* and *siccata* are both of a pale grizzle of which the modifying component is yellow and gives the general effect of 'olive green', while in *yunalis* and *assamensis* the modifying colour of the grizzle is red and there is no green tinge, general effect being dark brown.'—R.C.W.

Vernacular name—MAD (Garó); KMUD (Khasi)

(36) TUPAIA BELANGERI VERSURÆ, Thos., subsp. nov.

The Mishmi Tree Shrew

1922. *Tupaia belangeri versuræ*. Thomas, J.B.N.H.S., xxviii, p. 428.

Mishmi Hills—Dening (2,240'), ♂ 8, ♀ 3; Dreyi (6,000'), ♂ 1, ♀ 1.

Mr. Wells notes 'I understand this *Tupaia* breeds during May and June and makes its nest in holes, under roots of trees, etc. Three to five young are born naked and blind.'

Vernacular name—A' CHUBUA (Mishmi)

(37) TALPA MICRURA, Hodgs.

The Short-tailed Mole

(Synonymy in No. 23)

Mikir Hills (1,400'), ♀ 1; Mishmi Hills—Dening (2,240'), ♀ 2.

One specimen taken at Dening was of a blue gray colour with a short tail (10 mm.) black, with numerous hairs at the end. The snout is flesh coloured with a small white streak on the face above.

(38) PARASCAPTOR LEUCURUS, Blyth.

The White-tailed Mole

(Synonymy in No. 36)

Khasi Hills—Cherrapunji (4,500'), ♂ 2, ♀ 2.

Jaintia Hills—Shangpung (4,500'), ♂ 1, ♀ 2; Jowai (4,500'), ♀ 2.

These moles frequent gardens and potato patches. They throw out an offensive smell when handled or confined in a box.

Vernacular name—KYNDAT (Khasi)

(39) SORICULUS CAUDATUS, Horsf.

Hodgson's Brown-toothed Shrew

(Synonymy in No. 15)

Mishmi Hills—Dreyi (6,000'), ♂ 1, ♀ 3, in al. 2.

(40) SORICULUS RADULUS, Thos., sp. nov.

The Mishmi Brown-toothed Shrew

1922. *Soriculus radulus*. Thomas, J.B.N.H.S., xxviii, p. 429.

Mishmi Hills—Dreyi (6,000'), ♀ 1.

(41) CROCIDURA RUBRICOSA, And.

Anderson's Assam Shrew

(Synonymy in No. 25)

- Naga Hills—Margherita (200'), ♂ 1, ♀ 2.
 Lakhimpur—Sadiya (500'), ♂ 2, ♀ 1.
 Sibsagar—Golaghat (300'), ♂ 2, ♀ 3, in al. 1.
 Garo Hills—Tura, (1,400'), ♂ 1, in al. 1.
 Khasi Hills—Laitkynsao (2,000'), ♂ 1.
 Jaintia Hills—Jowai (4,400'), ♂ 1, ♀ 1; Konshnong (3,000'), ♂ 1 Syndai in al 1; Shangpung (4,000'), ♂ 1.
 N. Kamrup—Angarakhata (300'), ♂ 21, ♀ 14.
 S. Kamrup—Rajapura (600'), ♂ 1.
 N. Cachar—Lanka (400'), ♂ 2, ♀ 2.
 Mishmi Hills—Dening (2,240'), ♂ 4, ♀ 1; Dreyi (6,000'), ♂ 1, ♀ 3.
 Vernacular name—CHUCHU (N. Kamrup) TA'PEKING (Mishmi) for all shrews.

(42) PACHYURA, sp.

The Musk Rat

- Naga Hills—Margherita (200'), ♂ 1, ♀ 2.
 Sibsagar—Golaghat (300'), ♂ 5, ♀ 5.
 Garo Hills—Tura (1,440'), ♂ 2, ♀ 3; Dura Bandar (1,200'), ♀ 2, Duragiri (3,000'), ♀ 1.
 Khasi Hills—Nangpoh (1,200'), ♀ 2.
 Jaintia Hills—Shangpung (4,000'), ♀ 4.
 N. Kamrup—Angarakhata (300'), ♂ 26, ♀ 14.
 S. Kamrup—Rajapura (600'), ♂ 9, ♀ 7.
 N. Cachar—Lanka (300'), ♂ 1, ♀ 3.
 Mishmi Hills—Tiki (1440'), ♂ 2.

At Golaghat Mr. Wells noted that these shrews have large burrows easily recognized by the wide mouth as though the earth had fallen in for a few inches before the actual hole commences. The species was not found at Dening though he got it at Tiki, 1,000' lower, under similar conditions.

Vernacular name—CHUCHANDRA (N. Kamrup.)

(43) PACHYURA HODGSONI, Blyth.

The Himalayan Pigmy Shrew

(Synonymy in No. 15)

- Jaintia Hills—Shangpung (4,000'), ♂ 2, ♀ 1.

(44) ANOUROSOREX SQUAMIPES, A.M. Edw.

The Assam Short-tailed Shrew

(Synonymy in No. 25)

- Garo Hills—Tura, ♀ 1.
 Jaintia Hills—Jowai (4,500'), ♂ 8, ♀ 4; Shangpung (4,200'), ♂ 8, ♀ 11;
 Konshnong (3,000'), ♂ 1.
 Mishmi Hills—Dening (2,240'), ♂ 19, ♀ 8; Dreyi (6,000'), ♂ 10, ♀ 8.

(45) FELIS AFFINIS, Gray.

The Jungle Cat

(Synonymy in No. 1)

- N. Kamrup—Angarakhata (300'), ♂ 1.
 Mishmi Hills—Dening (2,240'), ♂ 1.
 Vernacular name—A'CHAGUN (Mishmi.)

(46) FELIS BENGALENSIS, Kerr.

The Leopard Cat

(Synonymy in No. 11)

- Garo Hills—Tura (1,450'), ♂ 1.
 Khasi Hills—Shillong (5,200'), ♂ 1.
 Jaintia Hills—Jowai (4,300'), ♀ 1 juv.
 Vernacular name—KUI (Khasi) for all wild cats.

(47) FELIS TEMMINCKI, Vigors and Horsf.

The Golden Cat.

(Synonymy in No. 14)

Mishmi Hills—Teju (3,000'), ♂ 1.

The Mishmis say that this cat breeds in hollow trees and has two young, also that it kills fawns and other small animals.

(48) PEIONDON PARDICOLOR (Hodgs.)

The Indian Tiger Civet

(Synonymy in No. 23)

Mishmi Hills—Dening (2,240'), ♀ 1.

(49) VIVERRA ZIBETHA, L.

The Large Indian Civet.

(Synonymy in No. 14, also see No. 17)

Lakhimpur—Sadiya (800'), ♂ 1.

Sibsagar—Golaghat (300'), ♂ 2, ♀ 3, juv. 1, unsexed 1.

Garó Hills—Tura ♂ 1; Duragiri (3,000'), ♀ 1; Dura Bandar ? 1.

Jaintia Hills—Shangpung (4,000'), ♀ 1; Konshnong (3,000'), ♂ 1, ♀ 1.

S. Kamrup—Rajapura (600'), ♂ 2, ♀ 2.

Vernacular name—MATNI (Garó.)

(50) VIVERRICULA MALACCENSIS, Gmel.

The Small Indian Civet

(Synonymy in No. 3)

Sibsagar—Golaghat (300'), ♂ 1, ? 1.

N. Kamrup—Angarakhata (300'), ♂ 3.

Mr. Wells notes that this civet can with ease climb a vertical tree trunk.

Vernacular name;—MATJOL (Garó) : BSHAD (Khasi).

(51) PARADOXURUS STRICTUS, Hodgs.

The Sikkim Palm Civet(Synonymy in No. 16, under *hermaphorditus*)

Sibsagar—Golaghat (300'), ♂ 1, flat skin 1, juv. 1.

Garó Hills—Dura Bandar (1,300') ? 1.

N. Kamrup—Darranga River (400'), ♂ 1.

(52) PAGUMA GRAYI, Benn.

The Himalayan Palm Civet

(Synonymy in No. 15)

Garó Hills—Duragiri (300'), ♂ 1.

Mishmi Hills—Dening (2,240') 1 dried skin.

(53) HERPESTES NEPALENSIS, Gray.

The Nepal Mongoose

(Synonymy in No. 19)

Lakhimpur—Kazirunga (250'), ♀ 4.

Sibsagar—Golaghat (300'), ♀ 1.

Jaintia Hills—Konshnong (3,000'), ♂ 1 juv.

N. Kamrup—Angarakhata (300'), ♂ 2, ♀ 5.

The survey took this mongoose in Midnapur, Bengal.

'Gray gave the type locality as "North India see Report No. 27."'—R.C.W.

Vernacular name:—ANGREBU (Garó) ; KSAR (Khasi) :

(54) HERPESTES URVA, Hodgs.

The Crab-eating Mongoose.

(Synonymy in No. 23)

Garó Hills—Tura unsexed 1.

S. Kamrup—Rajapura (600'), ♀ 1 weighing 3 lbs.

(55) CANIS INDICUS, Hodgs.

The Jackal

(Synonymy in No. 1, under *C. aureus*)

Jaintia Hills—Shangpung (4,000'), flat skin 1.

Jowai (4,500'), juv. 2.

Naga Hills—Margherita, ♂ 1.

Sibsagar—Golaghat (300'), ♂ 1.

S. Kamrup—Rajapura (600'), ♀ 1.

Mr. Wells made some interesting notes on the 'calling' of these animals. 'At Lanka in October 1920, he wrote, 'I heard the Jackals calling for the first time since the beginning of the hot weather. The local natives confirm the fact that one hears them only in the cold weather.'

(56) CHARRONIA FLAVIGULA, Bodd.

The Northern Indian Marten

(Synonymy in No. 15)

Garó Hills—Tura (1,400'), ♂ 1.

This is a very sly and wary creature, who turns hives over and eats the honey.

Vernacular name—MAPRA (Garó Bsong) (Khasi).

(57) MELOGALE PERSONATA, Geoff.

The Burmese Ferret Badger

(Synonymy in No. 16 under *Helictis personata*)

Khasi Hills—Laitkynsao (2,500'), ♂ 1.

Mr. Thomas has recently revived the generic name *Melogale*. This animal is not common anywhere in these hills, so the Khasis say. A shikari told Mr. Wells that there is a belief amongst the Khasis that when a 'Shrong' or village Tomcat grows very large it runs away to the jungle and turns into a K'SA, as this *Melogale* is called by the people of Shella, a small hill state.

Vernacular name—WAKSAGEL (Garó) : B'SHAD (Khasi) meaning a kind of civet.

(58) ARCTONYX COLLARIS, F. Cuv.

The Hog Badger

(Synonymy in No. 25)

Jaintia Hills—Hotsprings (2,400'), ♀ 1.

(59) LUTRA LUTRA NAIR, F. Cuv.

The Common Otter

(Synonymy in No. 11)

Lakhimpur—Sadiya (500'), ♂ 1 ; Kaziranga (250'), ♀ 1 flat skin.

S. Kamrup—Rajapura ♂ 1.

Jaintia Hills—Shangpung (4,000'), 1 flat skin.

Jowai (4,500') juv. 2.

These otters are becoming a trouble to the Assamese fishermen clearing their lines every evening, leaving only the heads of the fish. At Rajapura a small ♀ otter, immature (weight 3'5 lbs.) was brought in from a 'bhil,' which had a most pathetic cry when handled, something like a kitten.

Vernacular name—KSIH (Khasi).

At Shillong Mr. C. H. Holder showed Mr. Wells some skins of the smooth Indian otter (*Lutrogale barang tarayensis*), which he had caught in the neighbourhood. He told Mr. Wells that the otters made their appearance in the district only when he started breeding trout and he has killed quite a number lately. There must be some special reason to account for their coming, 'as the only water anywhere near is the mountain stream and no otter could live in that as there is less depth than 6''.'

(60) AMBLONYX CINEREA, Illig.

The Clawless Otter(Synonymy in No. 11 under *Aonyx cinerea*)

Sibsagar—Golaghat (300'), ♂ 1, ♀ 1.

Garó Hills—Tura unsexed 1.

Jaintia Hills—Hot springs (2,400'), juv. 2; Jowai (4,500'), ♀ 1.

(61) HELARCTOS MALAYANUS, Raff.

The Malay Bear

(Synonymy in No. 14)

Garó Hills—Duragiri (3,000'), ♀ 1 juv.

These bears are frequently seen on the very high trees, seeking berries.

NOTE.—*Melursus ursinus*. While in the Lohit Valley Mr. Wells wrote, 'Although I have not seen any, the Sloth Bear is evidently common as I have often come across large holes dug, by these animals (so the natives tell me) in their search for ants' and bees' nests, etc. To-day I saw four such 'holes within 20 yds.—one nearly 3 ft. 6 in. diameter and about 5 ft. deep.'

(62) PETAURISTA YUNNANENSIS, And.

Anderson's Flying Squirrel

(Synonymy in No. 36)

Mishmi Hills—Tiki (1,440'), ♂ 2; Dreyi (4,000'), ♂ 1.

At Dreyi both this and the following species occurred fairly commonly but it was difficult to capture specimens. These animals have a peculiar 'miawing' cry when in flight, generally uttered just before they alight on their objective. It would appear that they live in small colonies, probably 2 ♂s and some ♀s.

Vernacular name:—K'MAI (Mishmi).

(63) PTEROMYS (HYLOPETES) ALBONIGER, Hodgs.

The Particoloured Flying Squirrel

(Synonymy in No. 20)

Jaintia Hills—Konshnong (3,000'), ♂ 14, ♀ 14.

Khasi Hills—Laiterai (2,400') unsexed 1.

Mishmi Hills—Denning (2,240'), ♂ 2, ♀ 2.

For their home these squirrels usually seem to choose a tree with a trunk clear to 40 or 50 ft. up. They are used for food.

Vernacular name—HAGRAI (Mishmi).

(64) RATUFA GIGANTEA, McCL.

The Assam Giant Squirrel

(Synonymy in No. 14.)

Lakhimpur—Sadiya (500'), ♀ 1.

Garó Hills—Tura (1,300'), ♂ 1, ♀ 1; Duragiri (1,600'), ♂ 1.

S. Kamrup—Rajapura (600'), ♂ 1.

Cachar Hills—Hatikhali (1,600'), ♂ 1; Langting (1,500'), ♂ 1, ♀ 1.

N. Kamrup—Matanga River (2,500'), ♂ 3, ♀ 1.

Mishmi Hills—Tiki (1,440'), ♂ 2, ♀ 2.

Denning (2,240'), ♂ 1, ♀ 2; Dreyi (6,000'), ♂ 1, ♀ 2.

This squirrel is common at Tura where the natives call it the Palm Marten.

Vernacular name—TA'RAI (Mishmi.)

(65) DREMOMYS LOKRIAH GARONUM Thos. subsp. nov.

*The Orange-bellied Garó Hill Squirrel*1922. *Dremomys lokriah garonum*, Thomas, J.B.N.H.S., xxviii, p. 430.

Garó Hills—Tura (1,200'), ♂ 1; Duragiri (3,000'), ♀ 1.

Jaintia Hills—Konshnong (3000'), ♂ 1, ♀ 2.

Jowai (4,500'), ♀ 1.

Khasi Hills—Laitkynsao (2,500'), ♀ 2 juv.

S. Kamrup—Rajapura (600'), ♂ 1.

(66) DREMOMYS LOKRIAH SUBFLAVIVENTRIS, Horsf.

*The Orange-bellied Mishmi Hills Squirrel*1843. *Sciurus subflaviventris* McClelland, Gray. List., M.B.M., p. 144.1922. *Dremomys lokriah subflaviventris*, Thomas, J.B.N.H.S., xxvii, p. 429.

Mishmi Hills—Dreyi (6,000'), ♂ 14, ♀ 18.

'The cry of this squirrel is quite distinct from any other, being a sharp 'squeaky chatter, continually repeated.'

The fur of one shot at Dreyi in June, 1921, was 'stained with the juice of a calladium fruit on which most squirrels and rats are at present feeding.' While *Callosciurus erythræus* seems to prefer the higher creeper-clad trees, *Dremomys lokriah* appears to be mostly a ground feeder, more frequently below than above the 15 ft. level from the ground, making free use of the creeper when moving about. Of the two, *Dremomys lokriah* seems to be the more active, though both can move with remarkable rapidity when occasion demands, e.g., whilst courting.—H. W. W.

(67) CALLOSCIURUS ERYTHRÆUS ERYTHRÆUS, Pallas.

*Pallas' Squirrel*1778. *Sciurus erythræus*. Pallas, Glires, p. 377.1891. *Sciurus erythræus*. Blanford, Mamm. No. 245 (partim).

Garó Hills—Tura (1,300'), ♂ 11, ♀ 2; Duragiri (3,000'), ♀ 1.

Khasi Hills—Tharia (1,100'), ♂ 1; Nangpoh (1,400'), ♀ 1.

S. Kamrup—Rajapura (600'), ♂ 7, ♀ 4; Kulsí (150'), ♀ 1.

Note by Mr. Wroughton—'This species established so long ago has in recent years been regarded with some doubt as to its exact identity. This fine series quite confirms Blyth's identification of the Garó Hills as the type locality and may now be confidently accepted as *erythræus*, Pallas.' Mr. Wells found the species fairly common at Tura in June, but very scarce everywhere in April. The Khasis snare them with a bamboo and gut trap, using jackfruit as a bait. Several caught at Sadiya in December, 1920, had their tails bitten off.

Vernacular name—RISANG (Khasi); KELKUTA (Garó).

(68) CALLOSCIURUS ERYTHRÆUS ERYTHROGASTER, Blyth.

*The Manipur Squirrel*1842. *Sciurus erythrogaster*, Blyth, J.A.S.B., xi, p. 970.1891. *Sciurus erythræus*. Blanford, Mamm. No. 245 (partim).

Jaintia Hills—Syndai (2,500'), ♂ 5, ♀ 1.

Khasi Hills—Moostah (2,000'), ♂ 1, juv. 1.

'I place this series under this name with some hesitation. On the whole they conform fairly well with the series representing *erythrogaster* in the Museum collection from S. Manipur. It is however, somewhat startling to find them living immediately close to the true *erythræus* type. Three of the six specimens have red ears, indication of their relationship to *erythræus* proper.'—R.C.W.

The two specimens from Moostah were added after the foregoing was written. They have redtipped tails but are otherwise similar to those from Syndai.

(69) CALLOSCIURUS ERYTHRÆUS NAGARUM, Thos. and Wrought.

The Naga Squirrel

(Synonymy in No. 20)

Naga Hills—Margherita (400'), ♀ 1; Powai (400'), ♂ 1.

Lakhimpur—Digboi (500'), ♂ 1, ♀ 3; Mikir Hills (600'), ♂ 1, ♀ 1.

N. Lakhimpur (1,500'), ♂ 1.

Cachar Hills—Langting (1,500'), ♀ 1.

These squirrels are most easily got at sunset as they then make for the tops of 'nahora' trees, on whose berries they feed.

(70) CALLOSCIURUS ERYTHRÆUS AQUILO, Wroughton.

The North-Eastern Squirrel

1921. *Callosciurus castaneoventris aquilo*. Wroughton, J.B.H.N.S., xxvii, No. 3, p. 601.

1921. *Callosciurus erythræus aquilo*. Wroughton, J. B. N. H. S., xxvii, No. 4, p. 775.

Lakhimpur—Dibong River (500'–800'), ♂ 7, ♀ 4.

Mishmi Hills—Tiki (1,440'), ♂ 2, ♀ 2; Deniag (2,240'), ♂ 30, ♀ 24.

Dreyi (6,000'), ♂ 10, ♀ 9.

Mr. Wroughton finally placed this squirrel as a subspecies of *erythræus*. Mr. Wells notes in May at Deniag 'several busy in a simul trees, carrying 'off mouthfuls of the cotton in various directions, some to holes in trees, and 'returning for more. They are very partial to a fruit similar in shape and 'size to an elephant apple but with a smooth skin and sweet pulp. *Hylobates* 'and *Ratufa* also eat this fruit.'

Vernacular name—TWAJA (Mishmi).

(71) CALLOSCIURUS ERYTHRÆUS WELLSI, Wrought.

Well's Squirrel

1921. *Callosciurus erythræus wellsi*. Wrought, J.B.N.H.S., xxvii, p. 775.

Jaintia Hills—Jowai (4,500'), ♀ 1; Shangpung (4,000'), ♂ 3, ♀ 4.

Konshnong (3,000'), ♀ 3.

'In view of the kaleidoscopic character of the patterns in this genus in 'Assam it seems to me every opportunity ought to be taken to give names 'to the local races wherever possible. In the present case the whitish tail 'tag, though only a minor character is so constantly present I have not 'hesitated to name this form as a definite local race.'

(72) TOMEUTES BLYTHI, Tytler

Blyth's Squirrel

1843. *Sciurus assamensis*, Gray. List Mamm., B.M., p. 143.

1854. *Sciurus blythi*, Tytler. A.M.N.H., 2, xiv, p. 172.

1891. *Sciurus lokroides*, Blanford. Mamm. No. 251 (partim).

1915. *Tomeutes lokroides*, Thomas, A.M.N.H., xv, p. 385.

Naga Hills—Margherita (1,200'), ♂ 1, ♀ 1; Namdang (200'), ♀ 1 juv.

Sibsagar—Golaghat (300'), ♂ 2, ♀ 4; Kaziranga, ♀ 1.

Garo Hills—Rangapani (400'), ♂ 2; Tura (1,500'), ♂ 4, ♀ 2.

Duragiri (3,000'), ♂ 2, ♀ 1.

Khasi Hills—Laitkinsao (2,500'), ♂ 1, ♀ 1; Nongpoh (1,200'), ♀ 1.

Jaintia Hills—Syndai (2,500'), ♀ 1 juv.

Cachar Hills—Lanka (400'), ♀ 10; Tejnari (1,500'), ♂ 1.

Doiangmukh (600'), ♀ 2; Doiang River (1,600'), ♀ 1.

Lamsakhang (450–800'), ♂ 1, ♀ 1.

Hatikhali (1,500'), ♂ 2.

S. Kamrup—Rajapura (600'), ♀ 2.

N. Kamrup—Matanga River (2,500'), ♂ 3, ♀ 2; Boganadi (2,000'), ♂ 1, ♀ 1.

Angarakhata, ♂ 1, ♀ 2.

Note on the genus *Tomeutes* by Mr. Thomas:—

The study of these specimens has enabled me to clear up a point of nomenclature which affects all the *Tomeutes* of the *mearsi* type. In 1843 Gray used in his List the name of *Sciurus assamensis* but without a description, the name being therefore a *nomen nudum*. In 1854 Tytler described the squirrel of Dacca as *Sciurus blythi*, the typical specimens of which are now in the British Museum. In 1855 Blyth, writing of these squirrels used the name *Sciurus assamensis* with a secondary name or synonymy *S. blythi*, ignoring the fact that at that date *assamensis* was a *nomen nudum*, and added a description which has been accepted as giving validity to *assamensis*. But as his species *assemensis* is also *blythi*, the latter name, being earlier in date, must have preference. Both names have hitherto been provisionally put as synonyms of *Tomeutes lokroides*, Hodgs. But the present series includes some showing the thigh-patch whitish as in *mearsi*, instead of red as in *lokroides*.

Consequently if, as in Wroughton's synopsis, all the forms of the *lokroides* group with whitish or buffy patches are considered as one species with several

sub-species, the name of the species must be changed to *blythi* and thus the three Burmese species would be

Tomeutes blythi virgo,
 „ „ *bellona*,
 „ „ *mearsi*.

Whether this typical *T. blythi* of Dacca and other parts of Assam is quite the same as the nearest sub-species *T. blythi virgo* is a question which it will be better to leave for decision until specimens from the intervening countries are available. The two are undoubtedly very close to one another, but, *blythi* seems on the average to be of a paler buffy above than *virgo*, while the front aspect of its thighs is of a bluey greyish, which is replaced in *virgo* by buffy. Provisionally therefore I should call all four forms sub-species of *Tomeutes blythi*.

(73) *TOMEUTES STEVENSI*, Thos.

Steven's Squirrel

(Synonymy in No. 20)

Naga Hills—Ledo (200'), ♂ 1.

'I identify this specimen as *stevensi* with very considerable hesitation. A very young specimen, provisionally identified as this species, taken by the Survey in July, North of Hkamti on the Chindwin River, shows in a slight degree the dark bluish head and feet and the ochraceous hip patch. On the other hand, the type taken in the Abor Miri country and two other specimens collected in the same area in February-March show no smallest sign of either of these characteristics. Here the matter must rest until more material is available for examination.' R.C.W.

(74) *TAMIOPS MACCLELLANDI*, Horsf.

The Himalayan Striped Squirrel

(Synonymy in No. 20)

Lakhimpur—Sadiya (240'–600'), ♂ 6, ♀ 3.

Garó Hills—Tura (4,000'), ♂ 1.

Cachar Hills—Langting (1,500'), ♀ 1.

Mishmi Hills—Tetsam (700'), ♂ 1; Tiki (1,440'), ♀ 2.

Dening (2,240'), ♂ 8, ♀ 7; Teju (3,000'), ♂ 1.

Dreyi (6,300'), ♂ 5, ♀ 3.

These little squirrels usually frequent the large open trunks of the *nahor* and *simul* trees. Very seldom are they found in trees of dense growth. The Mishmis say that this species does a fair amount of damage to their Indian corn when ripening, and only a well-aimed stone can dislodge them. They say that *C. erythraeus* also feeds on Indian corn but not to the same extent as it is more easily driven away. This *Tamioops* is very fond of a tree, *Dicellostyles jujubifolia*, some of these trees holding 3 or 4 of the squirrels at a time. A female killed June 21 at Dreyi contained two fetuses.

(75) *BANDICOTA MEMORIVAGA*, Hodgs.

The Smaller Bandicoot Rat

(Synonymy in No. 29)

N. Kamrup—Angarakhata (300'), ♂ 1.

Mr. Thomas has recently decided that *B. elliotina* and *memorivaega* are identical and thus the former name should now be dropped.

(76) *GUNOMYS BENGALENSIS*, Gray and Hardwicke

The Bengal Mole Rat

(Synonymy in No. 19)

Sibsagar—Golaghat (300'), ♂ 9, ♀ 5 juv.

N. Kamrup—Angarakhata (300'), ♂ 38, ♀ 48.

Specimens of this Mole rat were collected in Kumaon and show in Report No. 15 as *G. tarayensis*, but later were included under *G. bengalensis*.

(77) DACNOMYS WROUGHTONI, Thos., subsp. nov.

*The Mishmi Hills Giant Rat*1922. *Dacnomys wroughtoni*, Thomas, J.B.N.H.S., xxviii, p. 430.

Mishmi Hills—Dening (2,240'), ♂ 1; Dreyi (6,000'), ♂ 4, ♀ 1.

Mr. Thomas has dealt with this rat under 'Results' in above paper.

(78) RATTUS FULVESCENS, Gray.

The Chestnut Rat

(Synonymy in No. 15)

Naga Hills—Margherita (200'), ♀ 1.

Lakhimpur—Sadiya (500'), ♂ 1, ♀ 1; Kharjan (400'), ♂ 6, ♀ 2.

Digboi (200'), ♂ 2, ♀ 1.

Jaintia Hills—Jowai (4,500'), ♂ 1, ♀ 5; Konshnong (3,000'), ♂ 2.

Shangpung (4,000'), ♂ 1; Jarain (3,000'), ♀ 1.

Khasi Hills—Shillong (3,800'), ♀ 1; Cherrapunji (4,500'), ♂ 2.

Nongpoh (1,200'), ♀ 1; Laitkynsao (2,500'), ♂ 2, ♀ 5.

Mou Phlong (5,500'), ♀ 2.

S. Kamrup—Rajapura (600'), ♂ 1, ♀ 1.

Mishmi Hills—Dening (2,240'), ♂ 27, ♀ 22; Dreyi (6,000'), ♂ 6, ♀ 7.

(79) RATTUS LISTERI, Thos.

The Sikkim Giant Rat

(Synonymy in No. 26)

Mishmi Hills—Dening (2,240'), ♂ 2; Dreyi (6,000'), ♂ 3, ♀ 2.

Vernacular name—KN'ZUNG (Mishmi).

(80) RATTUS LISTERI GARONUM, Thos., subsp. nov.

*The Garo Hills Giant Rat*1921. *Rattus listeri garonum*. Thomas, J.B.N.H.S., xxviii, p. 27.

Garo Hills—Tura (1,400'), ♂ 3, ♀ 2; Duragiri (3,000'), ♂ 1.

Jaintia Hills—Syndai (2,500'), ♀ 1; Nongpoh (1,200'), ♂ 1.

(81) RATTUS MANIPULUS, Thos.

The Manipur Rat

(Synonymy in No. 25)

Sibsagar—Golaghat (400'), ♂ 2, ♀ 2.

Jaintia Hills—Shangpung (4,000'), ♀ 1; Jowai (4,500'), ♀ 1.

(82) RATTUS NITIDUS, Hodgs.

The Nepal Shiny Rat

(Synonymy in No. 15)

Sibsagar—Golaghat (400'), ♀ 1.

Garo Hills—Tura (4,000'), ♀ 1; Duragiri (3,000'), ♀ 2 juv.

Jaintia Hills—Konshuong (3,000'), ♀ 4; Hotsprings (2,300'), ♂ 1, ♀ 3 juv.

Shangpung (4,000'), ♂ 2, ♀ 2 juv. 4.

Jowai (4,500'), ♀ 1 juv. 6.

Mishmi Hills—Denning (2,240'), ♂ 10, ♀ 14.

(83) RATTUS RATTUS SIKKIMENSIS, Hint.

*The Darjiling Tree Rat*1918. *Rattus rattus sikkimensis*, Hinton, J.B.N.H.S., xxvi, p. 61.

Naga Hills—Margherita (200'), ♂ 2.

Lakhimpur—Digboi (250'), ♂ 3, ♀ 1; Lohit Valley (700'), ♂ 1.

Kazirang (250'), ♂ 3, ♀ 5; Kharjan (400'), ♂ 1.

Dibong River (500'), ♂ 1.

Garo Hills—Tura (1,440'–4,000'), ♂ 5, ♀ 5; Duragiri (3,000'), ♀ 2, juv. 2; Rangapani (400'), ♀ 2.

Khasi Hills—Cherrapunji (4,650'), ♂ 1, ♀ 1; Nongpoh (1,200'), ♂ 1, ♀ 4; Nongprieng (2,400'), ♀ 2.

Jaintia Hills—Konshnong (3,000'), ♂ 3, ♀ 5, juv. 2; Shangpung (4,000'), ♀ 2; Jowai (4,500'), ♂ 2, ♀ 1; Hotsprings (2,400'), ♀ 1; Jarain (3,000'), ♂ 1.

Mishmi Hills—Tiki (1,440'), ♂ 1, ♀ 4; Dening (2,240'), ♂ 12, ♀ 5; Dreyi (6,000'), ♂ 3, ♀ 5.

(84) *RATTUS RATTUS TISTÆ*, Hint.*The Himalayan Tree Rat*1918. *Rattus rattus tistæ*, Hinton, J.B.N.H.S., xxvi, p. 61.

Naga Hills—Margherita (200'), ♂ 12 (2 juv.), ♀ 2.

Lakhimpur—Sadiya (500'), ♂ 1; Digboi (250'), ♂ 1 juv. Kharjan (400'), ♂ 1; Kaziranga (250'), ♀ 2; Dibong River (500'), ♂ 3; Lohit Valley (700'), ♂ 4.

Sibsagar—Golaghat (250'), ♂ 8, ♀ 1.

Garó Hills—Tura (1,300'), ♂ 2; Duragiri (3,000'), ♂ 1.

Khasi Hills—Cherrapunji (4,500'), ♀ 1; Nongpoh (1,200'), ♂ 1.

Jaintia Hills—Konshnong (3,000'), ♂ 1, ♀ 1; Jowai (4,500'), ♂ 1, ♀ 1; Syndai (2,500'), ♀ 1; Jarain (3,000'), ♀ 1.

S. Kamrup—Rajapura (600'), ♂ 2, ♀ 10.

N. Kamrup—Angarakhata (300'), ♂ 11, ♀ 15.

Cachar Hills—Denning (2,240'), ♂ 24, ♀ 32 (2 juv.)

(85) *RATTUS RATTUS WELLSI*, Thos.*Well's Rat*1921. *Rattus wellsi*, Thomas, J.B.N.H.S., xxviii, p. 27.

Khasi Hills—Mou Phlong (5,500'), ♀ 1.

(86) *MUS MUSCULUS*, L.*The Common House Mouse*

(Synonymy in No. 1)

Jaintia Hills—Shangpung (4,000'), ♂ 3, ♀ 9.

S. Kamrup—Rajapura (600'), ♂ 1, ♀ 9.

(87) *MUS DUBIUS*, Hodgs*The Nepal House Mouse*

(Synonymy in No. 15)

Naga Hills—Margherita (1,500'), ♂ 1.

N. Cachar—Lanka (400'), ♂ 2, ♀ 1.

S. Kamrup—Rajapura (600'), ♂ 1, ♀ 2.

N. Kamrup—Angarakhata (300'), ♂ 14, ♀ 24.

This mouse has in some instances been classed as *M. manei*, in others as *M. urbanus*, but it now seems preferable to apply the name *dubius* to it until the whole group can be properly worked out.

(88) *LEGGADA BOODUGA*, Gray*The Southern Field Mouse*

(Synonymy in No. 1.)

Sibsagar—Golaghat (300'), ♂ 9, ♀ 10, (4 juv.), in al. 4.

Jaintia Hills—Konshnong (3,000'), ♂ 6, ♀ 11; Shangpung (4,000'), ♂ 6, ♀ 8.

S. Kamrup—Rajapura (600'), ♂ 3, ♀ 8.

N. Kamrup—Angarakhata (300'), ♂ 33, ♀ 34.

(89) *LEGGADA JACKSONIÆ*, Thos., sp. nov.*Mrs. Jackson's Jungle Mouse*1921. *Leggada Jacksoniæ*, Thomas, J.B.N.H.S., xxvii, p. 596.

Garó Hills—Tura (3,000'), ♂ 2, ♀ 1.

Khasi Hills—Laitkynsao (2,000'), ♂ 1.

Jaintia Hills—Konshnong (3,000'), ♂ 1, ♀ 2.

Mishmi Hills—Denning (2,240'), ♂ 16, ♀ 9; Dreyi (6,000'), ♂ 2, ♀ 5.

This species is named in honour of Mrs. Jackson of Tura, to whose kindness and help Mr. Wells owes much of the success which attended his work in the 'Garó Hills.'—THOMAS.

(90) *LEGGADA NAGARUM*, Thos. sp. nov.*The Naga Jungle Mouse*1921. *Leggada nagarum*, Thomas, J.B.N.H.S., xxvii, p. 596.

Sibsagar—Golaghat (300'), ♂ 1.

Jaintia Hills—Konshnong (3,000'), ♂ 3, ♀ 4; Shangpung (4,000'), ♂ 3, ♀ 2.

Khasi Hills—Laitkynsao (2,000'), ♀ 1; Cherrapunji (4,500'), ♀ 1.

Mishmi Hills—Denning (2,240'), ♂ 18, ♀ 8; Dreyi (6,000'), ♀ 1.

(91) LEGGADA PAHARI, Thos.

The Sikkim Hill Mouse

(Synonymy in No. 23)

Lakhimpur—Sadiya (600'), ♂ 1.

Sibsagar—Golaghat (300'), ♂ 1.

(92) VANDELEURIA DUMETICOLA, Hodgs.

Hodgson's Tree Mouse

(Synonymy in No. 16)

Jaintia Hills—Konshnong (3,000'), ♀ 1, juv. 1.

Shangpung (4,000'), ♀ 4.

North Kamrup—Angarakhata (300'), ♂ 9, ♀ 6, in al. 1.

Mishmi Hills—Dening (2,240'), ♀ 1.

(93) HADROMYS HUMEI, Thos.

*The Manipur Bush Mouse*1911. *Hadromys Humei*. Thomas, J. B. N. H. S., 4, xx, p. 999.

North Kamrup—Angarakhata (300'), ♀ 1.

(94) GOLUNDA ELLIOTI, Gray.

The Indian Bush Rat

(Synonymy in No. 1)

North Kamrup—Angarakhata (600'), ♀ 1.

(95) CHIROPODOMYS GLIROIDES, Blyth.

*The Climbing Mouse*1855. *Mus gliroides*, Blyth, J. A. S. B., xxiv, p. 721.1891. *Chiropodomys gliroides*, Blanford, Mamm. No. 270

Jauibia Hills—Konshnong (3,000'), ♀ 1.

'The type specimen of this species, a young animal with half its tail missing, has been lost. These specimens from practically the same locality may confidently be accepted as topotypes. Comparison with Tenasserim specimens, which we have hitherto labelled as *peguensis*, shows that there is no character which they are separable, so that *gliroides* as the older name must be used for both.'—R.C.W.

(96) APODEMUS SPECIOSUS ORESTES, Thos.

*The Szechuan Long-tailed Field Mouse*1911. *Apodemus speciosus orestes*, Thomas, Abster. P. Z. S., p. 49.

Mishmi Hills—Dening (2,240'), ♀ 1; Dreyi (6,000'), ♂ 6, ♀ 2.

(97) EOTHENOMYS MELANOGASTER Libonotus, Hinton, subsp. n.

*Pere David's Vole.*1871. *Arvicola melanogaster*. Milne Edwards, Nouv. Arch. Mus. vii, Bull., p. 23.1891. *Microtus melanogaster*. Blanford, Mamm. No. 307.1923. *Eothenomys melanogaster libonotus*, Hinton, Aun. Mag. Nat. Hist. (9) xi, p. 151.

Mishmi Hills—Dreyi (6,000'), ♂ 3, ♀ 2, unsexed 1.

When revising the members, of the genus *Eothenomys* this, the most south-western representative of that genus, was found to be worthy of recognition. The species is widely distributed in Southern China and Yunman.

(98) RHIZOMYS PRUINOSUS, Blyth.

The Assam Bamboo Rat

(Synonymy in No. 36)

Khasi Hills—Cherrapunji (4,500'), ♂ 2, ♀ 6, juv. 4, in 13.

Laitkinsao (2,000'), ♂ 3, ♀ 2, unsexed 1.

(99) CANNOMYS BADIUS, Hodgs.

The Bay Bamboo Rat

(Synonymy in No. 20)

Garó Hills—Dura Bandar (1,600'), ♀ 2.

Jaintia Hills—Konshnong (3,000'), ♂ 15, ♀ 16.

Hotsprings (2,240'), ♂ 7, ♀ 2.

South Kamrup—Rajapura (600'), ♂ 15, ♀ 7.

(100) ACANTHION HODGSONI Gray.

The Crestless Himalayan Porcupine

(Synonymy in No. 27, also Report No. 37)

Sibsagar—Golaghat (300'), ♀ 2.

Garó Hills—Tura (1,440'), ♂ 1 juv.

(101) ATHERURUS ASSAMENSIS, Thos

The Assam Brush-tailed Porcupine

1921. *Atherurus assamensis*, Thomas. J. B. N. H. S., xxvii, p. 598.

Khasi Hills—Cherrapunji (4,500'), ♂ 1.

Mishmi Hills—Dreyi (6,000'), ♂ 1.

Vernacular name—DYNKHIAT (Khasi).

(102) LEPUS RUFICAUDATUS, Geoff.

The Common Indian Hare.

(Synonymy in No. 15)

Sibsagar—Golaghat (300'), ♂ 2 juv.

North Kamrup—Angarakhata (300'), ♂ 2, ♀ 6 (3 juv.)

The two immature specimens from Golaghat are here included, but are really too young for identification.

(103) CAPRICORNIS SUMATRENSIS RUBIDUS, Blyth.

The Aracan Serow

(Synonymy in No. 36)

Khasi Hills—Moosmai (4,000'), 1 horn only.

Mishmi Hills—Dreyi (6,000'), ♀ 1.

Vernacular names—RAMSAGEL (Garó); KHIAT (Khasi).

(104) BUDORCAS TAXICOLOR, Hodgs.

The Takin.

1850. *Budorcas taxicolor*. Hodgson, J. A. S. B., xix, p. 65, pl. p. 1-3.

1891. *Budorcas taxicolor*. Blanford. Mamm., p. 515.

Mishmi Hills—Dreyi (6,000') ? 1 and 1 horn only.

See also a note on the *Mishmi Takin* in J. B. N. H. S., xxix, p. 550.

Vernacular name—TARKHAN (Mishmi).

(105) MOSCHUS MOSCHIFERUS. L.

The Indian Musk Deer

(Synonymy in No. 23)

Mishmi Hills—Dreyi (9,500') ? ♀, skull in.

(106) MUNTIACUS VAGINALIS, Bodd.

The Barking Deer

(Synonymy in No. 2)

N. Kamrup—Bogranadi (2,000'), ♂ 1; Matanga River (2,500'), ♂ 1.

Mishmi Hills—Dening, B 1 (juv.), ♀ 1.

Vernacular names:—CHOTA HARIN (Garó), SKEI (Khasi).

(107) HYLAPHUS PORCINUS, Zimm.

The Hog Deer

(Synonymy in No. 37)

Lakhimpur—Dibong River (500'), ♀ 2.

N. Kamrup—Angarakhata (300'), ♀ 2.

(108) RUSA UNICOLOR, Bechs.

The Sambhar

(Synonymy in No. 5)

Assam, ♀ 1.

Note by Mr. Wells :— ' I have been asked by a number of people in Assam why it is that Sambhar always have a sore place on the neck. It has been noticed on all Assamese species, but I cannot find out whether this is peculiar to other parts of India also. '

(109) MANIS PENTADACTYLA, L.

The Eastern Pangolin

(Synonymy in No. 37)

Sibsagar—Golaghat (300'), ? 1 flat skin.

' The Survey has taken this species from Mount Popa and Pegu, but it was ' listed as *aurita*. '—R. C. W.

SUPPLEMENT TO REPORT No. 41 :—

A small collection of mammals from the Chilka Lake, Orissa, made by Mr. H. W. Wells.

These specimens were sent along with those from Assam and the Mishmi Hills hence their description may be conveniently tacked on to the Report on those areas.

Chilka Lake is a pear-shaped expanse of water 44 miles long, of area 344 sq. miles in the dry season and 450 sq. miles in the rains, in the district of Puri, Bihar and Orissa. It lies between 19° 28' and 19° 56' N. and 85° 6' and 85° 86' E., and is separated from the Bay of Bengal by only a long sandy ridge about 200 yds. wide. The water in the rains is mainly fresh, but in the dry season is saline.

The collection numbers 17 specimens which belong to 8 genera in 9 species.

(1) PTEROPUS GIGANTEUS, Bruenn.

The Common Flying Fox

(Synonymy in No. 2)

Chilka Lake, ♂ 1, ♀ 2.

(2) SCOTOPHILUS WROUGHTONI, Thos.

Wroughton's Bat

(Synonymy in No. 2)

Chilka Lake, ♀ 1.

(3) SCOTOPHIUS KUHLI, Leach.

The Common Yellow Bat

(Synonymy in No. 1)

Chilka Lake, ♀ 1.

(4) LYRODERMA LYRA, Geoff.

The Indian Vampire Bat

(Synonymy in No. 1)

Chilka Lake, ♂ 1, ♀ 1.

(5) ANATHANA PALLIDA, Lyon.

*The Orissa Tree Shrew*1913. *Anathana pallida*, M. W. Lyon, Proc. U.S.N.M., lxxv, p. 124.

Chilka Lake, ♂ 1, ♀ 1.

(6) CANIS INDICUS, Hodgs.

The Jackal(Synonymy in No. 1 under *C. aureus*)

Chilka Lake, ♀ 1.

(7) FUNAMBULUS PALMARUM, L.

The Palm Squirrel

(Synonymy in No. 2)

Chilka Lake, ♂ 3, ♀ 2 (1 juv.).

(8) BANDICOTA Sp.

Chilka Lake, ♂ 1.

More specimens from this area must be obtained before the species can be definitely determined. It is probably a sub-species of *B. malabaricus*.

(9) RATTUS RATTUS ARBOREUS, Buch-Hamilton.

The Bengal Tree Rat

(Synonymy in No. 47)

Chilka Lake, ♂ 1.

SUPPLEMENT II. COLLECTION MADE BY MR. H. WHISTLER IN KANGRA.

1. *Petaurista inornatus* from Upper Dharmsala ... June 1, 1923.
2. *Eoglaucmys fimbriatus* ♂ from Upper Dharmsala ... ,, 9, 1923.
3. *Ochotona roylei* ♀ from Kareri Lake, Kangra ... ,, 25, 1923.
4. *Felis affinis* ♂ and ♀ from Dharmsala ... ,, 25, 1923.
5. *Vulpes bengalensis* ♀ from Kangra Valley.
6. *Paguma grayi* juv. ♀ from Dharmsala ... June 21, 1923.
7. *Marmota himalayana* ♂ 2 juv. from S. Issoo, Lahul... July 13, 1923.
8. *Felis affinis* ♀ from Dhelu, Mandi State ... ,, 1, 1923.
9. *Canis pallipes* from Lahul (10,500').

Mr. Pocock has kindly given the following note on this wolf :—

Canis pallipes. This wolf is of great interest. The jaws and teeth, retained in the skin, show that the animal was about the same size as the common Indian wolf, *C. pallipes*, but the coat, in accordance with the high altitude at which the wolf was shot, is extremely thick and woolly, exactly as in the larger Himalayan wolf, *C. lupus laniger*. It entirely contradicts Blanford's statement that *C. pallipes* may be distinguished from *C. lupus* by the coat being shorter and with little or no under fur. Though shot in July, the animal retained most of its winter coat. More specimens from this area would be most desirable.

REPORT No. 42, KASHMIR AND PUNJAB AREAS

BY

MARTIN HINTON AND OLDFIELD THOMAS, F.R.S.

(By permission of the Trustees of the British Museum)

COLLECTION	No. 42.
LOCALITY	Kashmir and Punjab.
DATE	July, August, 1921, and October, November, 1923.
COLLECTED BY	Major Stockley, D.S.O., O.B.E., M.C.

The specimens in this collection were obtained from four distinct areas :—

(1) SRINAGAR (5,600')—with Tral (5,800'), Arapul Valley (6,300'), Sardallu (8,700').

Major Stockley says in his notes that the specimens here all come from the 'edge of the cultivation in the Kashmir Valley. The Sardallu specimens were 'taken in a heavily forested nullah, the trees being mainly walnut, chestnut, 'birch, maples in the ravines, with a main forest growth of pine, and a strong undergrowth of skimmia.' The rats and mice were taken at two 'guzar' huts on the edge of the open grass at 9,000'.

(2) RAWALPINDI—with Campbellpore, Chakdulla (2,000'), Chak Lala (1,100'). Here collections were made on the N. E. portion of the Kala Chitta Range, mainly gravel and limestone with a considerable amount of thorn scrub, chiefly babul and ber, with tussocks of speargrass interspersed. Oleander

and pampas grass grow in the nullahs. The heat in summer is intense and water is scarce, being, confined in winter to only a few pools.

(3) BHATTU HISSAR (600')—This is on the N. E. edge of the Bikanir desert, a country of sand dunes with canal irrigation. In this district Major Stockley noted a large increase in the number of Gerbils, probably due to improved conditions brought by the irrigation. But this increase has also caused increase in the number of foxes and birds of prey.

(4) KOHISTAN (Sind)—with Kotri (300') at the S. end of the Khirthar Range and the Barun Valley, a barren country of rocks and sand dunes.

The 249 specimens obtained belong to 28 genera in 38 species as follows :—

(1) HIPPOSIDEROS CINERACEUS, Blyth.

The Little Leaf-nosed Bat

(Synonymy in No. 37)

Bhattu Hissar—♀ 1.

(2) HIPPOSIDEROS FULVUS PALLIDUS, K. And.

The Bicoloured Leaf-nosed Bat

(Synonymy in No. 3, under *H. fulvus*)

Chak Lala (Rawalpindi), ♂ 5, ♀ 6.

(3) PLECOTUS WARDI. Thomas.

The Kashmir Long-eared Bat

1911. *Plecotus wardi*, Thomas, A. M. N. H. (8), vii, p. 209.

Tral (Kashmir)—♂ 2.

The type locality of this species is Ladak.

Major Stockley says that these two specimens 'entered the Tral shooting hut together, and I caught them in a butterfly net.'

(4) PIPISTRELLUS MIMUS, Wroughton.

The Common Dwarf Pipistrel

(Synonymy in No. 1)

Chak Lala—♂ 1; Bhattu Hissar, ♂ 8, ♀ 5.

(5) SCOTOPHILUS KUHLI, Leach.

The Common Yellow Bat

(Synonymy in No. 1)

Rawalpindi—Chak Lala, ♂ 3, ♀ 1; Toupri, ♀ 3.

Bhattu Hissar—♂ 5, ♀ 2.

(6) RHINOPOMA KINNEARI, Wroughton.

The Greater Indian Mouse-tailed Bat

(Synonymy in No. 3)

Ara (Rawalpindi)—♂ 3, ♀ 9.

(7) RHINOPOMA HARDWICKEI, Gray.

The Lesser Indian Mouse-tailed Bat

(Synonymy in No. 3)

Ara (Rawalpindi)—♂ 4, ♀ 2.

(8) PARAECHINUS BLANDFORDI, And.

Anderson's Hedgehog

(Synonymy in No. 24.)

Rawalpindi—Toupri, ♂ 1; Karung, ♂ 1.

(9) PACHYURA, sp.

Arapul Valley (Kashmir)—♂ 1; Bhattu Hissar—♂ 1.

(10) CROCIDURA RUBRICOSA, And.

Anderson's Assam Shrew

(Synonymy in No. 25)

Saradallu (Kashmir)—♂ 2, ♀ 1.

(11) *CROCIDURA ILENSIS*, Miller.

The Illi River Shrew

1901. *Crocidura ilensis*, Miller, Proc. Bio. Soc. Washington, xiv, p. 157.

Bhattu Hissar—♂ 1, ♀ 1.

The two specimens in this collection bear such a strong resemblance to *Crocidura ilensis* that the name may be provisionally applied to them. But the type specimens of this species come from Turkestan, a long way from the Punjab, and the other side of the Himalayas. Thus when more material is available for thorough examination, it may be necessary to reconsider the matter.

(12) *FELIS AFFINIS*, Gray.

The Jungle Cat

(Synonymy No. 1)

Kashmir—Tral, ♂ 1; Arapul Valley, ♀ 1.

Rawalpindi—Toupi, ♀ 1.

(13) *FELIS BENGALENSIS*, Kerr.

The Leopard Cat

1792. *Felis bengalensis*, Kerr. Animal Kingdom, p. 151.

Saradallu (Kashmir), ♀ 1.

(14) *HERPESTES E. PALLENS*, Ryley.

The Pale Grey Mongoose

(Synonymy in No. 12)

Chak Lala—♀ 2.

(15) *HERPESTES AUROPUNCTATUS*, Hodgs.

The Small Indian Mongoose

(Synonymy in No. 27)

Tral (Kashmir)—♂ 1.

(16) *CANIS I. KOLA*, Wroughton.

The Dekkan Jackal

(Synonymy in No. 1 under *O. aureus*)

Rawalpindi—Chak Lala, ♀ 1, Rawal, ♀ 1, Toupi, ♂ 1, ♀ 2.

(17) *VULPES LEUCOPUS*, Blyth.

The Indian Desert Fox

(Synonymy in No. 3)

Ara (Rawalpindi)—♂ 1, ♀ 1; Chakdalla (Campbellpore)—♂ 1, ♀ 1, Bhattu Hissar—♂ 2, ♀ 1.

(18) *PETAURISTA INORNATUS*, Geoff.

The Large Red Flying Squirrel

1844. *Pteromys inornatus*. Geoffroy. Voyage iv. Mamm., p. 62.

Sardallu (Kashmir)—♂ 1, ♀ 1.

(19) *EOGLAUCOMYS FIMBRIATUS*, Gray.

The Smaller Kashmir Flying Squirrel.

1837. *Sciuropterus fimbriatus*. Gray, Charlesworth, Mag. N.H., i, p. 584.

1889. Blandford, Mamm. No. 233.

Saradallu (Kashmir)—♂ 4, ♀ 1.

(20) *FUNAMBULUS PENNANTI ARGENTISCENS*, Wr.

The Northern Five-striped Squirrel

(Synonymy in No. 24)

Rawalpindi—Chak Lala, ♂ 2, ♀ 2; Rawal, ♂ 5, ♀ 2; Toupi, ♀ 1, Karung, ♀ 1.

- (21) GERBILLUS GLEADOWI, Murray.
The Little Hairy-footed Gerbil
 (Synonymy in No. 12)

Bhattu Hissar—♂ 1.

- (22) CHELIONES HURRIANAE, Jerd.
The Indian Desert Gerbil
 (Synonymy in No. 3)

Bhattu Hissar ♀—1; Kohistan (Sind)—♀ 4.

- (23) TATERA INDICA, Hardw.
The Indian Gerbil
 (Synonymy in No. 1)

Chak Lala, ♂ 2, ♀ 2; Rawal, ♂ 2. Jhalar (Campbellpore), ♀ 1
 Bhattu Hissar—♀ 2; Kohistan (Sind), ♂ 1, ♀ 1.

- (24) GUNOMYS BENGALENSIS, Gray and Hardw.
The Bengal Mole Rat
 (Synonymy in No. 19)

Chak Lala, ♂ 8, ♀ 5.

- (25) GUNOMYS WARDI, Thos.
The Kashmir Mole Rat

1908. *Gunomys wardi*, Thomas. J.B.N.H.S., xviii, 4, p. 745.
 Srinagar (Kashmir), ♂ 2.

- (26) RATTUS ALEXANDRINUS, Geoff.
The Egyptian House Rat

1812. *Mus alexandrinus*, Geoff. Desc. de l'Egypte Hist. Nat., ii, p. 733.
 1881. " " Thomas. P.Z.S., p. 533.
 Chakdalla, ♂ 1.

- (27) RATTUS RATTUS RUFESCENS, Gray.
The Common Indian Rat
 (Synonymy in No. 1)

Chak Lala, ♂ 6, ♀ 10.

- (28) RATTUS VICEREX, Boeh.
The North Asian Rat
 (Synonymy in No. 15)

Rawalpindi—Talala (1,800'), ♂ 2, ♀ 1.
 Kashmir—Tral, ♂ 1, ♀ 2; Arapul Valley, ♂ 2, ♀ 1; Saradallu, ♂ 3, ♀ 1;
 Chasma Shahi, ♂ 2, ♀ 2 (juv.)
 Chakdalla, ♂ 1, ♀ 1.

- (29) MILLARDIA MELTADA, Gray.
The Soft-furred Field Rat
 (Synonymy in No. 1)

Bhattu Hissar, ♂ 2, ♀ 1.

- (30) MUS GERBILLINUS, Wroughton
The Sind Wild Mouse
 (Synonymy in No. 24)

Rawalpindi—Chak Lala, ♂ 1; Ar A ♀ 1.
 Campbellpore—Chakdalla, ♂ 1; Jhalar, ♀ 1.
 Bhattu Hissar, ♂ 2.

(31) MUS HOMOURUS, Hodgs.

Himalayan House Mouse

(Synonymy in No. 15)

Kashmir—Srinagar, ♂ 2; Sardallu, ♂ 2.
Bhattu Hissar, ♂ 1.

(32) MUS URBANUS, Hodgs.

The Nepal House Mouse

(Synonymy in No. 15)

Bhattu Hissar, ♂ 1, ♀ 1.

(33) APODEMUS S. GRISEUS, True.

The Kumaon Long-tailed Field Mouse

(Synonymy in No. 15)

Kashmir—Saradallu, ♂ 10, ♀ 7; Sutor, ♀ 1.

(34) ACOMYS FLAVIDUS, Thos.

The Indian Spring Mouse

1917. *Acomys flavidus*, Thomas. J.B.N.H.S., xxv, p. 205.
Kohistan (Sind), ♀ 1.

(35) MICROTUS (HYPERACRIUS) WYNNEI, Bland.

The Murree Vole

1880. *Arvicola wynnei*, Blandford, J.A.S.B., xlix, pt. 2, p. 244.

1889. *Microtus wynnei*, „ „ Mamm. No. 303.

Saradallu (Kashmir), ♂ 1, ♀ 5.

(36) LEPUS DAYANUS, Blandf.

The Sind Hare

(Synonymy in No. 3)

Rawalpindi—Chak Lala, ♂ 1, ♀ 1; Toupri, ♂ 4, ♀ 2.

(37) OVIS VIGNEI CYCLOCEROS, Hutton.

The Afghan Urial

(Synonymy in No. 32)

Taliala, (1,800'), Rawalpindi, ♂ 1, ♀ 1.

Nilo, N. of Jhelum, ♂ 2.

(38) GAZELLA BENNETTI, Sykes.

The Indian Gazelle

(Synonymy in No. 1)

Rawal, ♂ 1.

INDIAN DRAGONFLIES

BY

MAJOR F. C. FRASER, I.M.S., F.E.S.

Part XXIV

With 2 plates and 5 text figures

(Continued from page 171 of Vol. XXXI)

Genus—BURMAGOMPHUS Williamson.

Burmagomphus—Will., Proc. U. S. Nat. Mus., vol. xxxiii, pp. 298-301, 1908.

Laid., Rec. Ind. Mus., vol. xxxiv, p. 399, 1922.

A genus of medium-sized Gomphine dragonflies, coloured black, vividly marked with citron yellow or greenish yellow. Species studied :—

B. pyramidalis Laid., *B. sivalikensis* Laid., *B. siamensis* sp. nov., *B. laidlawi*, Fras., *B. cauvericus* sp. nov., *B. hasimaricus* sp. nov., and *B. V-flavum* sp. nov. In addition to these, I have the able descriptions of *B. williamsoni* (Will.) nom. nov., *B. insularis* Laid., *B. jacobseni* Ris, and the somewhat brief description of the geno-type, *B. vermiculatus* (Mart.), to assist me in a study of the genus as a whole.

Of the eleven species quoted above, seven are found within Indian limits, whilst *siamensis* comes from Siam, as the name implies, *vermiculatus* from Tonkin, *insularis* from Borneo and *jacobseni* from Java.

B. siamensis and *V-flavum* are known only from females, most of the remainder from both sexes.

The thoracic markings denote a cleavage into two groups, which is further supported by the character of the genitalia. The first of these groups,—*vermiculatus*, is well-defined, and especially characterized by the fusion of the upper half of the antehumeral thoracic stripe with the lower end of the humeral, so as to form a single vermiform stripe, or one with a bayonet-twist at its middle. The lower half of the antehumeral stripe is absent, whilst the upper half of the humeral is represented by a small isolated spot.

In the second group, the thoracic markings are by no means so uniform and vary widely in the species; the hamules are built on the same plan as in the first group, although differing more widely than is found to be the case in that group.

Possibly some of the forms in group *vermiculatus* are mere races of that species, but as we do not know the shape of the genitalia of the geno-type, I prefer to give all specific rank. I am justified in this by the example afforded by *hasimaricus*, which is almost identical in its markings to *Pyramidalis*, which again is almost identical in its markings to *vermiculatus*, and yet the genitalia of *hasimaricus* is widely different to that of *pyramidalis*. I had actually classed the former as a race of the latter, until I discovered this important difference.

All species, so far as known, are jungle habitants, frequenting mountain streams in ravines, in montane or submontane areas of the tropics of Southern Asia. Males are found resting on stones in midstream or on rocks or foliage beside these waters. Females rest high in trees, descending only for purposes of ovipositing. Larvæ are of the torpedo-shape, resembling rather closely those of *Onychogomphus*; all breed in running water.

Wings: reticulation rather open; nervures between *Mi-iii* and *Miv* in forewing 2, only 1 in the hind; pterostigma braced, rather swollen, about one-fourth the length of distance between node and distal end of pterostigma, rather longer and narrower in the hind wing; trigones, hypertrigones and subtrigones all entire, trigone in forewing with costal side shortest, basal and distal sides nearly equal, the latter usually slightly angulate at its middle; trigone in hind wing rather elongate, its basal side shortest, costal and distal




Fig. 1. Wings of *Burmagomphus pyramidalis* Laid. ♂

sides subequal; discoidal field in forewing 2-celled to beyond level of node; *Miii* and *Miv* distinctly sinuous in forewing; *Cui* and *Cuii* in hindwing parallel nearly to border of wing, dilated in the fore; *Cuii* in forewing subpectinate, 3 rows of cells between it and border of wing; 1 to 2 rows of postanal cells in forewing, 3-4 in the hind, where the first postanal cell is entire and extends basad to middle of subtrigone; only 2 rows of cells between *Mi* and *Mia*; only 1 cubital nervure in all wings; arc at the 2nd antenodal nervure; sectors of arc parallel or slightly divergent from origin; the 1st and 4th the primary antenodals; no basal incomplete antenodal nervure; anal triangle 3-celled; base of wing slightly excavate; tornus angulate.

Head moderately large, face sloping, frons rounded, occiput variable, notched, rounded or spined at centre.

Legs moderately long, hind femora extending to slightly beyond the apical border of 2nd abdominal segment, armed with closely set, small, evenly-sized spines.

Abdomen tumid at base, thin and cylindrical as far as apical half of segment 7 from where it dilates considerably as far as 9, but without lateral wings, segments 7 to 10 diminishing gradually in length, 10 very small. Oreillets moderately large.

Anal appendages resembling those of *Gomphus*, of equal length and almost equally divaricate, both superior and inferior black.

Genitalia. Lamina depressed, arched or notched; anterior hamules short stilletes; posterior hamules broad, flattened or curled plates with an anterior robust spine, and often with 1 or more accessory marginal spines; lobe prominent, globular or funnel-shaped.

Geno-type.—*B. vermiculatus* (Mart.).

Group I. *vermiculatus*.

Burmagomphus pyramidalis Laid., Rec. Ind. Mus., vol. xxiv, pp. 400 and 401, 1922; Fras., Rec. Ind. Mus., vol. xxvi, p. 476, 1924.

Male: Abdomen 30 mm. Hindwing 23-24 mm.

Head: labium yellow, margined with black; labrum greenish-yellow with its anterior border narrowly, and base, more broadly black, and a short medial prolongation of black running from the base line; bases of mandibles black, the genæ, anteclypeus and postclypeus black, the latter marked with a median spot and a lateral triangular larger spot greenish-yellow; frons greenish-yellow, its base above black; vertex and occiput black, the latter slightly concave.

Prothorax black marked with a greenish-yellow spot on each side.

Thorax black marked with greenish-yellow as follows :—a complete mesothoracic collar, a small spot in the alar sinus above, a sinuous dorsal stripe running from near the alar sinus above, down and out to the second pair of legs. This stripe is formed by a union of an antehumeral with a humeral stripe, the upper antehumeral portion is very narrow and broadens abruptly at its middle where it takes a bayonet turn outwards to fuse with the humeral portion. A small upper humeral spot representing the upper part of an obsolete humeral stripe ; the sides broadly. The latter marked with a narrow, complete stripe on the postero-lateral suture and an incomplete one on the lower half of the anterior suture.

Legs black, coxæ yellow, as also the inner surface of the anterior pair of femora ; armed as for genus.

Wings hyaline often slightly tinted with saffron at the bases. Nodal index 10-12|13-9 10-14|12-11
10-9|9-9' 11-9|9-10, pterostigma yellow between black nervures ; 4 rows of postanal cells in hindwing.

Abdomen black marked with yellow as follows :—segment 1 with a triangular mark on dorsum at apex and a broad baso-lateral spot, 2 with a lanceolate dorsal stripe not quite reaching the apical border, and a very broad spot on the sides, 3 to 8 with narrow basal rings, rather broader on 7 and very narrow on 8 where however it is produced laterally as an elongated spot on the basal half of segment, segment 3 has also a large baso-lateral triangular spot confluent with the dorsal ring, segment 8 often with an additional apico-lateral spot, 9 has nearly its apical half yellow, but laterally this mark much narrower and confluent with a ventral bordering of yellow ; at its apex, at the middorsum a very robust spine, also yellow. (This spine has been mentioned by Martin in his short description of *vermiculatus*, but has escaped the notice of Laidlaw and of Williamson ; it appears to be a group character of *vermiculatus*.) Segment 10 entirely black.

Anal appendages black, superiors as long as segment 10, divaricate, tapering to a point, to the outer side of which is a stunted spine which gives a bifid appearance to the apices when viewed a little to one side. Inferior deeply cleft into two widely divaricate branches, which are rather longer than the superiors, and taper to a fine point, the apex of which curls rather abruptly upward.

Genitalia. Lamina depressed, broadly arched ; anterior hamules short, stillette-shaped with an inconspicuous backwardly-turned point ; posterior hamules very robust, broad, projecting markedly, pyriform, with a robust forwardly directed spine at apex ; lobe tumid, rounded, shallowly grooved in front.

Female. Abdomen 33 mm. Hindwing 27 mm.

Similar to the male, differs as follows :—The basal ring on segment 2 more regular and the yellow on its sides more extensive ; on the sides of segment 3 an elongate yellow spot, not confluent with the basal triangular spot ; segment 9 with only its apical fourth yellow and 10 with a pair of small dorso-apical spots.


Anal appendages small, conical, black, the protuberance between them yellow. Nodal index usually slightly higher. Vulvar scale glossy back, its apical third broadly bifid, the whole scale broadly triangular.

Distribution. Confined to Western India from south-west of the Deccan to Malabar, Coorg, Nilgiris, Kanara and Poona. Specimens from the latter place have the ground colour bright citron yellow, whilst from the moister zones of Malabar and Coorg, specimens have the ground colour more grass-green. The measurements given are of those from Poona, from where the type comes, those from the Western Ghats are from 3 to 4 mm. larger. They emerge from May to July in the latter parts, and from August to September at Poona. At the latter place I took most specimens settled fairly high up on evergreens near the river-side, whilst in the Western Ghats they are rarely seen except settled on rocks in mid-stream, and then only during bursts of sun shine. It is a shy, jungly creature, but not uncommon. Type in the Indian Museum.

Burmagomphus williamsoni Will. nom. nov. *Burmagomphus vermiculatus* Will., l.c., pp. 301, 303, Figs. 27, 28 and 29. 1908.

Male : Abdomen 28 mm. Hindwing 23 mm.

Closely resembles *pyramidalis* Laid., and *vermiculatus* Mart., for the latter



Lateral views of:—1. *Burmagomphus siamensis* sp. nov. ♀ 2. *Burmagomphus vermiculatus* (Mart). ♂ 3. *Burmagomphus pyramidalis* Laid. ♂ 4. *Burmagomphus sivalikensis* Laid. ♂ 5. *Burmagomphus laidlawi*. Fras. ♂ 6. *Burmagomphus cauvericus* sp. nov. ♂ 7. *Burmagomphus flavum* sp. nov. ♀

of which Williamson mistook this species. Differs from the former as follows :—

Nodal index $\frac{8-10}{10-8} \frac{10-9}{8-8}$; labium not bordered with black; labrum black with two large transversely elongate greenish spots at base; mesothoracic collar slightly interrupted; segment 2 with a long trilobed dorsal stripe; narrow middorsal lines on segment 3 to 5 confluent basad with the basal rings; segment 6 with a fine triangular dorsal spot at base; genitalia specialized, posterior hamules broader, the spine more forward and smaller, and in addition two smaller spines on the outer border, lobe more prominent and more deeply notched.

Distribution. Burma. Described by Williamson from 3 males collected by Mr. R. A. Earnshaw. Type in Williamson collection. Female unknown.

Burmagomphus hasimaricus sp. nov.

Male: Abdomen 32 mm. Hindwing 27 mm.

Closely similar to *B. pyramidalis* Laid., but differs markedly by its genitalia. I note also the following differences :—midlobe of labium entirely black, labrum black with two elongate basal spots as in *williamsoni*, a mere vestige of the central spot on epistome and the large lateral spots against the eyes on the same structure entirely absent; occiput and vertex entirely black, the large yellow spot here entirely absent, the borders of the occiput flat, not curled up as in *pyramidalis*; the mesothoracic collar very finely divided; the mid-lateral black stripe longer and nearly confluent with the short oblique stripe which runs towards the hinder stripe above; pterostigma darker, pale brown; a trilobed dorsal stripe on segment 2 and a fine middorsal line on 3 and 4, segment 8 entirely black; anal superior appendages longer and finer, and without the outer spine, the outer side bevelled for its apical two-thirds, inferior appendage with the apices of branches scarcely upturned. Posterior hamules very bulky, the base a little constricted, the apical portion produced squarely back with distinct angles, its anterior spine more robust and directed more forward.

The female differs less in some respects and more in others than does the male from *pyramidalis*. The labrum is similar to *pyramidalis* and the large triangular lateral spots of epistome are present. The median spot on this organ is present as two fine transversely linear spots. The occiput is entirely black. The thorax is much less broadly black than in the male and approximates to *pyramidalis*, the lateral stripes being very fine and the median short and not nearly reaching to the upper segmental stripe.

The dorsal surface of segment 2 is entirely without the middorsal stripe and has only a basal ring and the sides broadly yellow, segments 3 to 6 have the basal rings produced slightly apicad, and in addition to the lateral elongate spot on segment 2, there is a similar, but smaller one on 3. Segment 8 is unmarked as in the male. The protuberance between the anal appendages is black. Lastly the vulvar scale is longer and is cleft very deeply, ending in two long fine spine-like processes, very different to what is seen in *pyramidalis*.

Distribution. One male and two females from Mr. H. V. O'Donel, from Hasimara, Duars, Bengal. Mr. O'Donel in sending me these specimens remarks that the species is very close to *pyramidalis*, but differs by having a dorsal stripe on the abdomen. The differences in the genitalia of both sexes are however sufficient to constitute a good species. Type at present in the Fraser collection.

Burmagomphus siamensis sp. nov.

Female: Abdomen 27 mm. Hindwing 24 mm.

It is convenient to describe this new species here, as it forms a connecting link between *williamsoni* and the genotype *vermiculatus*, and because, on account of its proximity to Lower Burma, it may possibly extend to Burma, like many other Siamese species.

It resembles *williamsoni* closely, but differs as follows :—Nodal index $\frac{9-12}{9-8} \frac{12-9}{8-8}$; labrum greenish-yellow, its base and anterior border bordered with black and its centre traversed by a narrow black line cutting the yellow into two very large spots; mid-spot and lateral spots on epistome small and widely separated by the black; occiput yellow, raised into a long prominent median spine at its hinder border and with a smaller black spine at either end against the eyes but not quite on the free border. Thoracic markings similar but the

mesothoracic collar complete. Abdomen with a fine middorsal stripe on segment 2 continued as a very fine line on 3 and 4; a complete lateral broad stripe on 3 finely divided by the jugal suture, a similar on segment 4, but broadly broken at the level of the jugal suture and incomplete at the apical end, the basal rings on 5, 6 and 7 produced laterally for a short distance, and on 5 a medial lateral linear spot, 8 with two basal dorsal lunules narrowly connected over the dorsum, 9 with its apical border finely yellow. Vulvar scale broadly triangular, cleft to its base into two triangular processes.

Distribution. Bangkok, Siam. A single female in the Fraser collection. This specimen may of course be the female of *williamsoni* but the yellow occiput and the restricted markings on the face seem to separate it from that species. The genotype *vermiculatus* differs from it by its larger size,—abdomen 30–33 mm., hindwing 24 mm.; the nodal index is higher, 14 to 15 antenodal nervures in forewings, and 9 to 11 postnodals; 10 antenodals in hindwing and 9 to 11 postnodals. The lateral spots on the epistome are absent as in *hasimanicus*. The mesothoracic collar is interrupted; the first lateral black stripe is said to be forked above, but I think that this is an error, the second being meant, as in all other species. Lastly there are no lateral markings on any of the segments after 2. *B. vermiculatus* comes from Tonkin.

Group *laidlawi*.

In this group I place all those species in which the antehumeral and humeral stripes are not combined to form an oblique sinuous stripe and in which the male does not bear a dorsal spine at the apex of segment 9.

Burmagomphus laidlawi Fras. *Gomphus* sp. Fras., Rec. Ind. Mus., vol. xxiv, p. 419, 1922; id. ib., vol. xxvi, pp. 475, 476, 1924.

Male: Abdomen 33 mm. Hindwing 27 mm.

Head: labium with midlobe black, lateral lobes greenish white; labrum black marked with two transversely oval yellow spots; bases of mandibles yellow; ante- and post-clypeus black with a small linear spot, variably present, at lower border of latter; frons bright greenish yellow, rest of head black. Occiput slightly raised at the middle, fringed with long black hairs. Eyes bottle green.

Prothorax black with a large lateral spot and an anterior collar of yellow.

Thorax marked with greenish yellow or greenish white as follows:—a complete mesothoracic collar, oblique antehumeral stripes extending from alar sinus but not meeting the mesothoracic collar, a minute upper humeral point and the vestiges of a lower humeral stripe, the alar sinus and the whole of the sides, which latter are marked with two narrow black stripes on the sutures. These two stripes occasionally confluent above and below, thus enclosing an elongate yellow spot, but usually only the upper part of stripes are confluent. In one specimen from South Kanara, the stripes are so confluent as to almost obliterate the included yellow, and the labial spots are also absent in the same specimen.

Wings hyaline, slightly tinted with yellow at the base; 4 rows of postanal cells in hindwing; pterostigma dark brown, over 4 to 5 cells; nodal

index $\frac{13-16}{11-10} \frac{15-12}{10-12} \frac{10-14}{10-9} \frac{12-10}{10-11}$

Legs black, anterior femora yellow internally, hind femora armed as for genus.

Abdomen black marked with yellow as follows:—segment 1 with a dorsal spot extending from base to apex, and its sides broadly, 2 with a bilobed middorsal stripe extending from base to apex, and the sides broadly, the subdorsal black descending as a vertical stripe behind the oreillets, segment 3 with the middorsal carina finely yellow, but usually only at the base, and a large baso-lateral spot, segments 4 to 6 with basal dorsal triangular spots and baso-lateral lunules confluent with the former, 7 with a broad basal ring occupying about one-fourth the length of segment, 8 unmarked or with a minute basal dorsal triangular spot, and in some with a still smaller apical spot, 9 with its apical half yellow, this colour sometimes extending irregularly along the dorsal crest, 10 unmarked.

Anal appendages black, both superiors and branches of inferior markedly divaricate, the inferior more so than the superiors. The apices of superiors

long and thin and twisting in and a little down. Branches of inferior curling upward. Slightly shorter than segment 10.

Genitalia: lamina very narrow, very depressed; anterior hamules short stillettes with an inconspicuous hook at apex; posterior hamules very robust, projecting, squared and with a robust spine on the anterior corner projecting forward, a smaller one on the hinder angle projecting vertically. In some specimens there are 2 or 3 such posterior spines, but usually only one; lobe very tumid and prominent, rounded, deeply cleft.

Female: Abdomen 32 mm. Hindwing 29 mm.

Very similar to the male, markings better defined. The humeral spot larger and continued as a broken line below, or similar to the male. Often a small spot on each side of the postclypeus. A very prominent horn on each side of head just behind the lateral ocelli; occiput simple, nearly straight. The baso-lateral spot on sides of segment 3 continued after a slight interval, by a broad lateral stripe which extends nearly to apex of segment; 8 with a very fine basal yellow ring, the yellow on 9 more restricted, 10 with its apical border lined with yellow. The superior anal appendages small, conical, pointed, black with a yellow spot above. Vulvar scale very broad, of even width, slightly notched at its middle, glossy black. (The Kallar female has segment 10 unmarked, and the horns on vertex are poorly developed). Wings in all adult females are enfumed pale brown, teneralis are rather deeply saffronated at the base.

Distribution. Western Ghats of India only. The original pair were taken *in cop*, flying along the bed of the Kallar river, Nilgiris, 2,000 ft., August 13, 1922. A female was taken at Gudalur, Nilgiri-Wynaad, 3,500 ft., and several males in Coorg and South Kanara during September 23. Its habits are similar to *pyramidalis*, with which it is occasionally found in company. Type in the British Museum.

***Burmagomphus cauvericus* sp. nov.**

Male: Abdomen 35 mm. Hindwing 28 mm.

Very similar to *laidlawi*, differs as follows:—Midlobe of labium greenish, narrowly bordered with black; a small triangular spot of greenish yellow on each side of postclypeus; the upper humeral spot usually entirely absent, or if present, then a mere point; lateral markings on thorax similar to those of *pyramidalis*, the stripe on the first suture is incomplete above, but runs up higher than it does in *pyramidalis*, and is almost confluent with a short oblique stripe which runs back from below forewing to join the upper part of the stripe on the second suture, enclosing a small spot above. In some specimens this short stripe is so broadly confluent, that the stripe on the second suture appears to be expanded above, and marked with a tiny point of citron yellow. The hinder border of metepimeron bordered above narrowly with black. Segment 2 with a dorsal lanceolate spot extending to about the middle of segment 10 with a fine vestigial line along its middorsal ridge. Genitalia; posterior hamules less broad than in *laidlawi*, the anterior spine smaller and two smaller spines situated close together at hinder angle. Lobe smaller, less deeply notched.

Female: Abdomen 35 mm. Hind wing 31 mm.

Very similar to the male, differs as follows:—The black markings on sides of thorax more restricted, so that the enclosed spot above the front of second lateral suture is very large and the black stripe on the suture at this level is reduced to a fine line, the basal ring on segment 3 produced laterally as a broad triangular spot, followed by a large elongate lateral isolated spot, segment 4 with a similar lateral but smaller spot. Rest as for male. Segment 8 in both sexes unmarked. Vulvar scale more broadly notched so that its border exhibits two broad triangular processes. Anal appendages entirely black.

Distribution. Coorg only, along the banks and tributaries of the river Cauvery. This handsome species may be immediately recognized by the combination of the anterior thoracic markings of *laidlawi* with the lateral thoracic markings of *pyramidalis*. The occiput of the female is notched in the middle and rounded on either side of the notch. The horns on the vertex are shorter and with a broader base and are situated a little to the outer side of the hinder ocelli, instead of directly behind them. Type in the British Museum.

Burmagomphus sivalikensis Laid. l. c., pp. 401, 402, 1922.

Male. Abdomen 34 mm. Hindwing 25 mm.

Head: labium pale greenish white, the midlobe narrowly bordered with black; labrum black marked with two very large transversely elongate greenish-white spots; bases of mandibles coloured similarly; anteclypeus black with a median linear spot of palest green and a large spot on each side the same colour; frons black in front, greenish above, its base narrowly black, this colour prolonged forwards at the middle as a very short triangle; vertex black; occiput greenish white, slightly notched at its middle and convex on either side, fringed with pale hairs.

Prothorax black with an anterior collar, a large lateral spot and a small median dorsal spot citron yellow.

Thorax black marked with greenish-yellow as follows:—a mesothoracic collar finely divided by the dorsal black carina, very oblique long antehumeral spots well separated from the collar below and the alar sinus above, a long irregular humeral stripe constricted just below its upper part and continued below on to the base of the middle pair of legs. Laterally two narrow black lines on the lateral sutures, rather narrower than the black stripe separating the humeral stripe from the yellow stripe following it. The first lateral stripe is sinuous above and the second constricted above and angulated backwards below. Lastly there is a fine black line on the upper and back part of metepimeron.

Legs black, the inner surface of the anterior pair of femora yellow, armature of hind femora as for genus.

Wings hyaline; pterostigma pale yellowish brown between thick black nervures, over 3 to 4 cells; nodal index

7-12	13-8
6-9	9-7

Abdomen black marked with pale citron yellow as follows:—segment 1 with a triangular spot on dorsum, its base against the apical border of segment, segment 2 with a lanceolate middorsal stripe extending from base to apex, sides of these two segments broadly, the second segment with an extension of black behind the oreillet and a narrow black apical ring, segments 3 to 7 with a narrow basal ring, rather broader on 3 and, on all, constricted at the middorsum by invasion of black, 3 has also an elongate lateral spot each side, segment 8 unmarked, 9 with a very large triangular spot on its apical three-fourths, more extensive on dorsum than on sides, 10 unmarked.

Anal appendages black, as long as segment 10, superiors nearly parallel, pointed at apex and with a stout stunted spine near the apex on outer side as in *pyramidalis*, which gives a bevelled appearance to this part of appendage. Inferior deeply cleft into two very widely divaricate branches with upturned apices.

Genitalia: lamina slightly raised, hood-like, deeply cleft; anterior hamules short stillette-shaped organs; posterior broad and robust, longer and narrower than in other species and tapering outwardly to a blunt point, from the apex of which springs a long fine spine; lobe projecting, shaped like the spout of a milk-jug.

Female unknown.

Distribution. Dehra Dun and Hasimara, Duars, Bengal. The above description differs in several respects from that of Dr. Laidlaw's, which was made from an old broken and faded specimen. The present description has been made from a fresh specimen sent me by Mr. H. V. O'Donel, who took it at Hasimara; its ground colours are a beautiful greenish-yellow, in strong contrast to the dirty orange yellow of the type from Dehra Dun. The latter in the British Museum.

The character of the genitalia, as well as its markings, will easily distinguish this species from others of the genus.

Burmagomphus V-flavum sp. nov.

Female: Abdomen 31 mm. Hindwing 30 mm.

Head: labium pale greenish white; labrum black marked with two large basal transversely oval citron yellow spots; bases of mandibles yellow; anteclypeus black; postclypeus glossy black, narrowly bordered below with citron yellow and with a large spot of the same colour on each side against the eyes; frons black with a broad citron yellow stripe above which is cut by a median black line into two large oval spots; vertex black; occiput yellow

encircled with black, the hinder border narrowly black, behind broadly yellow, slightly concave.

Prothorax black broadly marked with citron yellow as follows:—the whole of the posterior lobe, a small geminate spot in front of it, a very large lateral spot nearly confluent with the yellow on the posterior lobe, lastly a broad anterior collar.

Thorax black marked with citron yellow as follows:—a complete mesothoracic collar slightly notched behind at its centre, antehumeral and humeral oblique stripes which converge and become broadly confluent below so as to form a large yellow V-shaped spot on each side of dorsum of thorax. A small spot in the front angle of the alar sinus. Laterally broadly yellow with an incomplete lateral black stripe on the first suture extending up as far as the spiracle and a narrow complete black line on the second suture.

Legs black, the inner sides of anterior pair of femora greenish-yellow; hind femora with a row of 9 to 10 short robust spines gradually lengthening distad, and numerous smaller spines at the proximal third; overlapping the base of segment 2 when apposed.

Wings: hyaline; pterostigma pale yellow between black nervures, rather large, more than one-fourth the length of distance between node and distal end of pterostigma, over 4 to 5 cells; nodal index $\frac{11-14}{11-11} \frac{13-11}{12-11}$; *Cui* and *Cuii* rather divergent in hindwing from a little more than half-way to wing border; no basal incomplete antenodal nervures.

Abdomen black marked with citron yellow as follows:—segment 1 with a small subdorsal apical black spot, otherwise entirely yellow, 2 with a broad middorsal trilobed stripe and the sides very broadly, 3 with a basal ring confluent with a fine dorsal stripe which extends to apex of segment, and laterally with a very broad stripe nearly interrupted at the jugal suture and not extending to apex of segment, 4 to 7 somewhat similar but the sides broadly black, only segments 4 to 5 having a small oval lateral yellow spot, 8 with a small triangle of yellow on its dorsum at the base, 9 with its apical half bright citron yellow, 10 with its apex narrowly bordered with three triangular spots of yellow.

Anal appendages short, conical, black. Vulvar scale very short and broad, less than one-fourth the length of segment 9, very shallowly notched at apex.

Distribution. Maymyo, N. Shan States, Upper Burma, June 15, 1924, two females collected by Col. F. Wall, I.M.S. This new species is not quite typical of genus *Burmagomphus*, as the pterostigma is too large and the legs too short and the armature of the hind femora differs a little. The large swollen pterostigma recalls that of *Cyclogomphus* but there is no incomplete basal antenodal nervure present in any of the wings of the two specimens. The formation of yellow Vs on the dorsum of thorax, by the fusion of the humeral and antehumeral stripes below, is sufficiently characteristic to distinguish this species from any other Indian species.

Genus: *Platygomphus* Selys.

Rather large sized Gomphine dragonflies characterized by the anal border of hind wing barely excavated and the anal angle subrounded, colours bright yellow narrowly marked with black. Only two species are known, both from within Indian limits and both very similar in appearance. They are found in the plains, in cultivated areas. Mr. T. B. Fletcher states that at Pusa, they are often seen settled, sunning themselves on the broad green leaves of cannas, in gardens.

Platygomphus martini a species described by myself in the Pusa Memoirs, turns out to be an *Indogomphus*. The type was returned to Pusa and a year later I received other specimens from the Duars, and failing to recognize my own species redescribed it again as *Burmagomphus duarensis* in the Records of the Indian Museum. By the great length and armature of its hind femora and curious proportion of its end abdominal segments, it is clearly an *Indogomphus*.

As Dr. Laidlaw remarks, *Burmagomphus* appears to shade off into *Gomphus* on the one hand and into *Platygomphus* on the other; this species appears to share characters of both these genera but more especially those of *Indogomphus*. (In Part XX of Indian Dragonflies this Journal, it is to be noted that in the explanation of the plate accompanying that part, '*Indogomphus duarensis*'

has been put instead of '*Indogomphus martini*,' although it is correctly designated in the text. I trust that this long chain of errors for which I have been responsible has now come to an end.




Fig. 2. Wings of *Platygomphus dolabratus* Selys., Male.

Wings : reticulation close, 2 nervures between *Mi-iii* and *Miv* in forewings, only 1 in the hind ; pterostigma braced, that of hindwing rather more elongate than the fore, equal in length to about one-fourth the distance between the node and distal end of pterostigma ; trigone in forewing with costal and basal sides equal, the distal side slightly longer ; in the hind trigone, the distal side considerably longer than the other two and a little sinuous, the basal shorter than costal side ; all trigones, hypertrigones and subtrigones entire ; discoidal field 2-celled nearly as far as node, then considerably dilated as far as termen ; *Miii* and *Miv* distinctly sinuous in the forewing ; *Cui* and *Cuii* in both wings parallel to border of wing ; *Cuii* not pectinate, 3 rows of cells between it and border of wing ; 3 to 4 rows of postanal cells in hindwing, *only 1 in the fore* ; first postanal cell entire, extending basad only to middle of subtrigone ; only 2 rows of cells between *Mi* and *Mia* at and near origin ; only 1 cubital nervure in all wings ; arc at the 2nd antenodal nervure ; sectors of arc parallel ; the 1st and 5th the primary antenodals ; no basal incomplete antenodal nervure ; anal triangle very narrow, 1-celled ; base of wing barely excavate ; tornus slightly rounded.

Head rather small, narrow, face sloping, frons rather rounded, occiput flat, straight.

Legs moderately long, hind femora extending to slightly beyond apex of segment 1 and armed with small, closely-set and numerous spines of even size.

Abdomen tumid at base, then thin and cylindrical as far as base of segment 7, then greatly dilated as far as segment 9, but not definitely winged as in *Ictinus* ; segments 7 to 10 diminishing gradually in length.

Anal appendages resembling those of genus *Gomphus*, of almost equal length and almost equally divaricate.

Genitalia ; lamina depressed, notched ; posterior hamules elongate narrow and subcylindrical, projecting perpendicularly from body, apex with a sharply curled spine ; lobe very tumid, large and projecting ; oreillets moderately large.

Genotype, — *Platygomphus dolabratus* Selys.

Platygomphus dolabratus Selys, Bull. And. Belg. xxi. (2) p. 44, (1854) ; id. Mon. Gomph., p. 113. (1857) ; Will. l.c., p. 303 (1901) ; Laid. l.c., pp. 398 and 399 (1922).

Male : Abdomen 41 mm. Hindwing 30 mm.

Head : labium, labrum and face bright yellow, the labum with a small brown median stripe projecting down from base, the postclypeus with a broad

transverse black stripe, the upper border of which is slightly convex and encroaches on the anterior surface of frons, the lower produced into two quadrate projections one on each side near the middle line; frons entirely yellow above and for greater part of front; vertex black with a large round spot of yellow behind the ocelli; occiput bright yellow, its border very slightly convex at the middle. Eyes bottle green.

Prothorax yellow at the sides, black on the dorsum except for a pair of twin spots at the middle of hinder lobe and the whole of the prominent anterior collar.

Thorax yellow marked with black as follows: the dorsum, where the black encloses a complete bright yellow mesothoracic collar and two broad triangular antehumeral spots of duller yellow. Behind these a moderately broad and even, humeral, bright yellow stripe which turns in a little at its upper part. On the sides a short fine vestige of a black line on the first lateral suture and a complete very fine black line on the second suture.

Wings hyaline, a little tinted with yellow occasionally; pterostigma yellow between black nervures, braced strongly, over 3 cells; nodal index 9-13|14-9

9-10|10-10

Legs yellow marked with black, a narrow distal streak on the outer side of the hind femora and the whole of the outer sides of the other two pairs, all the tibiae black. Hind femora with a row of short robust spines.

Abdomen yellow marked with black as follows:—segment 1 with a small subdorsal spot and a fine black line running obliquely back and outward from it, segment 2 with a broad subdorsal stripe extending the whole length of segment and enclosing an acorn-shaped spot the stalk of which is directed towards the base of segment, segments 3 to 7 black on dorsum; in adults this black encloses a lanceolate middorsal spot, broader at base and pointed towards the apical border; basad the spot is confluent with a basal ring in semi-adult specimens. (In all the specimens from Pusa, the black is present as an apical ring with a subdorsal branch running towards the base of segment on either side, enclosing a prolongation of the ground colour which runs from a basal ring). Segments 8 and 9 are dark brown above changing to yellow on the sides, 8 has a fine middorsal line at the base, segment 10 paler brownish yellow.

Anal appendages yellow or yellowish-brown, bordered with black, as long as segment 10; the superiors flat, divaricate, inner sides straight, outer right-angled with prominent outer point, apex pointed as seen from above, slightly bifid as seen in profile. Inferior deeply forked, its branches rather more divaricate than the superiors, their apices turned up a little.

Genitalia: lamina slightly raised, rounded, broadly arched; posterior hamules very long, projecting almost perpendicularly from abdomen, subcylindrical, the apex strongly curled and ending in a hooked spine; lobe very bulky, subrounded, deeply cleft, its border somewhat sinuous as seen in profile.

Female: Abdomen 40 mm. Hindwing 33.

Coloured and marked similarly to the male. The end segments of abdomen less dilated, more broadly yellow along the sides, segment 10 yellow. Vulvar scale very short, notched. Occiput low, its free border a little tumid, a little rounded at the middle.

Distribution: Bihar and Bengal. Type in the Selysian collection, paratype females in the British Museum, and paratypes of both sexes in the Pusa and Fraser collections.

Platygomphus feae Selys, *Odonates de Birmanie*, Ann. Mus. Civ. Genov., xxx. (1890); Will. l. c., pp. 303 and 304 (1908); Laid. l. c. p. 398 (1922).

Male. Abdomen 35-37 mm. Hindwing 30-31 mm.

Head: labium, labrum, face and frons dull yellow, unmarked except for some brownish suffusion of face and a basal black line on frons which extends forwards as far as the crest at the middle; vertex black; occiput yellow fringed with greyish hairs, black behind, straight.

Prothorax yellow, black at middle of dorsum.

Thorax yellow marked with black as follows:—the dorsum similar to *dolabratus*, the black enclosing a broad antehumeral oval or subtriangular spot on each side and a moderately narrow humeral yellow stripe. Laterally a single

black narrow stripe on the second lateral suture which divides above to form a 'Y', the anterior arm of which runs towards the base of forewing.

Legs black, inner surfaces of femora pale yellow, armed as for genus, short.

Wings hyaline; pterostigma brown between thick black nervures, over 3 cells; nodal index $\frac{6-13}{10-9} \frac{12-8}{10-9}$; costal border brownish.




Fig. 3. Genitalia of :—1. *Burmagomphus sivalikensis* Laid, 2. *Burmagomphus hasimaricus* sp. nov. 3. *Burmagomphus cauvericus* sp. nov., 4. *Burmagomphus pyramidalis* Laid, 5. *Anisogomphus occipitalis* (Selys), 6. *Davidioides martini* Fras, 7. *Burmagomphus laidlawi* Fras., 8. *Temnogomphus bivittatus* (Selys), 9. *Indogomphus martini* (Fras.) 10. *Indogomphus longistigma* (Fras.), 11. *Platygomphus dolabratus* Selys, 12. *Anisogomphus caudalis* sp. nov.

Abdomen black marked with yellow as follows :—segment 1 with a middorsal stripe and the sides broadly, 2 with a lanceolate stripe on middorsum extending from base to apex, and the sides very broadly, 3 to 6 with a narrow basal ring prolonged along the sides below, and on 3 and 4 a small median dorsal oval spot, 7 with its basal half yellow, 8 to 10 coloured as in *dolabratus*.

Anal appendages black, equal in length to the 10th abdominal segment. Superiors markedly flattened, the outer border strongly angulate but the point situated much nearer apex than in *dolabratus*. Apex shortly acuminate so that seen in profile it appears to be bifid. Inferior appendage strongly forked, with the apices of its branches blunt and turned slightly up.

Female unknown.

Distribution : Bhamo Burma, in July and August. Type in the Selys collection. Distinguished from *dolabratus* by the face unmarked, the vertex without a spot of yellow, the abdomen more broadly marked with black, the anal appendages and legs black. The forking of the hinder stripe on sides of thorax and segments 5 and 6 without a dorsal spot. Lastly by the shape of the superior anal appendages, which appear more bifid in profile than those of *dolabratus*.

Genus—DAVIDIODES Fras.

A monotypic genus closely allied to *Burmagomphus*, colours black marked rather broadly with bright yellow. The single species known is characterized by its close venation, by its elongate trigone of hind wing traversed by a nervure and by its characteristic genitalia. The position of the genus is difficult to locate ; it belongs to *Gomphus*, and its venation, apart from the traversed trigones and its close character, closely resembles that of *Burmagomphus* and *Platygomphus* ; the genitalia is however widely different so that it cannot be placed in either of these two genera. The anal appendages resemble those of genus *Gomphus*, the superiors being yellow and the inferior black.

Wings : reticulation very close ; 2 nervures between *Mi-iii* and *Miv* in forewing, only 1 in the hind ; pterostigma braced, short, less than one-fourth the length of distance between node and distal end of pterostigma ; trigone of forewing entire, with costal and basal sides subequal, the distal rather longer ; trigone of hindwing traversed by a nervure running from the costal to distal side, very elongate, costal and distal side subequal, the basal much shorter ; hypertrigones and subtrigones entire ; discoidal field in forewing 2-celled as far as level of node ; *Miii* and *Miv* distinctly sinuous in the forewings ; *Cui* and *Cuii* parallel in the hindwing, divaricate near the border of wing in the fore ; *Cuii* not pectinate, 3 rows of cells between it and border of wing in forewing ; 1-2 rows of postanal cells in forewing, 4 in the hind, the first cell here entire, extending basad to a little beyond the middle of subtrigone ; two rows of cells between *Mi* and *Mia* at origin ; only 1 cubital nervure in all wings ; arc at or near the second antenodal nervure ; sectors of arc parallel at origin ; the first and sixth the primary antenodals ; no basal incomplete antenodal nervure ; anal triangle 4-celled ; base of hindwing markedly excavate ; tornus very acute.

Head moderately large, face oblique, frons slightly rounded, occiput a little concave.

Legs moderately long, hind femora extending on to segment 2 and armed with short, closely-set, evenly-sized spines.

Abdomen tumid at base, thin, long and cylindrical as far as segment 8 which with 9 is rather dilated laterally, 10 very small.

Anal appendages resembling those of genus *Gomphus*, of about equal length and almost equally divaricate, superiors yellow, inferior black.

Genitalia : lamina slightly raised ; anterior hamules very long, thin and flattened, nearly as long as the posterior hamules, which are very robust, broad at base, sinuous, tapering to a fine point, projecting markedly ; lobe bulbous with a markedly prominent, trowel-like lip.

Distribution. Western Ghats of India. Habits similar to those of *Burmagomphus* but found at lower levels.

Davidioides martini Fras., l. c., pp. 472 and 473 (1924).

Male : Abdomen 38 mm. Hindwing 33 mm.

Head : labium dirty yellow ; labrum black with a large basal spot on each side, widely separated, yellow ; bases of mandibles yellow ; anteclypeus yellow,

postclypeus black; frons black on lower part of front, yellow on upper part and above, its base above broadly black especially at the middle; vertex and occiput black, the latter simple, slightly concave.

Prothorax black marked with yellow as follows:—a minute spot on centre of posterior lobe, a large lateral spot on each side of middle lobe, a twin spot on the middorsum and an anterior collar.

Thorax black on dorsum, yellow on the sides, a slightly interrupted yellow mesothoracic collar, slightly oblique antehumeral stripes separated widely from the collar below and not quite reaching the alar sinus above, a minute upper humeral spot. Laterally two narrow parallel black stripes on the sutures enclosing an equally narrow yellow line, lastly a narrow black posterior border to the metepimeron confluent above and below with the posterior black stripe. Yellow beneath.

Legs black, anterior and middle pairs of femora yellow within, armed with a row of very closely-set, short, even spines.

Wings hyaline; pterostigma blackish brown, braced, over 3 to 4 cells;

nodal index $\frac{12-14}{12-10} \frac{16-12}{10-13}$.

Abdomen black marked with yellow as follows:—segment 1 with a large spot on dorsum and the sides broadly, 2 with a middorsal fusiform spot not extending to base or apex, the ventral and lower part of apical borders narrowly, 3 to 7 with basal rings, narrow on 3 to 6, occupying the basal half 7, where also there is a small apico-lateral spot on each side, 8 to 10 unmarked.

Anal appendages. Superiors as long as segment 10, bright yellow, narrowly black at base, widely divaricate, conical, tapering to a point, with 5 or 6 minute black spines on the ventral surface. Inferior deeply cleft, its branches widely divaricate and equally so as superiors but shorter, curled slightly up at apex, black.

Genitalia described under generic characters.

Distribution. Western Ghats of India. Type in the Fraser collection from Kunnoth, N. Malabar, May 18, 1923.




Fig. 4—Wings of *Anisogomphus occipitalis* (Selys.) Male

Genus—ANISOGOMPHUS Selys

A genus of moderately large Gomphine dragonflies coloured black vividly marked with greenish or citron yellow. The genus was founded by Selys to include two species,—*occipitalis* and *bivittatus*. The latter species has been removed to a new genus by Dr. Laidlaw on venational grounds. An examination of its genitalia, which differs from all other known species of the family, shows that this was fully justified. A second species,—*orites* has been described by the same author and a third new species is described below,—*caudalis*. All three species are found in North-East India at the foot of the

Himalayas, in montane and submontane tracts, frequenting and breeding in mountain streams. But little is known of their habits but these seem to be similar to those of *Burmagomphus*.

Wings: reticulation close; pterostigma braced, a little swollen, short, less than one-fourth the distance between node and distal end of pterostigma, of nearly equal size in the two wings; 3 to 4 nervures between *Mi-iii* and *Miv* in the forewing, 1 or 2 in the hind; all trigones, hypertrigones and subtrigones entire; trigone in forewing with costal side slightly shorter than proximal, and the proximal considerably shorter than the distal; in the hind the costal considerably shorter than the basal and but little shorter than the distal side; discoidal field in forewing 2-celled to level of node; *Miii* and *Miv* slightly sinuous in forewing; *Cui* and *Cuii* in both wings parallel to nearly as far as wing border; *Cuii* in forewing pectinate, 2 rows of cells between it and border of wing; 2 rows of postanal cells in forewing, 4 in the hind, where the first cell is entire, or divided, and extends proximad nearly as far as proximal end of subtrigone; 2 rows of cells between *Mi* and *Mia*; 1 or 2 cubital nervures in forewing, 1 in the hind; arc at or a little distad of the 2nd antenodal nervure; sectors of arc parallel for a very short distance from origin, then divaricate; the 1st and 7th the primary antenodals; no basal incomplete antenodal nervure; anal triangle 3-celled; base of wing markedly excavate; tornus nearly right-angled.

Head: face oblique, frons slightly rounded but prominent, occiput flat, simple.

Legs moderately long, hind femora extending to apex of segment 1, armed with 4 to 6 robust long spines, the longest of which are near the middle of limb; at the base numerous short spines.

Abdomen tumid at base, thin and cylindrical to as far as segment 7; 8 to 10 slightly dilated, the end of abdomen somewhat squared; segments gradually decreasing in length from 7 to 10.

Anal appendages. Superiors short, pale coloured, directed straight back, parallel, armed beneath with a basal robust spinal process. Inferior black, deeply cleft into two widely divaricate branches which project laterally from beneath the superiors.

Genitalia: lamina slightly raised, arched; anterior hamules small, stillette-shaped; posterior hamules very robust, bulky, sinuous, S-shaped, ending in a very short spine, surface minutely spined; lobe tongue-shaped, curling back, tapering, coated with long hairs; oreillets moderately large.

Genotype—*Anisogomphus occipitalis* Selys.

Anisogomphus occipitalis Selys.

Male: Abdomen 52 mm. Hindwing 30 mm.

Head. Labium yellow, midlobe blackish brown; labrum glossy black with two small, basal, greenish-yellow spots; bases of mandibles citron yellow, rest of face glossy black; frons broadly greenish-yellow; vertex and occiput black; eyes bottle green.

Prothorax black with a large lateral spot and a median twin, smaller spot on midlobe, citron yellow.

Thorax black marked with citron yellow or greenish-yellow as follows:—a slightly interrupted mesothoracic collar, slightly oblique antehumeral stripes which are confluent below with the middle of each half of the mesothoracic collar, an upper triangular spot and the vestiges of a humeral stripe well below this.

Laterally two narrow stripes of black on the sutures, the first of which curls forward above and sends a short prolongation back below at the level of the thoracic spiracle, the stripe on the second suture sends a short prolongation forwards and downwards below the level of spiracle.

Wings usually hyaline or more or less enfumed brown according to the age of specimens; pterostigma dark reddish brown, over 3 to 4 cells, lying between thick black nervures; 2 cubital nervures in the forewing, only 1 in the hind;

membrane almost obsolete, greyish; nodal index

10-15	16-12	12-18	16-13
12-12	12-10	12-11	13-11

Legs black, inner sides of anterior femora yellow, armed as for genus.

Abdomen black marked with yellow as follows:—segment 1 with a large triangular dorsal spot tapering basad, and its sides very broadly, 2 with a trilobed middorsal stripe extending the whole length of segment, the middle lobe of the stripe globular, and on the sides, the oreillets and a very large

apical spot which is prolonged along the ventral border towards the base of segment, 3 with a baso-lateral triangular spot and its dorsal carina finely but more broadly at the base, 4 to 6 finely yellow along the middorsal carina, 7 similar but more broadly so. 8 with a mere basal vestige of this stripe, 9 and 10 unmarked. The 10th segment globular, compressed dorsally at the apex.

Anal appendages. Superiors closely apposed at the base, slightly divaricate thereafter, equal to about half the length of segment 10, pale greenish white at base, reddish thereafter, conical, pointed. Seen from the side, a stout, long, blackish brown process with blunt nose is seen springing from beneath near the base of each appendage and closely apposed to each other. The whole appendage bears a striking resemblance to a fox's head, the superior parts being the ears, the apposed lower parts the face muzzle. Inferior appendage black, deeply cleft into two widely divaricate branches. So widely divaricate are these, that their inner borders are almost in a straight line. Each branch very robust, bluntly pointed and projecting widely from beneath the superiors.

Genitalia: lamina depressed, broadly, arched; anterior hamules short broad at base, tapering to a fine point which is curled back; posterior hamules very robust, thick, sinuously curved, S-shaped, apex shortly pointed and directed slightly forwards, the surface coated with minute spines and hairs, the latter especially thick near the apex; lobe-shaped like the spout of a jug, long and tapering, coated with long hairs.

Female: Abdomen 33 mm. Hindwing 32 mm.

Very similar to the male, differs as follows:—labium very occasionally unmarked (this character sometime shared by the male); thoracic stripe more evident (sometimes confluent with the upper spot, especially in teneral of both sexes); abdominal markings more extensive, thus the dorsal stripe on segment 2 is broader and the lateral yellow on the same segment is uninterrupted, segment 3 has a broad lateral stripe extending the whole length of segment, and tapering towards the apex, segments 4 to 7 have similar stripes but not reaching the apical border of segments and broadly interrupted at the level of the transverse suture, segment 8 has a vestige of this stripe only. The femoral spines are more numerous, more robust and more widely-spaced (6 to 7 in number); nodal index $\frac{14-19}{13-14} \frac{19-14}{13-13}$; the 1st and 8th are the primary antenodals; occiput more deeply notched than in the male. Vulvar scale of great length, as long nearly as segment 9, very narrow, tapering to a blunt point, split nearly to its base into two closely apposed scales, which closely resemble those of *Onychogomphus M-flavum*.

Distribution: Northern Bengal and Assam, from June to August. Type in the Selysian collection. Paratypes in the British Museum, Pusa and Indian Museums, and in the Laidlaw and Fraser collections. The shape of the anal appendages of the male, the shape of the vulvar scale of the female, and the presence of 2 cubital nervures in the forewing will distinguish this species from the others.

Anisogomphus orites Laid., l. c., pp. 393 and 394 (1922).


Male: Abdomen 31 mm. Hindwing 30 mm.

Head: labium with midlobe black, the lateral citron yellow; labrum black marked with two small transversely oval basal greenish-yellow spots; bases of mandibles broadly citron yellow; rest of head black save frons which is broadly citron yellow.

Prothorax black marked with citron yellow as follows:—an anterior collar, a large spot on each side, a median spot on the posterior lobe and a small geminate spot just in front of it.

Thorax black marked with bright citron yellow as follows:—a broadly interrupted mesothoracic collar, oblique antehumeral narrow stripes confluent below with the outer part of the collar, a small triangular upper humeral spot, its long axis across the dorsum of thorax and its apex confluent or almost confluent with upper part of the antehumeral stripe, a vestigial fine humeral line situated well below the humeral spot. Laterally two narrow black stripes on the lateral sutures, the hinder of which is continued back for some distance along the ventral border.

Legs black, the first pair of femora yellow internally.



Lateral views of :—1. *Davidioides martini* Fras. ♂ 2. *Anisogomphus occipitalis* Selys ♂ 3. *Anisogomphus caudalis* sp. nov. ♂ 4. *Temnogomphus bivittatus* (Selys.) ♂ 5. *Platygomphus dolabratus* Selys ♂ 6. *Platygomphus feae* Selys ♂ 7. *Anisogomphus orites* Laid. ♀

Wings hyaline, sometimes palely enfumed in adults; pterostigma dark brown between thick black nervures, over 5 to 6 cells; nodal index $\frac{14-20}{13-13} | \frac{17-13}{12-12}$; trigones, entire, but in one specimen, those of the hindwings are traversed by a nervure which runs from the basal to the outer side; *only one cubital nervure in all wings*.

Abdomen black marked with citron yellow as follows:—segment 1 with a middorsal spot and the sides broadly, 2 with a trilobed middorsal stripe extending the whole length of segment, 3 to 7 with the middorsal carina finely yellow, more conspicuous on the latter segment. Segment 2 laterally, including the oreillets, and a basolateral spot on segment 3. Segments 8 to 10 unmarked.

Anal appendages. Superiors short, greenish white, conical, pointed, each bearing a ventro-basal process somewhat like that seen in *occipitalis* but narrow at base broadening triangularly below and ending in a point behind. Inferior black, cleft into two very widely divaricate branches resembling those of *occipitalis*, but curled up a little at apices.

Genitalia unknown, not described by the author.

Female: Abdomen 34 mm. Hindwing 32 mm.

Very similar to the male. The antehumeral stripe nearly always confluent with the humeral spot, sometimes broadly so. Segment 2 more broadly yellow laterally and segments 3 to 5 with lateral stripes as in *occipitalis*, segments 6 and 7 with only a basal spot left of this stripe, and 7 with a rather large basal dorsal spot formed by an expansion of the middorsal carinal stripe. Anal appendages small, greenish yellow. Vulvar scale very different to that of *occipitalis*, very short and broad, especially at the base, its apex about half the breadth of base and very shallowly concave, not split. Hind femora with a row of 6 long, robust, widely-spaced spines, this limb extending to middle of the 2nd segment.

Distribution. Northern Bengal, Assam and Sikkim. Distinguished from *occipitalis* by having only a single cubital nervure to all wings in both sexes. The female is at once distinguished by its short broad vulvar scale contrasting strongly with the long, narrow cleft scale of *occipitalis*. For the rest, the distinctive dorsal thoracic markings will separate it from the other two species. I have had no opportunity of examining the genitalia of the male.

The type comes from Shillong, Assam, 6,000 ft. and was taken by Mr. T. Bainbrigg Fletcher and is now in the Indian Museum. Paratype females, including the one with traversed trigones in my own collection. This feature is so regular in the two opposite wings that it can hardly be regarded as an abnormality; more material will be necessary to settle this point. Frequents and breeds in mountain streams, its habits apparently similar to those of *Burmagomphus*. The female taken by Mr. Fletcher was captured flying along a hedge some distance from any stream.

Anisogomphus caudalis sp. nov.

Male: Abdomen 36 mm. Hindwing 30 mm.

Head: labium with lateral lobes greenish-white, midlobe black; labrum black with narrow linear basal streak greenish-white, nearly divided at its middle; bases of mandibles pale greenish-white; ante- and post-clypeus glossy black with a small greenish-white spot at the sides of latter against the eyes; frons black in front, broadly greenish-yellow above; vertex and occiput black the latter nearly straight.

Prothorax black marked with a large transversely oval spot of yellow on each side.

Thorax black marked with greenish-yellow as follows:—a broadly interrupted mesothoracic collar, narrow, nearly parallel antehumeral stripes forming inverted 7s. by confluence with the outer ends of the mesothoracic collar below, an upper triangular spot and a lower vestigial stripe representing a broken humeral stripe.

Laterally two narrow black stripes on the sutures, the hinder of which fuses with an elongate triangular spot beneath thorax. The ground colour at the sides greenish-yellow.

Legs black, unmarked, hind femora with a field of tiny spines on the ventral surface of the proximal three-fifths and an inner and outer row of 5 robust spines on the distal two-fifths.

Wing hyaline; pterostigma black, short; 2 to 3 cubital nervures in forewing, only 1 in the hind; 4 to 5 rows of postanal cells in hindwing, the first entire; nodal index $\frac{12-17}{16-12}$
 $\frac{13-13}{12-13}$

Abdomen black marked with citron yellow as follows:—segment 1 with a dorsal triangular spot tapering basad, and its sides very broadly, 2 with a trilobed middorsal stripe extending the whole length of segment, the oreillets and the lower part of its apical border and apical two-thirds of its ventral border broadly, 3 with its middorsal carina finely and a large baso-lateral triangular spot, 4 to 6 similar but the lateral spots very small, 7 with the basal half of dorsum narrowly, the end of this spot square, segment 8 with a small triangular spot on its dorsum at the base. Lastly 7 and 8 each with a small round apico-lateral spot, rather larger on segment 7; 9 and 10 unmarked.

Anal appendages: Superiors short, about half the length of segment 10, greenish-yellow above on dorsum, black below and at sides, tapering to a fine point and closely apposed as seen from above, broadly triangular as seen from the side, this due to the fusion of a broad triangular plate into the underside of the appendage, beginning as a short spine near the underside of apex and sloping back towards the abdomen. This plate is not free as in the process seen in the other two species. Inferior appendage deeply cleft into two widely divaricate branches, the apices of which are markedly curled upward and rather sharply pointed.

Genitalia: lobe hood-shaped, rather depressed; anterior hamules exactly similar to those of *occipitalis*; posterior also shaped similarly but much smaller and not spined on surface; lobe similar to that of *occipitalis*.

Distribution: Assam. Type from Shillong, 6,000 feet, taken by Mr. T. Bainbrigge Fletcher, June 18, 1924, now in the British Museum collection. This species closely resembles *orites* but differs in the following points:—Lateral spots on postclypeus; humeral spots further removed from the upper ends of antehumeral stripes; lateral spots on segments 4 to 6, the basal half of 7 yellow on dorsum, apico-lateral spots on segments 7 and 8; lastly the formation of the superior anal appendages. It agrees by the extra cubital nervures in the forewings and the thoracic markings are very similar. The nodal index is almost the same, especially that of the hindwings. The abdominal markings appear to be the best guide for separating the two species.

Genus—TEMNOGOMPHUS Laid.

Anisogomphus Selys *pars*. Mon. Gomph., p. 120, (1857); Laid, l.c., p. 394 (1922).




FIG. 5.—(a) Anal appendages of *Temnogomphus bivittatus* (Selys), male, seen from above. (b) The same of *Anisogomphus caudalis*, sp. nov., male, seen from above and slightly from one side. (c) The same of *Anisogomphus occipitalis* Selys, male, seen from the side. (d) The same seen from above.

A monotypic genus, containing a single rather large Gomphine dragonfly coloured black vividly marked with bright yellow. In general facies and

size, closely resembling species of *Anisogomphus*, but distinguished by the presence of an incomplete antenodal nervure in all wings and by very different genitalia. Habits and distribution similar to that genus.

Wings: reticulation close; 2 to 3 nervures between *Mi-iii* and *Miv* in the forewing, only 1 in the hind; pterostigma braced, a little swollen, short, less than one-fourth the length of distance between node and distal end of pterostigma, equal in size in both wings; all trigones, hypertrigones and subtrigones entire; trigone in forewings with costal and proximal sides approximately equal, the distal considerably longer; trigone in hindwing with costal side considerably longer than proximal and only slightly shorter than the distal; discoidal field 2-celled, nearly to level of node, divaricate thereafter; *Miii* and *Miv* markedly sinuous in forewings; *Cui* and *Cuii* parallel in both wings nearly as far as border of wings; *Cuii* not pectinate in forewing, 2 to 3 rows of cells between it and border of wing; 2 rows of postanal cells in forewing, 4 in the hind; the first postanal cell in hindwing entire, extending nearly to proximal end of subtrigone; only a single row of cells between *Mi* and *Mia*; only 1 cubital nervure in all wings; arc between the 1st and 2nd antenodal nervures; sectors of arc divergent shortly after origin; the 1st and 6th or 7th the primary antenodals; a *basal incomplete antenodal nervure present in all wings*; anal triangle 3-celled; base of hindwing oblique; tornus angulate, but slightly obtuse.

Head moderately large; face slightly oblique, frons prominent, occiput slightly concave.

Legs long, hind femora extending as far as the oreillets on 2nd segment, armed with 4 or 5 long, robust spines, the longest near the middle of the limb, a field of minute spines on the proximal half.

Abdomen tumid at base, thin and cylindrical as far as middle of segment 7, dilated thereafter as far as middle of segment 9, the segments diminishing gradually in length from 7 to 10, latter very small.

Anal appendages black or dark brown, superiors short, parallel, directed straight back. Inferior deeply cleft into widely divaricate branches.

Genitalia: lamina slightly raised, arched; anterior hamules short, triangular, tapering to a long fine point; posterior hamules enormously enlarged, pale coloured, with a broad rounded lobe projecting back and a very large robust spine projecting forward; lobe tumid, rounded, prominent.

Genotype.—*Temnogomphus bivittatus* (Selys.).

Temnogomphus bivittatus (Selys) (*Gomphus bivittatus*), Bull. Acad. Belg., xxx. (2), p. 46 (1854); id. Mon. Gomph. p. 168 (1857); Will. l.c., p. 298 (1908); Laid. l.c., pp. 394 and 395 (*Temnogomphus bivittatus*) (1922).

Male: Abdomen 36 mm. Hindwing 33 mm.

Head: labium greenish-yellow, border of middle lobe black; labrum greenish-yellow as also bases of mandibles; anteclypeus glossy black; postclypeus greenish-yellow bordered with black above; frons greenish-yellow, its lower part in front black, this black confluent with the black of upper border of postclypeus. The face is thus crossed by a succession of alternating greenish-yellow and black stripes. Vertex black; occiput markedly convex, yellow fringed with black hairs.

Prothorax black marked with a moderately large subdorsal spot on each side of the posterior lobe and a large geminate spot at its middle above.

Thorax black marked with bright greenish-yellow as follows:—An interrupted mesothoracic collar confluent with very oblique antehumeral stripes on each side which form with it inverted figures 7, long sinuous humeral stripes slightly expanded at the upper end and continuous with the yellow at base of middle pair of legs below. Laterally bright greenish-yellow marked with two fine sinuous black lines on the lateral sutures. The yellow space between the humeral black and the line on the first suture, sends a hook-like extension above into the humeral black, this curious marking being invariably present. The wavy black stripes on the yellow ground colour of the sides give a curiously tigrine effect to this insect.

Legs black, the hind femora with a linear yellow stripe running the whole length of the outer side, the middle femora with a distal stripe at the inner sides, and the anterior femora entirely yellow on the inner sides. Hind femora armed as for genus.

Wings hyaline, costa greenish yellow in subadults; pterostigma light brown between black nervures, over 3 to 4 cells; nodal index :—

$$\begin{array}{r|l} 9-16 & 14-10 \\ 8-10 & 9-8 \end{array}$$

$$\begin{array}{r|l} 11-16 & 15-11 \\ 9-11 & 11-9 \end{array}$$

Abdomen black marked with citron yellow as follows :—a middorsal stripe running from segment 1 to 7, broad and triangular on 1, broad and trilobed on segment 2, narrow and fine from thence to the end of 7, and interrupted only narrowly at the apical border of each segment. The sides of segments 1 and 2 broadly, including the oreillets. Segment 3 with a broad lateral stripe tapering apicad, and interrupted slightly at the jugal suture, 4 to 7 with a small triangular baso-lateral spot, and 7 with an additional midventrolateral spot, segment 8 with a very large baso-lateral spot which runs narrowly along the ventral border nearly to apical border and ascends some way up the dorsum, segments 9 and 10 unmarked. (In one male that I have examined, there is a much larger spot on the side of segment 8, extending as far as apex of segment, and a similar spot on the side of 9, segment 10 has a much smaller lateral spot. In all other respects it resembles the type.)

Anal appendages: Superiors black at base, dark reddish brown as far as apex, conical, subcylindrical, with a small ventral spine beneath at about their middles, directed straight back, parallel, a little separated. Inferior black or very dark brown, deeply cleft into two widely divaricate branches which project back and outward from beneath the superiors, apices pointed and a little curled up at extreme apex.

Genitalia as for genus, described above. The posterior hamules are of unique shape, black, or bright yellow in some, cylindrical at base, and broadened at end so as to be roughly T-shaped, the hinder limb long and rounded at end, the anterior end bearing a sharp robust spine tipped with black. The whole process is shaped like a horse's head, the hinder part the muzzle, the spine representing its ears.

Female: Abdomen 38 mm. Hindwing 33 mm.

Coloured and marked exactly the same as male, except that there is a continuation of the lateral stripe on segments 4, 5 and 6, and that 8 and 9 are marked as described above for an aberrant male. Segment 10 is broadly yellow laterally and the anal appendages are also reddish yellow, very short and conical.

The vulvar scale highly specialized. There is first an elliptical ridge at the ventro-apical border of segment 8, followed by a flatter ridge, from the border of which springs the vulvar scale proper, this is narrow, about half the length of segment 9, very slightly bifid at its apex which is squared and nearly as broad as its base, the surface is raised on either side into two long ridges which taper to a point which projects slightly beyond the apex of scale. These two ridges appear to be the usual bilateral scales united by a chitinous web across the mid line.

The femora are armed with longer, more widely-spaced and more numerous spines (about 8 or 9 long ones, and some shorter near the base). Nodal index $\begin{array}{r|l} 13-17 & 15-10 \\ 12-10 & 11-11 \end{array}$; pterostigma longer, over 4 to 5 cells; 5 rows of postanal cells in hindwing.

Distribution: Northern Bengal and Sikkim. Darjeeling, May 26, 1924 (at Pashok, August 31, 1920 (at Mangpu, 2,400 ft.) Assam, Kumaon, May 1911.

I have no information as to the habits of this insect. It may be at once recognized by the alternating three black stripes and three bright yellow ones on the face, by the dilated terminal segments of abdomen combined with the character of anal appendages resembling those of an—*Anisogomphus*, as seen from above, and lastly by the incomplete antenodal nervure in all wings, which is never found in *Anisogomphus*.

(To be continued.)

THE IDENTIFICATION OF INDIAN BUTTERFLIES

BY

COL. W. H. EVANS, D.S.O., F.Z.S., F.E.S.

(Continued from page 83 of this Volume.)

Part X

(With a plate)

I. Hesperiidæ.—(contd.)

I. 47. *Carterocephalus*. The Mountain Hoppers. (Plate 31.) Above with prominent white or yellow spots on both wings (or yellow with black spots). Unh prominent yellow or silver markings.

1a (4a). Unh with yellow spots; no spot in 8. Upf and Uph black with large yellow spots or yellow with black spots. Cilia ochreous.

Palæmon Group.

1b (3). Uph always with a yellow submarginal spot in 6 in addition the large yellow spots in end cell and discal in 2 and 1.

1 (2). Upf veins black on the yellow area. Upf dark brown with large yellow spots, 2 in cell, base 2, discal in 1-4, small detached spots 4 and 5, conjoined apical in 6-8, complete marginal series. Unf mostly yellow, black veins at margin. Unh ochreous, spots black edged, veins black at margin; additional large spot in 7.

palæmon, Pallas (30). The European Hopper. Europe to Amur and N. America. (= *paniscus*, F.; *brontes*, Schiff).

2 (1). Upf and margins below veins not black (slightly so in some races of *silvius*).

α. ♂ upf yellow with black spots, mid cell, end cell, near base 2 and a dash mid 1 also a submarginal row. Unh as *palæmon*, but ground colour is brown; uph all spots show but except for the dorsum the wing is overlaid yellow, obscuring all but the submarginal spots and spots in 1 and 7. ♀ dark brown with spots as *palæmon*, but larger: discal series forms a continuous band confluent with the spots in 4 and 5 and the apical spots; submarginal spots small. Uph with a spot in 8.

silvius silvius, Knoch, (30). The Northern Hopper. N. Europe to Amur.

β. ♂ upf much redder ochreous, black spots larger, inner edge spot end cell continued to costa; submarginal spots much enlarged (except 4 and 5) and reach termen. Uph spots less rounded. ♀ as *silvius*, but spots reduced, discal spots not joined to the spots in 4 and 5; submarginal spots F absent, on H absent or incomplete. Unh ♂ ♀ appearing very different, yellow spots square and conjoined, wing appearing entirely yellow with small brown spots.

silvius houangty, Oh. W. China.

γ. ♂ upf reddish ochreous, black spots very large; spot mid cell conjoined to inner edge of spot in 2, which reaches dorsum; submarginal spots large, conjoined and merged to the dark termen, except for a single spot at the apex; dorsum narrowly dark. Uph dark brown with yellow spots as in the ♀, but with traces of submarginal spots and a spot in 7. ♀ upf spots still more reduced, sub-basal yellow band mid cell to v1, discal band v1-v4 meeting outer edge of large spot before end cell, small detached streaks in 4 and 5, apical conjoined spots 6-8; uph no submarginal spots or spot in 7, only discal in 1 and 2, large discal spots in cell, 2, 4-5 and submarginal spot in 6. Unh overlaid greenish ochreous with usual black spotting.

3 (1b). *silvius shoaka*, Evans. S. E. Thibet. Bhutan at 10,000 feet. R. Uph no yellow submarginal spot in 6; 3 large spots, mid cell, 2 and 4-5, small

I. 47. *Carterocephalus*.—(contd.)

spot in 1. Upf orange costal streak from base, large spot from base 2 to v1, discal in 1-3, before end cell to costa, in 4-5 near margin, apical in 6-8. Unh shining greenish ashy, spots as above and a spot mid 7.

abax, Ob. (30). The Yellow Thibet Hopper. W. China and E. Thibet.

4a (1a). Unh spots silver. Cilia ashy grey or white.

Argyrostigma Group.

4 (5a). Unh ochreous, outwardly with ferruginous patches, no pale spot in 7; silver streak from near base cell to near termen in 5, large spot mid 1, small spot in 2, submarginal band 1-5 and large submarginal spot in 6. Above dark brown with pale yellow spots arranged as in *Taractrocera maevius*; spot at base 2 to v1, discal row 1-5, apical 6-8, spot before end cell and streak on costa above it. Uph discal spots in 2 and 4-5, submarginal row 1-4 and large submarginal spot in 6.

pulchra, Leech (25). The Beautiful Mountain Hopper. W. China and N. E. Thibet. (= *ops*, Groum).

5a (4). Unh always a silver spot mid 7, continued by a pale or silver spot to costa (in *avanti* spot is small and costa pale).

5 (6a). Upf dark brown with large yellow spots, yellow areas = dark areas; spot near base cell continued to base 2 and v1, spot before mid cell continued to costa; discal band 1-3, subapical band 4-8, streak at apex. Unh ferruginous.

a. Upf with basal costal yellow streak discal band not joined to apical or cell spots. Uph yellow spot mid cell, discal band 1-6, large submarginal spots in 3 and 6, small in 2, 5 and 7. Unh silver spots arranged as the yellow spots above, but an additional large spot mid 7, continued to costa by a yellow spot.

argyrostigma argyrostigma, Evers. (25-30). The Orange and Silver Hopper. Amur, Altai, Mongolia; Tatung Mts. and Amdo, E. Thibet.

β. Upf spots yellow white; a spot base costa, discal band of 4 conjoined spots, double apical spot. Uph large central yellow spot and two beyond. Unh as last.

argyrostigma montana, South. Litang, 14,000 feet.

7. Upf spots orange and very large; as *argyrostigma*, but no costal streak, discal band conjoined to subapical and to spot near end cell. Uph single very large orange spot and a submarginal dash in 5. Unh silver spots as the yellow ones above, but no submarginal spots and with a small spot mid 7 and cell.

argyrostigma avanti, DeN. Chumbi Valley and Kumaon, 12,000 ft. VR.

6a (5). Above pale spots small, dark areas predominate. Uph at most 2 discal spots, no submarginal spots. Spots usually white.

6 (7a). Upf white spot in 2 far from spot in cell, midway between it and termen; usually 2 small spots in 1, under spot in 2 and a small spot in 3; conjoined spots 4-5 and 6-8; apex white. Unf white bar end cell, marginal spots at apex in 6 and 7. Unh dark ferruginous; silver spot mid cell, silver central band 1-8 and submarginal band 2-7 (spots in 3 and 6 enlarged), may be discontinuous. *a*. Unh inner edge discal band regular.

**dieckmanni dieckmanni*, Graeser. (25-30). The Silver Mountain Hopper. W. China, E. Thibet. Upper Burma (Fort White). VR. (= *gemmata*, Leech and *demea*, Ob).

β. Unh inner edge discal band irregular.

dieckmanni dulcis Ob. Amdo and Kokonor.

7a (6). Upf spots in 2 and 1 immediately under and in line with spot end cell. Unh submarginal band discontinuous, at most dots in 4 and 5, usually preceded by a pale streak; inner edge of discal band irregular.

7 (8a). Above spots yellow. Uph spot end cell and in 2 separate, latter shifted in. Unh overlaid sparse yellow scales and the spot mid costa in 8 is yellow. (♀ figured by Elwes as ♂ *niveomaculata*) spots upf large.

flavomaculatus, Ob. (30-35). The Silver and Yellow Hopper. W. China, E. Thibet, Kokonor. (= *postnigra*, South from Batang).

8a (7). Above markings white and unh the pale spot mid costa is silvery.

8 (9). Unh discal band posteriorly macular, spots in 2 and 3 rounded and detached, placed behind spots in 4 and 5.

a. Upf no basal markings, spots end cell and in 2 very large, much larger than the apical spots.

I. 47. *Carterocephalus*.—(contd.)

niveomaculatus niveomaculatus, Ob. (30). The Snow Mountain Hopper. W. China and E. Thibet.

β. Upf spots base cell and costa present though faint; discal markings narrow, no wider than the apical spots.

niveomaculatus thibetanus, South. S. E. Thibet.

9 (8). Unh discal band continuous and in a straight line.

α. Upf discal spots large, prominent spot base costa and cell, dot in 3. Unh discal band broad.

christophi christophi, Groum. (30). Christoph's Mountain Hopper. Thibet Sinin Mts., Amdo and Kokonor.

β. Above markings very narrow. Unh discal band nearly linear.

christophi micio, Ob. Tsekou, E. Thibet.

(I have been unable to place the following—*demeaplutus* and *flavostigma*, Ob.)

I. 52. *Elwesia*. Leslie's Hopper.

Above dull dark brown; cilia whitish. Upf non-hyaline small whitish spots, discal in 2 and 3 (often absent in ♂) apical in 6, 7 and sometimes in 8. Uph plain. Unf brown broadly overlaid whitish scales along costa and apex.

Unh brown, entirely frosted whitish scales, giving a dull frosted appearance. In the ♀ the spots on F are larger and unh there is a white spot mid 5. Antennæ prominently chequered brown and white. Very like *Eogenes alcides*.

lesliei, nov. (30-35). Chitral at low elevations in May to July. NR. (*lesliei* is Elwes' label name in the British Museum.)

I. 53. *Apostictopterus*. The Giant Hopper. (Plate 31.)

Above and below dark chocolate brown, unmarked. Cilia dark brown. Wings very elongate.

**fuliginosus*, Leech. (65). Manipur and Naga Hills. W. China. VR. (= *curiosa*, Swin.)

I. 54. *Astictopterus*. The Forest Hoppers. (Plate 31.)

Above dark velvet brown, usually becoming paler towards the termen F; Cilia brown. Upf rarely a small discal white hyaline spot in 3; from none to 3 apical spots 6-8, which may be small and separate, prominent and conjoined or only visible unf, always more prominent in ♀♀ and DSF. Unf dark brown, dorsum pale dull brown; WSF (*jama*) apical $\frac{1}{2}$ of termen broadly paler; DSF (*henrici*) costa and apex broadly dark to light ferruginous and termen broadly overlaid whitish scales. Unh WSF dark brown usually some whitish scaling about dorsum, lower part of disc and termen also the dark bands etc., present in the DSF may be more or less apparent; DSF pale ferruginous brown, more or less whitish scaled, prominent broad dark ferruginous discal band 1-6, large spot in cell, spot on either side in 7 and base 1, sometimes the spots forming the discal band are centred white scales.

α. Small form. DSF and WSF occur. DSF darker than the Indian form and with smaller apical spots F.

jama henrici, Holland (28-40). The Forest Hopper. S. and W. China, Hong Kong, Hainan. (= *chinensis*, Leech and *nubilus*, Mab.)

β. WSF large and usually with apical spots F, DSF with prominent conjoined apical spots and very prominent bands below.

jama olivascens, M. (30-40). S. India (Swin). Kumaon to Tavoy, Andamans, Siam, Tonkin. NR. (= *kada*, Swin; *unicolor*, Mab; *tonkinianus* and *permagnus*, Fruh.)

γ. Smaller, darker form, usually with no apical spots and no DSF.

jama jama, Fd. (28-35). Mergui, Malay Peninsular and islands. (a pair DSF in British Museum from Philippines). (= *melania*, Plotz; *quadripunctata*, Swin; *fuligo*, Mab.)

I. 55. *Ochus*. The Tiger Hopper. (Plate 31.)

Above dark velvet brown. Cilia brown, chequered yellowish at apex F and H. Upf sub-apical non-hyaline yellow band 5-8, may be reduced or absent. Unf black, costa narrowly yellow and veins at apex yellow; sub-apical yellow

I. 55. *Ochus*.—(contd.)

band more extensive. Unh ochreous yellow with numerous sharply defined black streaks and spots.

* *subvittatus*, M. (22-27). Kumaon to Dawnas. Tonkin. (= *subradiatus*, M. and *intricatus*, Fruh.)

I. 56. *Baracus*. The Hedge Hoppers. (Plate 31.)

Above dark brown; upf usually with non-hyaline discal spots in 1 to 3, irregularly placed and decreasing apical spots in 6-8 (rarely outer conjoined dot in 5); uph plain and cilia dusky brown. Below costa and apex F and all H overlaid ochreous scales; rest of F black.

1 (2). Unh not brown spotted; a more or less prominent whitish or pale ochreous streak from base cell to mid termen.

α. ♂ above lower discal portion F up to discal markings and all H, except for costa and termen very pale greyish ochreous; in ♀ dark brown, with the pale markings whitish. Unh dark ochreous brown, pale central streak prominent, whitish, and with pale streaks between veins at termen.

vittatus vittatus, Fd. (30-34). The Hedge Hopper. Ceylon. NR.

β. Above uniform dark brown; upf markings faint. Unh central streak prominent, whitish, also a pale streak along v1; ground colour below brownish ochreous.

* *vittatus subditus*, M. Nilgiris and Palni Hills. NR.

γ. Above ochreous spots more prominent, especially in ♀. Unh bright ochreous, pale central streak pale ochreous and inconspicuous also broad obscure streaks between veins.

vittatus septentrionum, WM and DeN. Assam to Dawnas, NR.

2 (1). Unf bright ochreous, profusely spotted dark ochreous brown; general appearance of alternating series of large ochreous and brown spots. Upf ochreous spots prominent, traces of a spot in the cell and some yellow scales on the costa. Unf ochreous area brown streaked between the veins.

hampsoni, El. (26-32). Hampson's Hedge Hopper. Coorg and N. Kanara, NR. (May be a low elevation race of *vittatus*, *subditus* appears to be confined to the plateaux and *hampsoni* to the low country. *plumbeola*, Fd is a small pale dull blue species from the Philippines).

I. 57. *Ampittia*. The Bush Hoppers. (Plate 31.)

Above dark brown with non-hyaline markings F and H, yellow. Upf usually discal spots in 2 and 3, apical 6-8 and a spot in cell. Unh yellow with small black spots or dark brown with yellow spots.

1a (5). Unh yellow, more or less prominently black spotted. Upf never a yellow spot in the cell.

1b (4). Upf yellow spot in 1 absent or placed centrally under spot in 2.


1 (2a). Upf always a prominent spot in 1 against v1. Antennal club without apiculus. ♂ upf yellow markings very wide; entire cell and costa yellow to beyond middle, extending into bases of 1 and 2; spot in 1 usually large; discal spots in 2 and 3 conjoined to a large patch; apical spots also form large patch and may be conjoined to the costal yellow area; narrow submarginal yellow fascia below the apex; uph with a broad yellow discal area, divided by dark veins and extending from 1 to 5 or 6, may be detached submarginal spots in 5 and 6. ♀ very variable; normally dark brown with small yellow spots, on upf end cell, in 1, 2 and 3 and conjoined apical in 6-8, on uph discal in 2 and 3 and submarginal in 5, but specimens occur with markings as wide as in the ♂. Unh yellow with ill-defined black spotting. ♂ brand upf prominent.

* *dioscorides*, F. (22-28). The Bush hopper. Ceylon, India, Burma, Siam, Shanghai, Tonkin, Malay Peninsular, Sumatra, Bali. C. (= *maro*, F, *camertes*, Hew and *palemonides*, Snell).

2a (1). Upf no spot in 1 against v1 (may be very faint in *virgata*). Antennal club with well-defined apiculus. Unh black spotting sharply defined.

2 (3). Upf ♂ brand very prominent and extends below v1 nearly to the termen. ♂ upf yellow markings narrow; discal spots separate, apical conjoined; narrow basal streak along costa and often streaks between veins behind apical spots; lower edge cell spot continued along mv. towards base; yellow suffusion on basal $\frac{1}{3}$ of 1. Upf suffused discal spots in 2-3 and 4-5.

virgata, Leech, (30). The Chinese Bush Hopper, C. China. R.



I. Hesperiidæ. *Celanorhinæ* (Part).

Hesperinæ. *Panophilinæ* (Part).

I. 57. *Ampittia*.—(contd.)

3(2). Upf ♂ brand absent or small, circular, obscure.

α. Upf conjoined discal spots in 2-3, conjoined apical 6-8, elongated spots one above the other mid cell and a streak on costa over the outer end of the upper spot. Uph prominent discal spot in 4-5 at the end cell and suffused spot in 2-3 below. No ♂ brand.

trimacula trimacula, Leech. (22-25). The Trimaculate Bush Hopper, W. China.

β. Upf markings broad as in *dioscorides*; cell and costa entirely yellow, but not extending at base below cell and not joining the apical patch along the costa. Uph large suffused discal patch 2-5. Cilia H entirely yellow, ♂ with obscure brand.

trimacula maroides, DeN. Assam to Dawnas. R.

4 (1b). Upf always a small spot in 1 against v1, but this spot is placed well behind the discal spots in 2-3, which are small and separate; double spot in cell, lower part longer; separate apical spots 6-8. Uph small spots in 2 and 3 or only in 3. Unh entirely yellow, black spotting absent or very sparse. Cilia H pale yellow. No ♂ brand.

maga, Leech. (30). The Plain Bush Hopper. C. China. Malay Peninsular (Arizan, Ozaki, 2 ♀ F.M.S. Coll). (= ? *myakei* and *arisana*, Matsum and *formosana*, Fruh, Formosa).

5 (1a). Unh dark brown with prominent yellow spots in cell and bases 1 and 7, discal and submarginal series.

α. Small. Above spots obscure or absent on F and absent on H. Below dark and the spots small.

dalailama nanus, Leech. (22). The Dark Bush Hopper. China, Ichang and Ningpo.

β. Larger. Upf prominent triangular spot before end cell, discal spots in 2 and 3 close together and apical 6-8. Uph discal row 2-5.

dalailama dalailama, Mab. (30). W. China, Thibet. (= *lyde*, Leech).

I. 58. *Aeromachus*. The Scrub Hoppers. (Plate 31.)

Above dark brown. Below with a discal and submarginal row of small pale spots of which the discal row may show upf.

1a (5). Antennal club with a small apiculus.

1 (2a). Below the pale spots purple. Above unmarked. Unh dark brown overlaid with a few ochreous scales.

α. ♂ brand upf mid 1 to base 3. Below the spots prominent.

catocyanea catocyanea, Mab. (25-30). The Blue spotted Scrub Hopper. E. Thibet. VR.

β. ♂ brand a small fold mid v1. Below spots not prominent. Apex F produced in ♂.

catocyanea kali, DeN. Sikkim to N. Shan States. R.

2a (1). Below the pale spots yellowish white.

2 (3a). Unh beyond cell and unf at apex the veins white, at least near termen H; often a prominent dark spot at end cell. Upf plain or with the discal spots showing. ♂ brand upf from mid 1 to base 3.

α. Wings rounded. Unf discal band sharply angled at v5. Unh clothed ochreous olive scales, but leaving very prominent postdiscal dark spots in 1-7 between the discal and submarginal pale spots; an inner dark spot in 7 flanked by white spots and a spot end cell, the upper half of which is white. ♂ brand usually present, but vs 6 and 7 H not hair-pinned.

inachus inachus, Men. (24-32). The Veined Scrub Hopper. Japan, Amur. (= *propinquans*, Alph.).

β. Unh as *inachus*. Above and unf as *stigmata*.

inachus chinensis, El. W. China.

γ. Apex F in ♂ produced. Unf discal band evenly curved. Unh sparsely clothed olive scales, the dark markings not so prominent and sometimes not apparent, discal band not nearly so curved, the inner black spot in 7, if present, near base. ♂ with brand and vs 6 and 7 H hair-pinned.

* *inachus stigmata*, M. (22-30). Murree to Burma. NR.

3a (2). Below veins not pale. ♂ brand absent (except *piceus*) or just a fold mid 1.

I. 58. *Aeromachus*.—(contd.)

3 (4). Below discal pale spots not conspicuously larger than the sub-marginal. Unh clothed olive ochreous scales.

γ. ♂ with a brand.

discreta piceus, Leech (22-28). The Gray Scrub Hopper. E. Thibet.

β. ♂ no brand.

discreta discreta, Plotz. Assam to Burma, Tonkin, Malay Peninsular, Sumatra. NR.

γ. Unf markings very obscure. ? no brand.

discreta javanicus, El. Java.

4 (3). Below discal band of pale spots conspicuously wider than the sub-marginal series.

γ. Below dark; unh overlaid ochreous scales, spots prominent.

jhora dubius, El. (22-28). Dingy Scrub Hopper. Palnis, Travancore, C.

β. Below brighter; unh overlaid ochreous scales and spots prominent.

jhora jhora, DeN. Sikkim. ? to N. Burma. R. (I have not seen *jhora* from outside Sikkim nor *discreta* from W. of Assam; it is possible that they are conspecific since the genitalia are alike).

5 (1a). Antennal club without apiculus. Wings rounded, small. Below markings obscure; unh overlaid olive grey scales.

γ. ♂ no brand.

pygmaeus pygmaeus, Fab. (20-22). The Pigmy Scrub Hopper. S. India. Assam to Burma, Malay Peninsular, Sumatra. NR. (= *indistincta* and *obsoleta*, M).

β. ♂ with brand.

pygmaeus musca, Mab. Luzon.

I. 59. *Pedestes*. The Bush Bobs. (Plate 31.)

Above with hyaline spots on F only.

1a (3). Upf with spot across cell, discal spot in 3 and a larger one in 2 running below the above two; 3 apical spots in 6-8.

1 (2). Above blue-black, central hyaline spots large and white, spot in 3 to base 3. Below costa and apex F and all H overlaid olive ochreous scales. Unh small white discal spots in 2 and 3.

* *masuriensis*, M. (30-32). The Mussoorie Bush Bob. Mussoorie to Assam. NR.

2 (1). Above dark brown, hyaline spots pale yellow, smaller, spot in 3 not to base 3. Below costa and apex F and all H overlaid ochreous scales. Unh unmarked, but faint traces of discal spots are sometimes visible. In a not infrequent variety the discal spots in 2 and 3 upf are widely separated from the cell spot.

* *pandita*, DeN. (25-30). The brown Bush Bob. Sikkim to N. Burma. NR. (= ? *tali*, Swin, Haut Yunnan and *parnaca*, Fruh, Tonkin).

3 (1a). Upf with 2 spots end cell, 3 discal, 3 apical. Unh dull red brown, several basal and a curved discal row of indistinct black spots.

bononoides, Druce. (34). Sarawak. (Not examined by me; placed by Druce in *Pedestes*, but position doubtful).

I. 60. *Arnetta*. The Bobs. (Plate 31.)

1a (3a). Above dark brown with small hyaline white spots upf, upper edge cell (may be absent in *vindhiana*), discal in 2 and 3 close together, apical 6-8 minute. Unh with more or less complete set of small white spots, discal row 1-7, base 7, mid cell, base 1 (double).

1 (2). ♂ unf no tuft. Unh ochreous brown to greenish ochreous, spots obscure.

* *vindhiana*, M. (25-32). The Vindhyan Bob. S to C. India. NR. (= *nil-giriana*, and *modesta*, M).

2 (1). ♂ unf with a tuft of black hairs mid dorsum and v1 distorted in middle. WSF unh dark brown, clothed dark ochreous scales with spots very prominent. DSF pale to dark brick red more or less clothed whitish scales, spots small and sometimes incomplete.

* *atkinsoni*, M. (25-32). Atkinson's Bob. Sikkim to Tavoy. NR. (= *subtestaceus* and *khasianus*, M.)

3a (1a). Unmarked above and unh.

1. 60. *Arnetta*.—(contd.)

3 (4). Unf unmarked except for dorsum being paler. Above and unh dark rich brown. ♂ with black tuft unf and vl. distorted as in *atkinsoni*.

kala nov. (30). Sumatra. (Described from 1 ♂ of which the antennæ and palpi are missing, but the genitalia are very similar to *atkinsoni*; in general appearance the insect resembles *Itys iadera* and *Iambrix unicolor*).

4 (3). Unf with a conspicuous triangular apical orange yellow patch from 6-8. Unf costa and apex and unh very sparsely overlaid with conspicuous ochreous scales.

verones, Hew. (31). Sumatra. Kuala Tahan, Pahang (Singapore Museum). (Usually considered as a variety of *Koruthaialos rubecula*, but is structurally quite different; the palpi antennæ and venation show a close alliance with *atkinsoni* and the genitalia though peculiar, are allied).

1. 61 *Iambrix*. The Bobs. (Plate 31.)

Normally without hyaline spots except sometimes in the ♀ of *salsala* on upf and in *tytleri*.

1a (4a). Unh with small silver spots. ♂ no brand.

1 (2a). Upf with prominent pale yellow hyaline spots, large spot towards end cell, tapering towards scv., larger spot in 2 with its inner edge in line with the inner edge of the cell spot, may be a dot in 3, apical spots 6-8, non-hyaline spot in 1 against vl. Below costa and apex F and all H overlaid bright ferruginous ochreous scales; unf the spots white and the dorsum pale; unh prominent small silver spots in 2, 3, 5, 7 and cell, more or less black edged and some black streaks towards the margin.

tytleri, Evans. (28). 'Tytler's Bob. Naga Hills, Siam, Borneo. VR. (= *woolletti*, Riley).

2a (11). Upf no prominent hyaline spots (except ♀ *salsala* which never has a spot in the cell).

2 (3). Unh black edged silver spots in 1, 5 and 7, that in 5 most prominent; rarely a faint spot in cell, but never so prominent as the spot in 5; spots in 1 and 7 often faint or absent. Above dark brown with some dark ochreous scaling on F, which in ♂ forms a more or less apparent series of discal spots; in ♀ these spots are more or less replaced by a non-hyaline spot in 1 and a series of hyaline spots in 1-8, bent in at v6. Unf costa and apex and all unh densely clothed ferruginous ochreous scales; unf in ♂ a more or less prominent silver spot end cell against upper edge and two small discal spots in 4 and 5.

α. Smaller. Upf ♂ only ochreous spots in 1 and 2.

salsala luteipalpus, Plotz. (26-30). The Chestnut Bob. Ceylon. South India. C.

β. Larger. Upf very variable, but usually discal spots in 1 and 2 are obscure, those in 3-6 being more conspicuous.

**salsala salsala*, M. Sikkim to Burma, China, Siam, Malay Peninsular and Islands. C. (= *vasuba*, Fruh, Tonkin).

3 (2). Unh normally with small silver spots in 1 (double), 2, 5 and mid cell, the spot in cell as prominent as the spot in 5. Above much darker velvet brown, very little ochreous scaling about apex F and no formed spots. Unf costa and apex and all unh overlaid dark ochreous: unf silver cell spot present, spots in 4 and 5 usually absent.

stellifer, But. (24-28). The Malay Chestnut Bob. Karens to South Burma. Annam, Malay Peninsular and Islands. Philippines. NR. (= *niasicus*, Fruh).

4a (1a). Unh unmarked. ♂ upf a glandular streak along 2nd quarter of vl from base, the vein being slightly deflected.

4 (5a). Above unmarked dark velvet brown, paler in ♀. Below as above; unf lower discal area to dorsum paler.

unicolor, DeN. (31). Perak, Pahang, Malacca, Sarawak. R.

5a (4). Upf with red discal band. Below costa and apex F and all H overlaid dark ochreous scales. ♂ unf a thin tuft of hairs mid dorsum overlying a black and white brand in 1; uph scales above cell to costa modified.

4 (5). Upf red band dull, narrow, appears composed of densely packed dark ochreous scales, does not extend into 1 and is not wider than the dark termen.

I. 61. *Iambrix*.—(contd.)

sindu, Fd. (25-28). The Small Red Bob. South Mergui VR. Malay Peninsular and Islands. NR. (= *obliquans*, Mab. and *yamanta*, Fruh. Felder's types of *sindu* at Tring are ♂ *Koruthaialos rubecula* and ♀ *kophene*; he describes *sindu* as near *Astictopterus jama*, but smaller; Felder's type of *jama* at Tring is *Koruthaialos xanites*, but his description refers to an insect without any markings; the conclusion is that the Tring types are unreliable).

5 (4). Upf red band much broader, clearer and brighter, much broader than the dark terminal area and extends into 1; in ♀ it extends into cell, the end of which is marked by a dark line.

latifascia, El. (27-30). Borneo. Sumatra (Fruh). Pahang (F.M.S. Museum).

I. 62. *Suastus*. The Palm Bobs. (Plate 31.)

Above dark brown, with or without hyaline spots F. Unh usually with dark spots.

1 (2a). Unh grey (brown, overlaid greyish ochreous scales) with a variable number of sharply defined black spots; spot against upper outer edge of cell always present and as large as any other spot; discal series 1, 2, 3, 5 (usually absent) and 6. Upf hyaline white spots usually prominent, but may be small or absent; spot across cell beyond middle (lower part often absent), spot in 3 and large spot in 2 with its inner edge under the centre of the cell spot and its outer edge under the inner edge of the spot in 3; apical spots 6-8; usually non-hyaline lower spot in 1 and sometimes an upper dot. Unf apex and costa grey and there may be some black spotting at apex. Cilia grey. Size very variable.

α. Much darker above and below dark grey.

gremius subgrisea, M. (32-42). The Indian Palm Bob. Ceylon. C.

β. Generally paler.

* *gremius gremius*, F. India, Burma, South China (= *divodasa*, M; *robsonii*, DeN; *nigroguttata*, Matsum, Formosa).

2a. (1). Unh with obscure ill defined, suffused black spots in a decreasing row from 1-3, 6 and upper edge of cell; spot in 1 always present and larger than the rest.

2b. (4). Uph no white about the tornus.

2 (3). Above black, unmarked. Cila F brown; H white, chequered brown. Unf a white patch mid 1; unh white to v6 and scv., some obscure brown marginal spots.

minuta, M. (28-32). The Ceylon Palm Bob. Ceylon. NR. (= *sinhalus*, Plotz).

3 (2). Above dark brown with hyaline spots F.

α. Upf small hyaline spots in 2 and 3, rarely a dot in 6. Cilia brown. below with more or less purple sheen; apex and costa F and all H overlaid sparse grey white scales, general appearance dark grey brown. Unf obscure discal white area in 1.

rama bipunctus, Swin. (26-34). The Small Palm Bob. Nilgiris. R.

β. Upf always hyaline spots in 2 and 5 and usually a double (or single lower) spot in cell and a spot in 6; the discal spots much closer together than in *gremius*. inner edges of spot in 2 and cell spot in line; traces of spot in 1 against v1, which unf shows as a broad white patch extending across 1. Cilia narrowly white becoming brown at apex F. Below white scaling much denser, general appearance pale grey.

rama rama, Mab. Bengal, Sikkim to Tavoy R. (= *sala* Auct nec. Hew; true *sala* is a *Plastingia*).

α. Above spots prominent. Cilia H grey, brown at apex H and on F. Below apex and costa F and all H overlaid pale ochreous scales; unf white area mid 1 more extensive, reaching dorsum.

* *rama aditus*, M. Andaman and Coco Islands. R.

4 (2a). Uph with white tornal area. Above black, no spots upf (may be faint in race *scopas*). Cilia white to apex H. Below as *minuta*.

α. Uph white area narrow and irregular, most extensive along vs 1a and 1, not reaching dorsum.

everyx everyx, Mab. (26-34). The Malay Palm Bob. Tavoy, VR. Malay Peninsular, Sumatra, Pulo Laut, Borneo. (= *tripura*, DeN).

I. 62. Suastus.—(contd.)

β. Uph white tornal area regular and sharp defined, extending $\frac{1}{4}$ along termen and dorsum.

everyx scopas, Stg. Java, Bali and Philippines. (= *albescens*, Mab.; *migreus*, Semp.; *chilon*, Doh, Sumba).

(*plana*, Swin, New Guinea, is described as belonging to this genus).

I. 63. Inessa. DeNicèville's Bob.

Above dark brown with a conspicuous purple gloss. Upf double spot in cell, lower portion prolonged nearly to base; small non-hyaline spot in 1 against v1, spots in 2 and 3 and dots nearer margin in 4 and 5, apical in 6-8. Uph small spot mid cell; continuous discal band 1-5, separated by veins and spots composing it elongate and irregular. Below as above but no purple sheen.

ilion, DeN. (32). Lombok. VR.

I. 64. Scobura. The Forest Bobs. (Plate 31.)

Above dark brown with hyaline spots F and H; upf always non-hyaline spot in 1 against v1, large spot in 2, double spot in cell, of which the upper part is smaller and may be absent, apical spots 6-8 (may be incomplete).

1a (3a). Upf no spot in 3 (rarely a dot in *cephala*) and the cell spot is immediately over the spot in 2. Uph large double hyaline spot in 4-5 and smaller one in 2 (may be absent). Cilia grey on H.

1 (2). Upf usually a small spot in 4. Unf costa and apex and all H greenish yellow. Unh there may be 3 small white spots in 1, 3 and 6 against the larger spots in 2, 4-5 and the spots are usually chestnut edged; in ♀ the chestnut colour is very pronounced and usually occupies the whole area from the spots to the termen and on the apex upf.

* *cephala*, Hew. (27-36). The Forest Bob. Sikkim to Burma, Malay Peninsular, Sumatra and Borneo. NR. (= *isota*, Swins.).

2 (1). Upf never a spot in 4. Uph spot in 4-5 always large and spot in 2 obscure or absent. Unf apex and costa and all H overlaid greyish green. Unh spots black edged and often a discal row of black spots and 2 spots in the cell, also black.

phiditia, Hew. (32) The Malay Forest Bob. Manipur to Burma. Malay Peninsular and Sumatra. Borneo. R. (= *martini*, El).

3a (1a). Upf prominent spot in 3.

3 (4). Upf no spots in 4 and 5, spot in 2 shifted out. Uph small equal hyaline spots in 2, 3 and sometimes 5, spot in 3 much nearer margin. Below basal half costa F and basal half H bright yellow; apex F and outer half H bright chestnut; unh small dark ringed white spots in 1, 2, 3, 5, 6 and dark spot base 7 (may be absent).

cephaloides, DeN. (36-40). The Large Forest Bob. Sikkim to Burma. Tonkin and Hainan. R.

4 (3). Upf figure of 8 spot in the cell, large spot base 2, small spots 1, 3, 4, 5 and 6-8. Uph a spot in 2 and a double spot in 4-5. Unh and apex unf veins pale.

coniata, Hering. China. (? = ab' of *cephala*).

I. 65. Suada. The Grass Bobs. (Plate 31.)

Above brown with hyaline white spots F, typically prominent lower spot in cell, large spots in 2 immediately below it, dot base 3, apical spots 6-8 and an elongate non-hyaline spot along mid v1. Unh unmarked.

α. F spots prominent; uph rather paler mid disc. Unf apex broadly and all H clothed dense olive ochreous scales.

* *swerga swerga*, DeN. (32-36). Sikkim to Burma, Malay Peninsular, Sumatra. NR. (= *moelleri*, M).

β. F spots prominent. Larger. Uph much paler mid disc in ♂ and in ♀ whitish, costa broadly and termen narrowly brown. Unf apex brown above v6 and then decreasingly white to tornus; unh all white to narrow brown costa.

swerga triplex, Pl. Java. (Plotz gives no locality and I prefer to use his name for the Javan race rather than invent a new name.)

I. 65. *Suada*—(contd.)

γ. F spots on disc small, separate and may be absent. Uph tornus broadly pure white extending half-way along dorsum and termen. Unf and unh as *triplex*.

swerga cataleucos, Stg. Borneo, Pulo Laut, Philippines. (= *albinus* Semp).

I. 66. *Koruthaialos*. The Veivet Bobs. (Plate 31.)

Above and below dark brown, no spots but usually with a prominent red discal band upf and unf; unh no scaling.

1a (3a). Palpi third joint long and thin.

1 (2). Upf discal red band sharply defined, 3 mm. wide, clear red and extending from upper edge of cell (or just beyond) to v2 (or just beyond). Upf band rather more extensive at either end, sides straight and parallel or convergent posteriorly. A fairly constant form.

rubecula, Plotz. (32-37). The Narrow-banded Velvet Bob. Assam to Burma. Malay Peninsular and Islands, Philippines, Luzon, Natuna and Celebes. C. (= *hector*, Wat and *namala*, Fruh. I have verified true *rubecula* from Assam to Malay Peninsular and Sumatra, but the other localities require check).

2 (1). Upf discal band absent, just indicated or sullied.

α. Upf unmarked; unf usually some faint red suffusion at upper end of cell and in ♀ a dusky suffused band.

butleri butleri, WM. (32-37). The Dark Velvet Bob. Sikkim to N. Burma. R. (I have verified the type specimen in the Indian Museum.)

β. Upf discal band sullied, dull, 2 mm. wide, from upper end cell to v3 or 2, dcu. prominently black. Unf variable, may be a trace of red only at end cell or a dusky red band from costa to below v1.

butleri avidha, Fruh. S. Burma. S. Annam. NR.

γ. Upf unmarked or with traces of a red spot at end cell. Unf a large dusky red area about upper edge of cell and a detached smaller area about base 3 (may be absent).

butleri niasicus, Fruh. Sumatra, Nias.

δ. Larger, otherwise as last.

butleri haraka, Fruh. Java.

3a (1a). Palpi third joint short, stouter, just protruding.

3b (5). Upf always a red band.

3 (4). Unf discal red band divided by dark veins, discontinuous at v1 and the red area is extensive in 1. Upf band dusky red, divided by dark veins and the dcu. black, extends from costa to tornus, 2 mm. wide in ♂, 5 mm. in ♀. Large.

focula, Plotz. (46-52). The Large Velvet Bob. Sumatra, Java. R. (= *kophene*, DeN).

4 (3). Unf discal red band not divided by black veins, extends from costa to dorsum and is more or less equally wide throughout, broad and clear red, inner edge below cell always concave.

α. Upf band orange red, from costa to tornus, usually wide (5 mm., but may be as narrow as 3 mm. and stop at v3; on unf in narrow banded specimens the inner edge of the band is irregular, but always concave). ♀ unh ochreous brown.

* *xanites gopaka*, Fruh. (34-40). The Bright Red Velvet Bob. Assam to Burma. Tonkin. NR.

β. Larger. Upf band deeper red and always broad.

xanites xanites, But. (37-41). S. Burma. Malay Peninsular. Borneo and Philippines. NR. (= *letitia*, Plotz; *palawites*, Stg.; *luzonensis*, Fruh).

γ. As last but larger still.

xanites kerala, DeN. (49). Sumatra.

δ. Upf discal band very dark red and reduced, may consist of only of a spot end cell. Unh band as broad as usual.

xanites javanites, Stg. (36-39). Java, Bali, Batu.

5 (3b). Above and below dark brown with a purple plum suffusion, no traces of a red band; Unf dorsum broadly pale pinkish brown. F v11 not so close to v12.

nigerrima, Swin. (36-42). The Plain Velvet Bob. Assam to S. Shan States. R. (I have verified the type.)

I. 67. *Sancus*. The Coon. (Plate 31.)

Above dark shining brown, unmarked. Below dark brown, dorsum F pale; on apex F a narrow oblique purple white patch, small whitish post-discal spots in 4 and 5 and subapical in 6-9. Unh dorsum broadly pale brown, a purple grey area about end cell, separated from a similar broad marginal area by discal spots in a straight line from 2-5, spots in 6, 7 and in middle of upper edge of cell.

a. The markings below prominent.

* *pulligo subfasciatus*, M. (36-46). South India, Cachar to Burma, Siam, Annam. Malay Peninsular. C.

β. Larger. Below the purple grey areas absent, the spots dull yellow.

pulligo pulligo, Mab. Sumatra, Java, Borneo and Bali. (= *forensis*, Plotz. *kethra*, Plotz = *ulunda*, Stg. is the Philippine race and *fuscata* Snellen = *cellundo*, Stg., the race from the Celebes).

I. 68. *Watsoniella*. Watson's Demon.

Above dark brown unmarked, becoming outwardly paler on F, in ♀ broadly and very markedly so. Below similar, but paler and unf dorsum to v2 very pale brown. Very like *Kerana diocles*.

swinhoei, Elwes. (48-50). Sikkim to Burma. R.

I. 69. *Ge*. The Circular Tufted Demon.

♂ above uniform dark brown, cilia grey; ♀ with pale yellow discal spots in 2 and 3 and apical in 6-8, conspicuously paler beyond the spots; uph in ♂ centrally paler. Below dark brown, dorsum F pale brown; in ♀ the outer half of the wing is conspicuously very pale brownish yellow.

geta, DeN. (35-37). S. Mergui. Malay Peninsular Sumatra, Batu, Java and Borneo. VR.

I. 71. *Kerana*. The Demons. (Plate 31.)

Above dark brown with or without a red band F.

1 (2a). Above dark brown unmarked, cilia brown; ♂ no band. Below dark brown, termen F and H broadly paler; on H the basal area is also paler, leaving a dark chocolate costa and discal band to v1; costa and apex F and all H clothed inconspicuous ochreous scales. F v5 bent at origin.

* *nigrita*, God. (48-60). The Chocolate Demon. Bengal, Kumaon to Burma, Siam, Malay Peninsula and Islands, Philippines. C. & Individuals from the Malay Islands have a more or less prominent whitish curved sub-apical fascia unf.

(= *diocles*, M; *maura*, Snell; *fumatus*, Mab.; *evaira*, *sumata*, *prabha*, Fruh).

2a (1). Upf broad red discal band (may be absent in *niasana*).

2b (5). F v5 bent down at origin. Body dark brown.

2 (3a). Below no spots. ♂ unf towards base a large black velvet patch from v1 to mid cell; uph a similar patch filling the cell.

a. Upf a very broad orange red discal band from mid costa to tornus at v1; narrower at costa and indented at the origin of v5.

armata armata, Druce. (49-60). The Red Demon. S. Mergui, VR. Malay Peninsula, Sumatra and Borneo.

β. Upf, band absent, obscure or broken into spots.

armata niasana, nov. Nias.

3a (2). Below with curious glistening pale purple spots only visible in a side light; unf in cell just before the red band, sub-apical in 5, 6 and 7, unh discal in 2, 3, 5 and 6. Upf and unf a broad orange red band from mid costa to v1 at tornus.

3 (4). Upf the red band narrows towards costa.

gemmifer, But. (35-38). The Gem Demon. Tavoy to S. Burma, Malay Peninsula, Sumatra, Borneo, Natuna, Philippines. R. (= *dombya*, Fruh).

4 (3). Upf discal band broadens towards costa.

vaijrada, Fruh. (36). Fruhstorfer's Demon. Borneo. VR.

5 (2b). F v5 only slightly bent down at origin. ♂ upf a very broad orange yellow band from mid costa below the scv. to mid dorsum; at v2 it is curved in and nearly reaches the base. Thorax and base H clothed orange yellow hairs.

I. 71. Kerana.—(contd.)

Abdomen with narrow white rings. Unf discal band to costa. In ♀ upf the red band is not continued towards the base posteriorly and the yellow hairs on the thorax and at base H are absent, very like *xanites*.

fulgur, Den. (37). The Yellow Demon. Malay Peninsular, Sumatra and Borneo. VR. (I am not satisfied with the position of this species but have only been able to examine damaged specimens.)

I. 73. Udaspes. The Grass Demon. (Plate 31.)

1 (2). Above dark brown with large hyaline white spots; upf a spot across cell, conjoined to a bent spot in 1-2, small detached spot in 3, conjoined spots in 4 and 5, conjoined sub-apical spots in 6-8; uph a large central hyaline area in 1-6. Cilia chequered brown and white. Antennæ white banded below club. Below dark brown, apex F, dorsum and termen H overlaid pale whitish scales, which on H leave a large dark spot mid 1 to base 2. Considerable seasonal variation in the colouring below.

**folus*, Cr. (40-48). Ceylon, India, Kangra to Burma, Siam, China, Malay Peninsular and Islands C. (= *cicero*, Fab.)

2 (1). Uph a prominent white spot in 4-5 and a small spot base 3. Unh striated.

stellatta, ob. (40-45). The Thibetan Grass Demon. Maenia, E. Thibet. VR.

I. 74. Notocrypta. The Banded Demons. (Plate 31.)

Above black; upf a broad curved hyaline white discal band consisting of large conjoined spots in 1, 2 and cell. Below dark brown usually with some whitish scaling on the paler areas at the apical part of the termen F and the dorsal portion H also on H from mid end cell to mid v1; often some ochreous scaling between band and apex on F and on basal half of H. Antennæ white banded below beginning of a club or only white streaked below.

1a (4a). F no traces of sub-apical spots in 6-8; never a well-defined spot in base 3 (only in *renardi*). Unf a yellow costal patch at end of the band.

1 (2a). F discal band very wide, width = basal black area and placed nearer base than apex, more compact and straighter than usual; unspotted except very rarely a faint dot mid 4 unf. F much squarer than usual, termen at right angles to dorsum. Below with more or less white scaling.

quadrata, Elwes. (40-45). The Broad banded Demon. Malay Peninsular, Borneo. R.

2a (1). F discal band narrower and placed rather nearer apex than base. F dorsum at an angle of 120 to 150 to termen.

2 (3). F unmarked except for the discal band. Apex F produced. Below white scaling only traceable at apex F; but margins broadly paler brown, also centre of disc H, thus resembling *Kerana nigrila*.

α. F discal band broad and compact.

volux volux, Mab. (39-42). The Dark banded Demon. Karens to S. Burma, Siam, Malay Peninsular and Islands, Philippines. NR. (This is a common species in S. Burma and passes as *albifascia*, which is a *Hyarotis*; figured in Lep. Ind. as *albifascia*; the correct name and synonymy in this and other species in this genus is doubtful.)

β. F discal band narrow, tending to be macular and the spot in 1 may be absent.

volux niasana, nov. Nias.

3 (2). F almost invariably a prominent spot mid 4, if not visible upf may show unf; very rarely a dot mid 5 and there may be a dot mid 3 against v3. Wings rounded and band more curved than usual.

α. Below white scaling usually prominent. Unh no silver spots.

paralysos alysia, nov. (33-40). The Common Banded Demon. Ceylon, S. India, Mussoorie to Karens. C.

β. Below white scaling faint and usually obsolete.

paralysos devadatta, Fruh. Dawnas to S. Burma, Malay Peninsular, Sumatra, Nias. C.

γ. Darker, larger, wings rather more pointed. Below very little or no white scaling.

I. 74. *Notocrypta*.—(contd).

paralysos sidha, Fruh. Java. (= *asanga*, Fruh. and *sukavata*, Fruh. Sumbawa).

♂. Unh a prominent silver spot mid cell and small spots in 2 and 5. Larger, band broader; below white scaling faint.

paralysos paralysos, WM. Andamans. NR.

(The race from the Philippines and Celebes is *clavata*, Stg.=*chunda* and *yaya*, Fruh.).

4a (1a). F always with 2 or 3 sub-apical spots, which are at least traceable unf; usually small spots mid 3, 4 and 5.

4 (5). Unf the discal band continued to costa by a more or less well developed yellowish white patch. The white spotting and scaling below is very variable; local races are very difficult to define.

α. Small. Spotting and scaling variable.

* *feisthamelii alysos*, M. (38-50). The Spotted Demon. Murree to Burma. W. China, Siam, Malay Peninsular, Sumatra and Borneo. C. (= *rectifasciata*, Leech).

β. Larger. White spotting and scaling below prominent.

feisthamelii avattana, Fruh. Java, Lombok and Sumbawa. (= *samyutta* and *dharana*, Fruh. The Philippine race is *alinkara*, Fruh and the Celebes race *celebensis*, Stg. Race *feisthamelii*, Bdv., flies from the Moluccas to Queensland and = *varians* and *chimaera*, Plotz.; *padhana*, *samana* and *satra*, Fruh.; ? *klossii*, Roth).

5 (4). Unf discal band not continued above scv. Above spotting and below white scaling prominent. Apical spots conjoined.

curvifascia, Fd. (38-50). The Restricted Demon. Ceylon, S. India, Sikkim to Burma, China, Andamans, Siam, Malay Peninsular, Sumatra. ? Java, Borneo and Philippines. C. (= *restricta*, M.).

(An allied species, *renardi*, Oz., with many named forms and possibly comprising more than one species flies from the Moluccas to Australia; it differs principally in having a well defined spot at base 3 and the spot in 1 tends to be detached).

I. 75. *Gangara*. The Giant Redeye. (Plate 31.)

Above brown with large discal hyaline yellow spots on F, across cell, in 2 and 3. Below some blue white scaling at apex F and bands composed of similar scales on H.

1 (2). F with apical spots in 6-8, spot in 6 out of line, nearer margin; spot in 3 against mid v4. Cilia H grey at apex.

α. Normally with a small non-hyaline spot in 1 against v1 under the spot in 2.

thyrsis clothilda, H5. (70-76). The Giant Redeye. Ceylon. NR.

β. No spot in 1.

* *thyrsis thyrsis*, F. India. Sikkim to Burma, Siam, Tonkin, Malay Peninsular and Islands. NR. (= *pandia*, Fruh).

γ. F spot in 2 not to base of v3; in other races spot in 2 is larger and reaches to middle of cell spot.

thyrsis yasodara, Fruh. Andamans.

δ. Upf only 2 apical spots in ♂.

thyrsis philippensis. Fruh. Philippines.

2 (1). F no apical spots; discal spots differently placed, spot in 3 reaches to behind origin of v4 and nearly to origin of v3; spot in 2 against mid v3 and not near cell spot. Unh a white spot near base 7. ♂ upf the glandular streak absent; unf yellow tuft present.

saguinocculus, Martin. (65-70). Martin's Redeye. Malay Peninsular, Sumatra, Borneo. R.

I. 76. *Erionota*. The Palm Redeye. (Plate 31.)

Above brown with large hyaline yellow discal spots on F, in cell, 2 and 3; spot in 3 against mid v4.

1 (2). Upf no hyaline spots (very rarely faint); spot in 2 reaches to middle (or further) of spot in cell. Unh pale to dark brown with an irregular, often obscure, paler discal area.

I. 76. *Erionota*.—(contd.)

a. Upf hyaline spots and apex white.

thrax grandis, Leech. (70-76). The Palm Redeye. W. China. VR.

β. Upf hyaline spots yellow.

* *thrax thrax*, L. S. India. Dun to Burma. Siam, Malay Peninsular and Islands, Philippines. NR.

γ. Smaller. Apex F above more or less white; antennæ below club much whiter and cilia above whitish. Unf a white apical patch.

thrax acroleuca, WM. and DeN. (52-55). Andamans and Nicobars. NR. Occurs as a scarce variety of *thrax* in Assam to Burma, Malay Peninsular and Borneo. (= *hiraca*, M and *lara*, Swin).

(*alexandra*, Semp, is a race or perhaps species from Luzon. *sakita*, Ribbe is the Celebes race and the Moluccan is *hasdrubal*, Fruh).

2 (1). Upf with a hyaline spot in 6-8, spot in 6 pushed out (not so much as in *thyrsis*); spot in 2 does not reach to the middle of the cell spot; apex may be more or less white tipped. Unf sub-apical pale area. Unh dark brown with a more or less purple gloss, margin broadly paler, conjoined irregular violet white spots about end cell and from 1-7, also a spot at extreme base cell and sometimes a spot near termen in 1.

sybirita, Hew. (60-70). The Sybarite Redeye. S. Mergui, VR. Malay Peninsular, Borneo and New Guinea.

I. 77. *Paduka*. The Banded Redeye, (Plate 31.)

♂ above dark brown, unmarked. ♀ with large hyaline yellow discal spots in cell, 2 and 3, spot in 2 with its inner edge against middle of cell spot and its outer edge against inner edge of spot in 3, which is against mid v4; small non-hyaline spot in 1 against spot in 2. Unf pale sub-apical area, overlaid sparse white scales and a more or less prominent spot on 7; dorsum whitish. Unh a broad prominent transverse band of violet white scales from v1 below cell to apex and expanding towards apex. Cilia above pale yellow.

a. Unh transverse band well defined throughout, violet white.

* *lebadea subfasciata*, M. (62-68). Ceylon. R.

β. Unh as last; no trace of the spot in 7 unf.

lebadea glandulosa, Dist. Sikkim to Burma, Tonkin, Malay Peninsular, Sumatra and Java. VR.

γ. Unh transverse band silvery white, dense from v1 to end cell and thence sparse, extending to middle of disc. Unf scaling to apex. ♀ spots whiter, smaller and more separated, spot in 1 larger.

lebadea andamanica, WM. Andamans. R.

δ. Upf ♂ brand more distinct. ♀ no spot in 1 upf.

lebadea lebadea, Hew. Borneo, Pulo Laut. R.

(*toradja*, Fruh. is apparently a distinct species from the Celebes, with 3 large white spots upf and some grey scaling unh, not forming bands).

I. 78. *Pudicitia*. The Spotted Redeye. (Plate 34.)

Above brown with large yellow hyaline spots F and H. Upf 2 spots in cell (lower nearer base and behind origin of v3) large spot in 2 immediately below, spot in 3, $\frac{1}{2}$ along v4; apical spots 6-8 in line; semi-hyaline spot above middle of v1 replaced by a yellow patch unf. Uph 3 hyaline yellow discal spots in 2, 3 and 4-5. Below as above.

* *pholus*, Den. (58-64). The Spotted Redeye. Bhutan to Naga Hills. VR.

I. 79. *Matapoides*. The Tufted Redeye.

Above dark brown; upf a green streak in certain lights below v2; otherwise unmarked; cilia orange at tornus H. Below rich dark brown, unmarked; unf ♂ a brand on the nacreous area mid 1. H lobed.

smaragdinus, Druce. (60). Borneo. VR.

I. 80. *Matapa*. The Branded Redeye. (Plate 31.)

Unmarked dark brown.

1 (2a). H. cilia grey or very pale yellow. Unh warm ferruginous brown. ♂ brand black, obscure.

I. 80. *Matapa*.—(contd.)

aria, M. (40-45). The Common Redeye. Ceylon, S. India to Dun to Burma, Andamans, China, Hainan, Siam, Tonkin, Malay Peninsular and Islands to Philippines. C. (= *pulla*, Plotz and *neglecta*, Mab.).

2a (1). H cilia and tip of abdomen bright orange.

2b (4b). ♂ brand black, obscure.

2 (3). Above purple glossed, especially at apex F. ♀ body and base wings clothed metallic bluish hairs. ♂ below dark brown, apex and dorsum F paler; ♀ uniform pale brown.

purpurascens, El. (48-54). The Purple Redeye. Sikkim to Karens. Tonkin, Sumatra, Borneo, R.

3 (2). Above no purple gloss, apex F grey. Below ochreous brown; apex and mid dorsum F and base H very pale brown.

druna, M. (48-54). The Dark-brand Redeye. Sikkim to Burma. Andamans, Malay Peninsular and Islands. NR.

4a (2b). ♂ brand grey and prominent.

4 (5). Below dull dark brown, veins conspicuously black.

sasivarna, M. (40-47). The Black-veined Redeye. Sikkim to Burma, Malay Peninsular, Sumatra, Pulo Laut. NR.

5 (4). Below ♂ dark ferruginous brown; ♀ pale yellow brown. Upf and unf apex rather paler.

* *shalgrama*, M. (46-52). The Grey-brand Redeye. Sikkim to Burma, Andamans, Tonkin, Java, Bali, Pulo Laut. NR.

(*celsina*, Fd. with a tear-shaped brand occurs in the Philippines, Celebes and Moluccas. It = *hyrmira*, Hew : *assur*, Mab and *ractaya*, Fruh).

I. 81. *Oerane*. The Demon Flitter. (Plate 31.)

Above black; upf compact broad white central band from upper edge cell to mid v2 and a small conjoined spot base 3; no apicals pots or spot in 1 or on H. Below dark brown, dorsum F paler; apex F and all H very sparsely overlaid greenish ochreous scales; unf discal band is continued in ♂ to v12 by greenish ochreous scales and in ♀ by white scales. ♀ paler and with wider band.

* *neaera*, DeN. (28-34). Dawnas to S. Burma. Malay Peninsular and Islands. R. (= *pusilla*, Fruh. *drymo*, Mab. is a variety with the spot in 3 missing. *microthyrsus*, Mab. is the race from the Philippines. In Rhop Java the ♀ is figured as unmarked).

I. 82. *Hyarotis*. The Flitters. (Plate 31.)

1 (2a). Cilia broadly chequered white and brown; antennæ white banded before apiculus. Above dark brown; upf with white hyaline spots, across cell, in 1 (non-hyaline) against v1, discal in 2 and 3, apical 6-8. Below dark brown, outer half dark ochreous, bearing centrally a broad diffused dark brown band; unf cell spot continued to costa and spot in 1 diffused: unh an irregular broken central white band from v1 across end cell to v8.

α. Upf spots small, those in cell and in 2 do not overlap.

adrastus adrastus, Cr. (38-48). The Tree Flitter. Ceylon, S. India. NR. (= *phoenicis*, Hew).

β. Upf spots larger, those in 2 and 3 overlapping.

* *adrastus praba*, M. Kangra to Burma, Andamans, Siam, Hong Kong. NR.

(*meluchus*, Fruh is the race from the Malay Peninsular. Pulo Condor, Sumatra, Borneo and Java. *mindaneusis* and *palawensis*, Fr., the Philippine races).

2a (1). Cilia uniform dark brown; antennæ plain brown. (White under club in ♀♀ and ♂ *pria*).

2 (3a). Unh base dark golden yellow. Upf and unf large hyaline white spot across cell, joined to a still larger spot in 2. Otherwise unmarked.

basiflava, DeN. (38-48). The Golden Flitter. S. India. R.

3a (2). Unh unmarked.

3 (4). ♂ upf with a hyaline white discal band consisting of a large spot across cell conjoined to an equally large spot in 2, which extends (non-hyaline) more or less into the upper part of 1.

I. 82. Hyarotis.—(contd.)

pria, Druce. (35-45). The Banded Flitter. Dawnas to S. Burma. Siam, Tonkin, Malay Peninsular and Islands. VR.

(= *albifascia*, M; *asawa*. Fruh; *inornata*, El; *singularis*, Mab. This seems to be an extremely rare species and I have not seen a ♀, unless a ♀ specimen from Selangor is the true ♀; it has no spot in 1, a completely detached dot in 3 and apical dots in 6 and 8).

4 (3). ♂ unmarked dark brown above and below; ♀ with a hyaline white discal band consisting of large conjoined spot in 2 and cell, usually a conjoined or semi-conjoined spot base 3 and sometimes a conjoined non-hyaline spot in the upper part of 1.

* *monteithii*, WM. (35-40). The Dubious Flitter. Cachar, Dawnas, Malay Peninsular and Islands. Philippines. ? Moluccas (Holland). VR. (= *noctis*, Stg.; *comoploea*, Swin.; *dissimilis*, Snell.; *perfusca*, Mab.).

I. 83. Itys. The Brush Flitters. (Plate 31.)

1 (2). Upf with small pale yellow hyaline spots, narrow spot across cell constricted in middle or divided in two, quadrate to nearly vertical linear spot in 2, small spot in 3 and apical in 6-7 or 8. Unf dorsum pale, diffused white discal spot in 1; apex F and all H dark purple washed in fresh specimens; unh bearing a very obscure dark central and postdiscal broad band, small yellow spots in 2, 3, 4 and end cell, one or more of which may be absent.

* *microstictum*, WM. (34-39). The Brush Flitter. Cachar, Dawnas, Malay Peninsular (Pulo Ubin). Luzon, Mindoro. VR. (= *binghami*, Swin and *biseriata*, Elwes).

2 (1). Above unmarked dark brown with a faint purple tinge, more prominent below. Uph costa and unh dorsum paler. ♀ paler and larger.

iadera, DeN. (32-36). The Dark Brush Flitter. Penang, Sumatra, Java and Borneo. VR. (= *ciliata*, Elwes).

I. 84. Zographetus. The Flitters. (Plate 31.)

Above dark brown with hyaline white spots on F; 2 cell spots (upper small and often absent), quadrate spot in 2 immediately under the cell spot, small spot in 3, apical 6 and sometimes 7, small non-hyaline spot in 1 in ♀ and ♂ of *satwa* against vl. Cilia greyish.

1 (2). Below basal half costa F and basal $\frac{3}{8}$ H bright yellow, outer $\frac{2}{8}$ H and apex F purple brown, paling outwardly. Unh small dark spot mid cell and mid 7.

* *satwa*, DeN. (30-36). The purple and Gold Flitter. Kumaon to Burma, Siam, Malay Peninsular, Sumatra and Java. NR. (*durga*, Plotz from Mindanao and Sumbawa is a race or allied species; it has a purple gloss, antennæ white below club and no glandular streaks in ♂).

2 (1). Unf costa and apex and all unh ferruginous to ochreous (paler in ♀), bearing unh a more or less prominent discal row of purple brown spots in 1-7, a spot mid 7 and mid cell. Antennæ white banded before apiculus in ♀.

α. Extremely variable in size, colour of underside, and spots unh, which are usually large and diffused, but may be separate and sharply defined, often upper discal spots absent.

* *ogygia ogygia*, Hew. (30-32). The Purple spotted Flitter. S. India. Sikkim to Burma, Malay Peninsular and Islands. R (= *flavipennis*, DeN.; *flavalum*, DeN.; *ogygioides*, Elwes).

β. Unh dark yellow of a peculiar shade; lower discal spots coalesced to a very broad dark band, connected by dark veins to a broad dark marginal area.

ogygia andamana, nov. Andamans. R.

I. 85. Isma. The Tufted Flitters. (Plate 31.)

Above dark brown with pale yellow detached spots upf; 2 spots in cell lower elongate, spot in 2 outwardly concave, small spot in 3, may be dot in 4 and apical in 6 and sometimes 7; ♀ may have a non-hyaline spot in 1 against vl. Unf dorsum white. Unh a more or less obscure discal row of dark spots in 1-6 and a spot end cell. Cilia dark brown at tornus H in ♂ and at apex F, elsewhere the long hairs grey or pale yellow.

85. *Isma*.—(contd.)

1 (2). Comparatively small. Below dull dark brown, apex and costa F and all H more or less sparsely overlaid ochreous green scales.

protoclea, HS. (30-34). The Plain Tufted Lancer. Karens to S. Burma, Malay Peninsular, Java, Borneo, Banguay, Borneo, Pulo Laut. R. (= *iapis*, DeN. and *obscura*, Dist.).

2 (1). Comparatively large. Below purple brown.

α. Below purple brown, the outer $\frac{1}{2}$ F and lower $\frac{3}{4}$ H greyish white, all H with a conspicuous purple gloss.

* *vulso bicolor*, nov. (38-40). The Purple Tufted Lancer. Karens and Dawnas, Malay Peninsular. R.

β. Below purple brown, overlaid ochreous scales, no purple wash, the dark spots unh in 2 and 3 may be white centred. Upf hyaline spot in 2 more quadrate.

vulso vulso, Mab. Sumatra, Java and Bali. R.

γ. Below with a feeble purple gloss on basal half H, otherwise rather as *protoclea*. The clasp of the genitalia is serrate on the upper edge, not plain as in rest.

vulso purpurascens, El. Borneo, Pulo Laut. (= *binotatus*, El.).

(*sewa*, Plotz from the Celebes probably belongs to this genus).

I. 86. *Sepa*. The Lancers.

1a (4a. 6a). ♂ with only a discal brand from mid v1 upf to base 3.

1 (2a). ♂ cilia at tornus H elongated as in *Isma*; brand upf angled at v2. ♂ upf no spot in 1 or in cell; a very narrow oblique linear pale yellow spot in 2, small spot in 3, may be a dot in 4, small spot in 6 and sometimes in 7. Below paler, but not overlaid ochreous scales, outer half of 1 pale; unh a small pale spot in 2 and indications of a spot in 4 end cell. ♀ paler, spots on F wider, may be 2 spots in cell; unf space 1 all dark. Cilia in ♂ dark ochreous, brown at tornus, dark brown in ♀.

cronus, DeN. (42-44). The Branded and Tufted Lancer. Malay Peninsular (Perak and Gunong Ijan), Sumatra and Borneo. VR.

2a (1). ♂ cilia at tornus not prolonged. ♂ brand upf not angled. ♂ upf nearly always a spot in 1, rhomboidal spot in 2, decreasing spots in 3 and 4, small spots in 6 and 7 and may be a dot in 8, 2 elongate spots in cell, lower linear; all spots white or very pale yellow. Uph cuneate hyaline spots in 2, 3 and 4. Below overlaid ochreous scales. Cilia ashy ochreous.

2 (3). ♂ upf spot in 1 rounded, not reaching v2. Below darker, spots equal, no spot in cell or in 1 and 5. ♀ spots in cell reduced or absent; uph spots small, round, spot in 4 may be absent.

feralia, Hew. (40-44). The Cicatrose Lancer. Peninsular Siam, Malay Peninsular, Sumatra, Java, Borneo, Pulo Laut. VR. (= *cicatrosa* and *umbrosa*, El.; *indistincta*, Druce. The synonymy is doubtful; all the species in the genus are very rare and it is very difficult to link the sexes).

3 (2). ♂ upf spot in 1 narrow, vertical, reaching v1 and may be divided by a black line. Below more ochreous; unh spots larger may be a spot in cell and often small spots adjoining the spots in 2 and 4 may show above. ♀ spots reduced.

fenestrata, El. (36-39). The Fenestrate Lancer. S. Mergui, Malay Peninsular, Sumatra, Borneo, Pulo Laut. (= *concinna*, El.). VR.

4a (1a. 6a). ♂ with curved linear stigma behind the spot in 2 upf and an oval patch above basal $\frac{1}{4}$ of v1.

4 (5). Upf spots white, large, arranged as in *feralia*, cell spots often conjoined. Uph ♂ very small discal spots in a curve in 2, 3 and usually 4; ♀ unmarked.

dawna, nov. (38). The Dawna Lancer. Dawnas to S. Burma. VR.

5 (4). Upf spots yellow, arranged much as *feralia*, but very variable F and H. ♂ stigma may be absent.

miosticta, DeN. (42-44). The Cinnamon Lancer. Malay Peninsular, Java, Borneo. VR. (= *guttulifera* and *cinnamonea*, El.).

6a (1a. 4a). ♂ upf no seam, but with an oval patch above basal $\frac{1}{4}$ of v1; spot in 2 is differently placed due to the absence of the seam, its inner edge in line with the inner edges of the cell spots; spots yellow. Unh obscure series of black discal spots in 1-5. ♀ spots reduced and cell spots usually absent.

I. 86. *Sepa*.—(contd.)

6 (7). Uph spots in 2-4 cuneate, prominent, those in 2 and 3 inwardly pointed and reaching bases 2 and 3; may be spot in cell and unh in 5.

inarime, DeN. (36-38). The Cuneate Lancer. Mergui, Malay Peninsula, Sumatra, Java, Borneo, Pulo Laut, Philippines. R. (= *zetus*, Mab.; *matanga*, Druce; *bipunctata*, El).

7 (6). Uph spots in 2, 3 and 4 small rounded, latter often absent. Smaller. *bononia*, Hew. (26-32). The Baby Lancer. Karens to S. Burma, Malay Peninsula, Sumatra, Java, Borneo, Pulo Laut. R. (= *idyalis*, DeN).

I. 87. *Plastingia*. The Lancers. (Plate 32.)

Above with prominent yellow or white spots (hyaline) upf; more or less developed pale basal and costal streak F and discal area H, ochreous or greenish.

1a (6a). F v2 origin much nearer base than the origin of v11; v5 bent down at origin. Upf always a hyaline spot in 4. Above markings and cilia bright yellow.

Callineura Group.

1b (5). Unh with purple white spots.

1c (4). Antennæ at least with the entire club yellow.

1 (2a). Uph tornus broadly yellow.

vermiculata, Hew. (40). The Vermiculate Lancer. Sumatra. VR.

2a (1). Uph tornus dark brown.

2 (3). Unf no purple white streak in 5.

α. Unh veins red. Upf cell spots one above the other, lower hardly produced towards base. Antennæ shaft yellow in ♂, club only in ♀.

callineura burmana, nov. (39-42). The Red-Vein Lancer. Cachar to S. Burma. R.

β. Unh veins yellow. Upf lower cell spot prolonged towards base and conjoined to spot in 2. Antennæ as last.

callineura perakana, nov. Malay Peninsular (Perak, Negri Sembilan, Pahang). R.

γ. Unh and apex unf with red veins. Upf cell spot as in last. Antennæ in ♂ and ♀ with the club only yellow.

callineura callineura, Fd. Sumatra, Java, Borneo.

3 (2). Unf with a purple white streak in 5. Unh veins yellow. Antennæ shaft yellow.

α. Upf cell spots conjoined or nearly so.

latoia margherita, Doh. (36-40). The Yellow-vein Lancer. Upper Assam (Doh). Tenasserim (1♂ B.M.) VR.

β. Upf cell spots separate, lower nearer base.

* *latoia latoia*, Hew. Dawnas to S. Burma. Siam, Malay Peninsula, Sumatra, Java, Borneo and Labuan, Philippines. R.

4 (1c). Antennæ with only the apiculus yellow. Above like *latoia*, but smaller. Unh mostly yellow. Very variable.

α. Upf basal yellow area restricted.

helena fruhstorferi, Mab. (32). The Small Yellow-vein Lancer. S. Mergui, Malay Peninsula, Sumatra, Java, Batoe Is. R. (= *niasana*, Fruh.).

β. Above basal yellow area upf much more extensive.

helena helena, But. Borneo, Labuan, Pulo Laut. R. (= *natuna*, Fruh. *flavia*, Stg. is the race from the Philippines and ? Celebes).

5 (1b). Unh no purple white spots. Upf the yellow spot in 2 reaches base 2 all the discal spots form a broad continuous band. Unh orange with small black spots.

aurantiaca, El (35-39). The Orange Lancer. Sumatra, Borneo, Pulo Laut. R. (Species allied to the above are *telesinus*, Mab. = *laenas*, Mab. Philippines with unh yellow green bearing feeble pale spots. *liburnia*, Hew, Philippines and Luzon, unh dark brown with prominent yellow spots).

6a (1a). F v2 from mid base and origin v11.

6b (11a). F v5 straight or nearly so. Upf no spot in 4. Unh with markings.

Sala Group.

6c (9a). Unh with a semi-circular row of small black discal spots, a round spot end cell and spots at base 2 and 7. Unf some dark spotting at the apex.

I. 87. *Plastingia*.—(contd.)

6d (8). Uph no prominent yellow area. Upf a rather obscure dark ochreous streak from base to middle over v1. Antennæ white ringed mid club.

6 (7). Antennæ shaft only chequered at extreme base. Upf lower cell spot larger than the upper spot; apical spots in 6 or 6 and 7; hyaline spots pale yellow. Uph in ♂ a recumbent tuft of shining dark brown hairs from base overlying cell. Below apex and costa F and all H dark grey brown with a strong purple wash. Tegumen with ears at the side.

* *sala*, Hew. (32-36). The Purple Lancer. Cachar to S. Burma. Malay Peninsular, Pulo Laut. R. (= *fuscicornis*, El.).

7 (6). Antennæ shaft chequered throughout. Upf lower cell spot no larger than the upper one and may be absent; no apical spots; spots white. Uph ♂ no tuft. Tegumen with spines at the side.

α. Below apex and costa F and all H pale yellow brown with a faint purple wash.

submaculata kanara, nov. (32-36). The Maculate Lancer. N. Kanara. R.

β. Unh, etc. shining greenish grey.

submaculata maculicornis, El. Ataran Valley, Burma, Siam, Pulo Laut. VR.

γ. Below yellow green.

submaculata submaculata, Stg. Palawan.

8 (6d). Uph with a prominent yellow discal patch and upf with prominent yellow basal streaks. Unh yellow. Uph tornal area brown. Upf cell spots equal.

α. Uph yellow discal area divided by dark veins. Unh spots complete and diffused.

* *noemi noemi*, DeN. (35-40). The Spotted Yellow Lancer. Sikkim to Assam. VR.

β. Uph yellow discal area not divided by black veins. Unh markings mostly consist of narrow rings and may be incomplete.

noemi tavoyana, nov. Karens (Doh). Tavoy, Mergui. VR.

(Species allied to *noemi* are *similis*, El., Borneo, Pulo Laut, much smaller. *flavescens*, Fd. = *atala* and *samanga*, Fruh, Celebes, with the tornal area uph broadly yellow, *viburnia*, Semp, Philippines, with unh prominently black veined).

9a (6c). Unh chequered all over with large black and yellow or white pale spots.

9 (10). Below pale spots white in ♂ and ♀. Above yellow markings obscure; upf lower cell spot small or absent.

* *naga*, DeN. (33-38). The Silver Spotted Lancer. Assam to Burma, Malay Peninsular, Sumatra, Java, Borneo, ? Philippines. (= *valenia*, Fruh.). R.

10 (9). Below spots pale yellow. Above yellow markings prominent; upf lower cell spot well developed.

tessellata, Hew. (33-40). The Straw spotted Lancer. Malay Peninsular, Sumatra, Java, Borneo, Philippines, Celebes. R. (= *eulepis*, Fd; *palawata*, Stg.; *pellonia* and *mangolina*, Fruh.).

11a (6b). F v5 bent down at origin. Unh unmarked.

Corissa Group.

11 (12). Upf no spot in 4. Uph discal yellow area small or absent cilia yellow. Below apex F and all H yellow, veins paler.

α. Above unmarked.

corissa tytleri, Evans. (30). The Plain Yellow Lancer. Nagas. VR.

β. Upf prominent discal yellow spots in 2, 3, apical in 6-7 and 2 cell spots, non-hyaline spot in 1.

* *corissa corissa*, Hew. (30-45). Dawnas to S. Burma, Malay Peninsular, Sumatra, Java, Borneo, Pulo Laut. R. (= *indrasana*, El.; *drancus*, Plotz.; *latonia*, Stg.; *patmapana*, Fruh.).

12 (11). Upf with a spot in 4, over the spot in 3, no spot in 1 and basal streak obscure; spots yellow. Uph plain. Below apex F and all H overlaid dark ochreous scales, unmarked.

pugnans, DeN. (34-36). The Pugnacious Lancer. Dawnas to S. Burma, Malay Peninsular and Islands. R.

(*extrusus*, Fd, Aru is a peculiar species, mimicking *Allora doleschalli*).

I. 88. *Lotongus*. The Palmers. (Plate 32.)

Above with prominent hyaline spots on F.

1a (3a). Unh no yellow band. Upf spots white.

Calathus Group.

1 (2). Upf hyaline spots large and conjoined, spot in 3 to base 3, non-hyaline yellow spot in 1 against v1, apical spots 6-8 present or absent. Uph costa and apex to v7 whitish yellow. Unf discal spots continued to costa and broadly to dorsum. Unh termen white from apex to mid 3.

* *calathus*, Hew. (44-46). The White Tipped Palmer. Tavoy to S. Burma, Malay Peninsular, Sumatra, Java, Borneo. Palawan. R. (= *zalates*, Mab. and *aliena*, Stg.).

2 (1). Upf hyaline white spots smaller and completely separated. ♂ double (♀ single) spot in cell; ♂ quadrate (♀ oblique) spot mid 2, small spot in 3, no apical spots and only obscure spot in 1 in ♀; spots may be even more reduced. Unf discal band continued to costa and a large white spot in 1; unh apex may be yellowish white.

schædia, Hew. (42-45). The Plain Palmer. Malay Peninsular, Sumatra, Nias and Borneo. R. (= *parthenope*, Weymer; *traviata*, Plotz; *surus*, Mab. and *maculatus*, Dist).

(*taprobanus*, Plotz. = *mythecoides*, DeN. is an allied species from the Celebes with a prominent white band unh).

3a (1a). Unh with a conspicuous yellow band.

Avesta Group.

3 (4a). Uph tornus and dorsum yellow joining a broad central yellow area; unh central yellow band very broad, from v1 to costa, parallel to termen. Upf double white spot in cell, spot mid 2 and a spot in 3 against its outer edge, apical spots in 6-8. Unf apex paler, cell spot joined to costa by a yellow area. Antennæ club white below. Resembles *Hasora schonherri*.

onara, But. (42-44). The Yellow Palmer. Sumatra, Borneo, Palawan. VR. (= *excellens*, Stg. and *adorabilis*, Fruh.).

4a (3). Uph dorsum dark brown. Below chocolate brown, dorsum F much paler; unf cell spot continued to costa by a yellow spot; unh a narrow yellow band, from basal $\frac{1}{2}$ dorsum to mid costa, interrupted at v1a. Upf no apical spots.

4 (5). Upf 2 spots in cell, conjoined to a single spot in ♀; large spot in 2, prolonged in ♀ under cell spot, prominent spot in 3, non-hyaline spot in 1. Uph tornal cilia yellow and central area from v1-7. Unh band rather broader and expanding to costa, slightly curved. H excavated at v1 and lobed. Below purple washed and veins pale. Hyaline spots white.

sarala, DeN. (45-50). The Yellowband Palmer. Assam to Bhamo, Tonkin, W. China. VR.

5 (4). Upf only rounded pale yellow spots in 2 and 3. Uph unmarked and cilia uniform dark brown. Unf an upper spot in cell. Unh band narrow and straight, nearer base. Below chocolate brown, no purple wash and veins not pale. H termen even.

avesta, Hew. (44-48). The Malay Yellowband Palmer. Dawnas to S. Burma, Borneo, Pulo Laut. VR. (= *tamiata*, Stg. *quinquepuncta*, Joicey and Talbot is a well marked race from Hainan).

(To be continued)

ON SOME INTERESTING FEATURES OF THE FAUNA OF THE WESTERN GHATS

BY

SUNDER LAL HORA, D.SC.,

Officiating Superintendent, Zoological Survey of India.

[Read at the thirteenth annual meeting of the Indian Science Congress at Bombay and communicated with the permission of the Director, Zoological Survey of India.]

The nearness of the Western Ghats to the place of our meeting this year and the charming peculiarities of its fauna have induced me to write this short note. It is written chiefly with the object of exciting some interest in the study of Natural History among the advanced students of zoology of the Bombay University. There are two striking peculiarities of the fauna of the Western Ghats, firstly the presence of a marine element in the fauna, and secondly the wonderful adaptations exhibited by several animals to the extremely wet and dry weather conditions that prevail at different times of the year in the Ghats. I shall illustrate my point by referring to four definite instances, which of late have attracted great attention and about which more information is badly needed. I want to refer to *Succinea arboricola* Rao, *Cremnoconchus syhadrensis* Blanford, *Lithotis rupicola* Blanford and *Limnocnida indica* Annandale. Besides these I shall have occasion to refer to several other interesting forms found in this area.

A very interesting mollusc, *Succinea arboricola* Rao,¹ was recently discovered by me aestivating on the bark of mango trees at Lonavla. A short note² on its habits was read at the last meeting of the Congress and has since then been published in the *Records of the Indian Museum*.³ Several of the specimens collected during August and September, 1924, are still with me aestivating on the bark, and I have not been able to find out at what time of the year they become active. In the midst of heavy rains I found *S. arboricola* in deep slumber, and in the last week of December last year I found them in a similar comatose condition. The only changes that have been observed are that the number of individuals found in December is much less and that the mollusc, instead of being found on the bark, was found aestivating on scars left by the falling of leaves from small twigs. This observation leads to the conclusion that the animals had been leading an active life for some time during the months intervening between September and December. It would be extremely interesting to make fortnightly observations on these molluscs with a view to

¹ Rao, *Rec. Ind. Mus.*, xxvii. pp. 394-400 (1925).

² Hora, *Proc. Twelfth Ind. Sci. Cong.*, p. 148 (1925).

³ Hora, *Rec. Ind. Mus.*, xxvii, pp. 401-403 (1925).

studying their behaviour under natural conditions from season to season, and to work out their entire life-history.

Cremnoconchus syhadrensis Blanford¹ is another peculiar mollusc of an essentially marine family—*Littorinidae*—found on rocks below the falls at Khandhalla. The species seems to possess a very localised distribution, for last year when I was specially looking for them, I did not find a single specimen anywhere between Bombay and Mahabaleshwar except below the two big falls near Khandhalla. During the rains the falls are not quite accessible, but at other times of the year it takes from twenty to thirty minutes to reach them from the station with a suitable guide. Towards the end of December, after a very careful and long search, only two individuals of *Cremnoconchus syhadrensis* were found by me on the rocks kept moist by spray of water from the fall. They were, however, quite plentiful on big rocks projecting out of the water at the edge of the pool in the neighbourhood of the fall. They were found aestivating in small pits well protected from the mid-day sun. Drs. Annandale² and Prashad³ have already referred to the peculiar mode of life of this species.

In the Nilgiris there is another mollusc of the family *Neritidae* which possesses more or less similar habits to those of *Cremnoconchus*. *Neritina perott etiana* Recluz⁴ is found on rocks kept moist by spray from a fall or on the edges of streamlets where they are occasionally washed by the current of water. The Nilgiris, unlike the portion of the Western Ghats in the immediate neighbourhood of Bombay, are never extremely dry, because they get heavy rainfalls during both the monsoons. In accordance with the climatic conditions prevailing in the Nilgiris *Neritina perott etiana* is probably not called upon to hibernate in the dry season as does *Cremnoconchus syhadrensis*, and it was found that the animals of the former species died a few hours after they were removed from their natural habitat, while those of the latter are capable of surviving long periods of dessication.

Lithotis rupicola Blanford⁵ is another highly peculiar mollusc of a very limited range. It was first described by Blanford from the edge of the waterfall at Khandhalla in 1863, but since then it has remained in obscurity. Very little was hitherto known regarding its habits and anatomy. From the material collected by me in 1924, Dr. H. S. Rao⁶ has been able to confirm its distinct position in the family *Succineidae*. *Lithotis rupicola* was found in great abundance along with *Cremnoconchus* during rains, but we know very little about its mode of life after the rainy season. The late Dr. Annandale,⁷ who visited the locality in March and April, records the entire absence of this mollusc from the cliffs below the Khandhalla falls. In December last year I

¹ Blanford, *Ann. Mag. Nat. Hist.* (3), xii, p. 184, Pl. IV. (1863).

² Annandale, *Rec. Ind. Mus.*, xvi, pp. 119, 148 (1919).

³ Prashad, *Proc. Twelfth Ind. Sci. Cong.*, pp. 138, 139 (1925).

⁴ Recluz, *Rev. Zool. Cuv.*, p. 333 (1841).

⁵ Blanford, *Ann. Mag. Nat. Hist.* (3) XII, p. 186, Pl. IV, figs. 8-10 (1863).

⁶ Rao, *Rec. Ind. Mus.*, xxvii, pp. 387-394 (1925).

⁷ Annandale, *Rec. Ind. Mus.*, xvi, p. 119 (1919).

made a thorough search for it but with no success. Not even a single dry shell of the species was seen on the spot. The Succineid molluscs are known to hibernate on rocks, trees, etc., during the unfavourable season, but the exact conditions under which *Lithotis rupicola* passes the dry season still remains to be investigated. Another species of *Lithotis*, *L. tumida* Blanford,¹ known from Poona and the adjoining country, has not been found again since its discovery in 1870, and we know very little either about the structure of its animal or its habits.

I should also like to mention here a small species of *Paludomus*² and a highly modified, air-breathing Ampullarid—*Turbinicola saxea* Reeve³—which are abundant during the rainy season all over the Ghats from Khandhalla to Mahabaleshwar. Very few specimens were seen of the latter species in December, and to work out the entire life history of these two fairly common molluscs will be a nice piece of work.

In the Ghats in 1911, Prof. Agharkar discovered a widely distributed freshwater medusa, and realising the significance of such a discovery, Annandale sent a short note to *Nature* and to the *Proceedings of the Asiatic Society of Bengal*.⁴ Next year when he examined better preserved material, he⁵ christened the medusa *Limnocrnida indica*. In the same year a note on its habits and distribution was published by Gravely and Agharkar.⁶ In 1913 Agharkar⁷ published a further note on the species. The medusae have so far been collected from the Krishna river at Dhoni, Yenna river at Medha, and the Koyna river at Tambi. I was informed by an old fisherman at Wai that these 'flowers' were also to be found in a big tank adjoining the bridge near Wai, and there is no apparent reason why these medusae should not be found all over the valleys of these rivers in suitable places. Several attempts have been made to find the asexual phase of this medusa or the hydroid stage, but so far without any success. The medusae appear in March and April and disappear suddenly with the first showers of the monsoon rains. The best way to find its hydroid stage is to pay occasional visits to these places from February to May and to look carefully for the growth on the rocks, or to try to breed the medusae under more or less natural conditions.

Besides the peculiarly modified individuals mentioned above, there is a great variety of interesting animals found in the Ghats and the report published by the late Dr. N. Annandale on the fauna of certain small streams in the Bombay Presidency, shows the wealth of material that is to be found in this area and the fascinating characters of its fauna.

¹ Blanford, *Journ. As. Soc. Bengal*, xxxix, p. 23, Pl. 3, Fig. 24 (1870).

² Annandale, *Rec. Ind. Mus.*, xvi, p. 147 (1919).

³ Prashad, *Mem. Ind. Mus.* viii, pp. 87, 88 (1925).

⁴ Annandale, *Proc. As. Soc. Bengal*, for May, 1911, also Annandale, *Nature*, xxxiii, p. 144 (1911).

⁵ Annandale, *Rec. Ind. Mus.*, vii, pp. 253-256 (1912).

⁶ Gravely and Agharkar, *Rec. Ind. Mus.*, vii, pp. 399-403 (1912).

⁷ Agharkar, *Rec. Ind. Mus.*, ix, pp. 247-249 (1913).

RECENT ADDITIONS TO THE INDO-CEYLONESE COCCID FAUNA,
WITH NOTES ON KNOWN AND NEW FORMS

BY

T. V. RAMAKRISHNA AIYAR

In 1921 the writer prepared a check list of the Indo-Ceylonese Coccidæ then known and the same was published in the Report of the Proceedings of the Fourth Entomological Meeting at Pusa the same year. Since then, a few additional papers¹ have appeared containing new records and further data on known forms. Of these the most important is Green's fifth volume on Ceylon Coccidæ. This extremely valuable addition to the literature on Asiatic Coccidæ from one of the foremost authorities on the group not only contains new records from Ceylon but includes a wealth of information on the whole family, especially with regard to the taxonomy and nomenclature of the various sub groups and species. With the idea of making the writer's list of 1921 more complete and up-to-date for purposes of reference, this paper is prepared as a supplement to the same. All new forms recorded from the region since the publication of the list of 1921 and a few that were omitted from that list by oversight are included here with such notes on known forms which appeared to the writer to be worth recording. In addition, the more important changes in taxonomy and nomenclature, especially those pointed out by Mr. Green, are also indicated. It is believed that the original list with this supplement may be of some use to Indian workers on this group until such time when a complete annotated catalogue of the family with all up-to-date data can be prepared.

NOTE.—Those forms with an asterisk in the following list are species which have been referred to in the previous list; others are new records noted since then. All new forms with manuscript names were collected by the writer, and were very kindly identified by Mr. Green. This supplementary list was also perused by Mr. Green and the necessary amendments made, and the writer hereby expresses his thanks for the continued help he has been getting in this manner.

Sub-Family.—MONOPHLEBINÆ.

Monophlebus, Burm.

M. phyllanthi, Green. p. 169. A. M. N. H. xii. 1923. This new species was confused with *M. contrahens* Wlk. and was identified as such by Green in his recently published fifth part of Ceylon Coccidæ. He has since found out the mistake and described it as *phyllanthi* in the above reference. Material of *phyllanthi* has been collected by the writer from Malabar and Godavari on *Cleistanthus* and *Croton*² respectively.

Icerya, Sign.

I. purchasi, Maskell. This pest which appears to be a recent introduction into Ceylon was omitted in the 1921 list; not noted in India as yet; p. 221, N. Z. Trans., xi, 1878 and p. 436, Green, C. of C. V. 1922. Found in Ceylon on *Acacia*, sp. *Casuarina*, *Citrus*.

-
- ¹ 1. Green, 'Coccidæ of Ceylon' Pt. v, 1922. Dulau & Co., London.
2. Green 'Supplementary notes on the Coccidæ of Ceylon', *Bombay Natural History Journal*, vol. xxviii. 1922, pp. 1007-1037.
3. Ramakrishna Aiyar, 'A Further Contribution to our Knowledge of South Indian Coccidæ,' Report of the 5th Ent. Meeting, Pusa, 1923, pp. 339-344.
4. Misra, 'A List of Coccidæ in the Pusa Collection' 5th Ent. Meeting, Pusa pp. 345-350, 1923.
5. Green, On *Monophlebus Contrahens* W, and a new sp p, 168. A. M. N. H., xii. 1923.
² Referred as *M. tamarindus* in previous list.

I. aegyptiaca, Douglas.* (1890) = *I. Tangalla*, Green (1896) *vide* Green, p. 439 of C. of C. V., 1922.

I. seychellarum, West.* (1855) = *I. Crocea*, Green (1896) *vide* Green, p. 440. C. of C. V., 1922.

I. seychellarum, var *nardi* Green. p. 441, C. of C. V., 1922 on *Andropogon nardus*. Diyatalawa (Ceylon).

Leachia festiva, Kief. Kieffer records this species on mango-Bengal on p. 163. Marcellia. xii. 1908.

I. jacobsoni, Green. p. 316. Tijdschrift V., Ent. lv. 1912 on *Shorea Bangalore*. (M. Hassan Coll.)

Walkeriana, Sign.

W. ovilla, Green. p. 448, C. of C. V., 1922, on *Eugenia subavenis* and *Michelia nilagirica*, Hekgala, Ceylon.

Aspidoproctus, Newstead.

Walkeriana cinerea,* Green (a) and *W. euphorbiæ*, Green.* (a) are brought under the genus *Aspidoproctus*. Green, pp. 450 and 453, C. of C. V., 1922.

Labioproctus, Green.

Walkeriana polei, Green. 1896, is brought by Green under this newly erected genus *Labioproctus*. p. 453, C. of C. V., 1922.

Nietnera, Green.

N. pundaluoya, Green 455 C. of C. V. 1922; on *Litsea*, on cinnamomum, etc., different places in Ceylon.

Sub-Family.—MARGARODINÆ

Kuwania Ckll.

Monophlebus zeylanicus, Green* (1896) is brought under the above genus, Green, p. 425 C. of C. V., 1922.

The Species is thought to be nearer the Margarodinæ.

Sub-Family—TACHARDINÆ

Chamberlin has recently revised the taxonomy of this group. He erects it into a family called Lacciferidæ. According to him the Indo-Ceylonese species are arranged as under, *vide* Bulletin. Ent. Res. xvi. 1925.

Laccifer, Oken.

L. albizziæ, Green } p. 340, Ramakrishna Aiyar. Check list of Coccidæ,
L. fici, Green } 1921.
L. lacca, Kerr }
L. ebrachiata, Chamberlin. On *Ficus*, Bangalore. *Vide* above reference.

Metacardia, Chamberlin

M. conchiferata, Green, p. 407. *Coccidæ of Ceylon*, vol. v, 1922. On *Mimosa*, *Anona*, croton—Kandy, Jaffna Matale, Ceylon.

Tachardina, Chamberlin

T. detorella, Maskell } p. 340, Ramakrishna Ayyar's list of 1921.
T. theæ, Green }
T. lobata, Green, p. 414, C. of C.V., 1922 and p. 340 of Ramakrishna Ayyar's list of 1921. On *Flacourtia* and *Fluggea* in Ceylon and on *Thespesea* and *Pongamia* in S. India. See also p. 208. Bull. Ent. Res. xiv. 1923.
T. ternata, Chamberlin. Included as doubtful species are *Mahdihassan's*, *silvestrii*, *pagoliensis*, *sindica mysorensis*, *communis*, and *chinensis*.
T. minuta, Mor, does not occur in India.

Sub-Family—ERIOCOCCUS

Eriococcinae

The genus *Kermes* Boit is brought under the above sub-family by Green p. 345, C. of C.V., 1922.

Eriococcus, Targ.

- E. nuerae*, Green, p. 349, C. of C.V., 1922. On unknown tree, Nuwara Eliya.
E. bambusae, Green, p. 350, C. of C.V., 1922. On bamboo, Udagama.
E. transversus, Green, p. 351, C. of C.V., 1922. On *Arundinaria*, Maskeliya.
E. tenuis, Green, p. 351, C. of C.V., 1922. On Grass. Pundaluoya.
E. rhodomyrti, Green, p. 352, C. of C.V., 1922. On *Rhodomyrtus*, Nuwara Eliya.
E. osbeckiae, Green, p. 353, C. of C.V., 1922. On *osbeckia*, Namunakuli.

Pseudopulvinaria, Atk.

- Lefroyia castenea*, Green (1908)* = *Pseudopulvinaria sikkimensis*, Atk, (1889)* and is brought under *Eriococcinae*, Green, p. 345, C. of C.V., 1922.

Sub-Family—ASTEROLECANIINÆ

Asterolecanium, Targ.

- A. minutum*, Green (Ms.) On bamboo, Dhony, Malabar.
A. gutta, Green, p. 1035, B.J., xxviii. 1922. On *Calophyllum walkeri*. Pattipola, Ceylon.
A. loranthe, Green, p. 1036. Do. do. On *Loranthus*. Hakgala, Ceylon.
A. pseudomiliaris, Green, p. 1036, B.J., xxviii. 1922. On Bamboo. Peradeniya, Ceylon.
A. lineare, Green, is renamed *A. lanceolatum* by Green p. 461, C. of C.V., 1922 as the former name appears to be preoccupied.

Polea, Green

- This new genus is erected for Green's *Pollinia ceylonica** (1909), p. 461, C. of C.V., 1922.

Cerococcus, Comst.

- Eriococcus paradoxa*, Msk., var *indica* Green = *Cerococcus indicus*, Green, these two are inserted separately in the previous list by mistake.

Anomalococcus, Green.

- A. hirsutus*, Green: (Ms.) On a wild tree. Sengleteri, Tinnevely.
A. cremastogastri, Green. This is given in the Pusa list¹ as from Coimbatore. This identification appears doubtful as our species is not *cremastogastri* but *indicus* Green. (Ms.)

Sub-Family—*Dactylopiinae**Halimococcus*, Ckll.

- H. borassi*, Green, p. 360. C. of C. V., 1922. On palmyra fronds, Peradeniya.

Geococcus, Green.

- G. radicum*, Green, p. 262. E.M.M., vol. 38. 1902 also p. 361, C. of C. V., 1922, on grass roots. Pundaluoya.

Pseudantonina, Green.

- P. bambusae*, Green, p. 363, C. of C. V., 1922. On bamboo. Nuwara Eliya.

Pedronia, Green

- P. strobilanthis*, Green, p. 364, C. of C. V., 1922. On *Strobilanthes*. Nuwara Eliya.

Erioides, Green.

- E. cuneiformis*, Green, p. 365, C. of C. V., 1922. On *Eugenia oligantha* and *Calophyllum*, Nuwara Eliya.
E. rimulæ, Green, 366, do. do. *Phyllanthus* bark and *Eugnymus* sp.; Nuwara Eliya

¹ p. 346. Rpt. of 5 Ent. Meeting, Pusa, 1923.

Tylococcus, Newstead.

T. formicarii, Green, p. 368, C. of C. V., 1922. From nests of *Cremastogaster* on *Grewia*. Maha Illuppalama, Ceylon.

T. simplicior, Green, p. 369, C. of C. V., 1922, on unknown plants with the ant *Ecophylla*, Maha Illuppalama, Ceylon.

[NOTE :—Both the above species are now referred to the genus *Farinococcus* of Morrison].

Pseudococcus, West.

P. kandyensis, Green, p. 373, C. of C. V., 1922. On grass. Kandy.

P. pulverarius, Newst, p. 373. Do. do. do. Pundaluoya.

P. pulverarius, sub sp. *bambusæ*, Green, p. 374, C. of C. V., 1922. On bamboo. Pundaluoya.

P. citriculus, Green, p. 377, C. of C. V., 1922. On leaves of *Citrus*.

P. monticola, Green, p. 378. Do. do. On elephant grass, Peradeniya.

P. corymbatus, Green, (*Ms.*) and *P. nipæ*, Green, on Casuarina, potato, etc. Maskeliya.

are = *P. filamentosus*, Ckll. var *corymbatus*, Green, vide p. 379, C. of C. V., 1922.

P. crotonis, Green,¹ = *P. lilacinus*, Ckll, Green, p. 380, C. of C. V., 1922.

P. maritimus, Erhorn, p. 315, Canad. Ent., 1900, and Green, p. 384, C. of C. V., 1922. On *Passiflora* and Tomato. Pundaluoya.

P. comstocki, Kuwana, p. 52. Proc. of Cal. Ac. Sc., iii, 1902. Green, p. 386. C. of C. V., 1922, on *Loranthus*, Pundaluoya.

P. debregeasiae, Green, p. 388, C. of C. V., 1922. On *Debregeasia*, Pundaluoya.

P. detorquens, Green (*Ms.*) On bamboo, shoots, Walayar, Malabar.

Ferisia, Fullaway.

Pseudococcus virgatus, Ckll. is brought under this new genus by Fullaway, p. 311, Proc. Haw. Ent. Soc., v. 1923.

Phenacoccus, Ckll.

P. glomeratus, Green, p. 389, C. of C. V., 1922. On *Hibiscus esculentus* and *Thespesia*. Peradeniya.

P. ornatus,² Green, p. 392, C. of C. V., 1922. On *Jasminum sp.* Ceylon. Also recorded by the writer on Jasmine from S. India.

P. spinosus, Robinson, p. 145, Philip. Jour. Sc. xii, 1918 and p. 394. Green, C. of C. V., 1922, on *Ficus*, Peradeniya.

P. hirsutus, Green. Has been recently noted in S. India on *Ficus sp.* and in shoots of feak ; rather badly on the latter—Walayar.

P. ballardi, Newst.=*P. mangiferae*, Green.

Antonina, Sign.

A. maritima,³ Green, p. 396, C. of C. V., 1922. On *Cyperus*, Colombo.

A. bambusæ, Msk., p. 237, N. Z. Tr., xxv, 1892 and Green, p. 397, C. of C. V., 1922. On Bamboo. Peradeniya, also found in S. India. Nilgiris. (E. E. Green Coll.)

Lachnodius, Msk.

L. humboldtiæ, Green, p. 400. C. of C. V., 1922. on *Humboldtia*, Yatiyantota, Ceylon. This species is to come under *Lachnodiella*, Hemp, according to Morrison.

¹ p. 46, Proc. M. S. Nat. Mus., vol. 60, 1922.

² pp. 41 and 45. A contribution to our knowledge of S. Indian Coccidæ by T. V. Ramakrishna Ayyar, published as Bull. No. 87 of the Pusa Agrl. Res. Inst., 1919.

³ pp. 41 and 45. A contribution to knowledge of S. Indian Coccidæ by T. V. Ramakrishna Ayyar, published as Bull. No. 87 of the Pusa Agrl. Res. Inst., 1919.

Xenococcus, Silvestri.

- X. annandalei*, Silvestri, p. 311, Rec. Ind. Mus., xxvi, 1924; in nest of ant (*Acropyga acutiventris*) in roots of *Ficus*, *sp.*, Madras.

Sub-Family—LECANIINÆ

Aclerda, Sign.

- A. japonica*, Newst. This has recently been noted on sugarcane in Salem, S. India.
A. ischæmi, Green, (Ms.) On *Ischæmum hirsutum*, Sengleter, S. India.

Cribrolecanium, Green.

- C. formicarium*, Green, p. 639, A. M. N. H., viii, 1921. On *Stereospermum* branches with ant. *Cremastogaster* *sp.* Ceylon.
C. radicola, Green, p. 642, A.M.N.H., viii, 1921. On roots of *Acacia auriculiformis*, Coimbatore. The food plant given as *Cassia* was the writer's mistake.

Lecanium, Burn.

- L. mangiferae*, Green.* On mango, Coimbatore. The first record for Continental India.
L. bicruciatum, Green.* On mango, Tinnevely. First record for Continental India.
L. desolatum, Green, p. 1020, B.J., xxviii, 1922. On *Ficus gibbosa*. Peradeniya.
L. fusiforme, Green, do. do. On unknown plant. Ambalangoda.
L. illuppalamæ, Green, p. 1021, B.J., xxviii, 1922. On unknown plant. Mahaiillupalama.
L. ixoræ, Green, p. 1022, do. do. On *Ixora Coccinæ*. Henaratgoda.
L. latioperculatum, Green, p. 1022, do. do. On a wild shrub. Peradeniya. Recently noted by the writer on mango in Coimbatore, S. India.
L. limbatum, Green, p. 1023, B.J., xxviii, 1922. On *Ixora coccinea*, Baticaloa, Ceylon.
L. mancum, Green, p. 1023. On *Calophyllum*. Badulla, Ceylon.
L. piperis var *namunakuli*, Green, p. 1024, B.J., xxviii 1922. On *piperis*. *sp.* Badulla, Ceylon.
L. tessellatum var *obsoletum*, Green, p. 1024. On *Myrtus*. Matale.
L. tessellatum,* noted on palm in hot house. Nilgiris.
L. trifasciatum, Green, p. 1024, B.J., xxviii, 1922. On *Hemicyclea*. Baticaloa.
L. tripartitum, Green, p. 1025, do. do. On *Calophyllum* Badulla.
L. ramakrishnae, Green, (Ms.) noted on pear. Kulu, N. India, vide Pusa list.
L. marsupiale, Green,* noted pretty bad on *Ficus indicus*, Kollegal. S. India.
L. imbricans and *Hemilecanium imbricans*,* Green, are same and not different as would appear from Pusa list.

Lecanopsis, Targ.

- L. ceylonica*, Green, p. 1026, B.J., xxviii, 1922. Pattipola. Ceylon.

Exaeretopus, Newst.

- E. farinosus*, Green, p. 1027, B.J., xxviii, 1922. On *Psychotria bisulcata*, Badulla.

Ceronema, Msk.

- C. fryeri*, Green, p. 1028, B.J., xxviii, 1922. On unknown plant, Maha Illupalama.
C. iceryoides, Green, p. 1029, do. do. do. Puttalam.

Ctenochiton, Msk.

- C. cinnamomi*, Green, p. 1030, B.J., xxviii, 1922. On *Cinnamomum*. Colombo.
C. fryeri, Green, p. 1031, do. do. On unknown tree. Vavuniya, Ceylon.
C. olivaceum, Green p. 1032, do. do. On *Pterospermum suberifolium*. Matale, Ceylon.

Cardiococcus, Ckll.

- Inglisia bivalata*, Green and *I. castilloa*, Green, are brought under this genus, vide Green, p. 1034, B.J., xxviii, 1922; the former was recently found badly infesting *Pongamia glabra* shoots in Coimbatore, S. India.

Lecaniodiaspis, Targ.

- L. mimusopis*, Green, p. 1034, B.J., xxviii, 1922. On bark of *Mimusops hexandra*. Hambantota, Ceylon.

Pulvinaria, Targ.

- P. polygonata*, Ckll. Pusa list, p. 347, Rept. 5th Ent. Meetg. On mang. Pusa.
P. cellulosa, Green, do. do. do. do.
P. maxima, Green, a memoir on—Ramakrishna Aiyar: Pusa. Ent. Mem. 1925; Vol. viii (12) pp. 127-155.

Takahashia, Ckll.

- T. japonica*, Ckll. Pusa list, p. 347, Rept. 5th Ent. Meetg. On Mulberry. Bhimtal, N. India.

Neolecanium, Parr.

- N. crustuliforme*, Green. This is referred to the genus *Platysaissetia* by Cockerell, vide p. 461, Green, C. of C. V. 1922.

Sub-Family—DIASPINÆ

Chionaspis, Sign.

- C. gynandropsidis*, Green, p. 1017, B.J., xxviii, 1922. On *Gynandropsis*. Peradeniya.
C. acuminata var *atricolor* Green, p. 1017, B.J., xxviii, 1922. On *carissa*. Maha Illuppalama.
C. linearis, Green, p. 1018, B.J., xxviii, 1922. On bamboo leaves. Peradeniya.
C. tenera, Green, p. 1019, do. do. On unknown shrub. Mahaluppalama.
C. pusa, Green, (Ms.) Pusa list, p. 348, Rept. of the 5th Ent. Meetg. On orange leaves, Pusa.
C. centripetalis, (Ms.) Green, do. do. do. On apple leaves. Kashmir.
C. spp. megaloba, Green, and *Hedyotidis*, Green, are recorded from N. India (Pusa list) from Pusa and Kangra respectively.
C. longissima, Green (Ms.) recently noted on bamboo. Dhony, Malabar.
C. elongata, Green,* recently recorded on bamboo. Dhony, Malabar; first record for India.
C. herbæ, Green.* On grass. (Coonoor, S. India.) Do. record for India.

Hemichionaspis, Ckll.=*Pinnaspis*, Ckll (an earlier name).

- P. buxi*, Bouche. In Pusa list on *Dracaena* leaves. Bombay.
P. ramakrishnae, Green. (Ms.) on *Eugenia Calophyllifolia*. Coonoor, S. India.
P. bauhiniae, Green (Ms.) On *Bauhinia racemosa* Papanasam, Tinnevely.

Neoleucaspis, Nov. gen. Green (Ms.)

- N. parallela*, Green. (Ms.) recently found by the writer on bamboo. Dhony, Malabar.

Diaspis, Costa.

- D. barbatus*, Green, (Ms.) on *Ischemum hirsutum*. Sengleteri, Tinnevelly.
D. boisduvali, Sign, p. 342. Ann. Soc. Ent. Fr., ix, 1869. Green, p. 464, C. of C. V., 1922, on Orchid, Maskeliya noted in S. India also on Orchid. Ootacamund.
D. antiquorum, Green, p. 1011, B. J., xxviii, 1922, on *Euphorbia antiquorum* with *Parlatoria mangiferae*. Green. Elephant Pass, Ceylon.
D. bambusæ, Green, p. 1012, B. J., xxviii, 1922. On bambo. Yatiyantota, Ceylon.
D. heneratgoda, Green, p. 1012 do. On unknown tree. Heneratgoda.
D. orientalis, Green, p. 1013 do. On *Sapindus*. Sigiriana. Ceylon; recently noted by the writer on *Hemigyrosa*. Tinnevelly, S. India.
D. phœnicis, Green, p. 1014, B. J., xxviii, 1922. On *Phoenix zeylanica*. Mahailuppalma.
D. mihiriya, Green, p. 1014 do. On *Dichopsis*. Bogawantalawa.
D. grandilobis, Green, p. 1015 do. On *Diospyros*. Peradeniya.
D. amygdali, Tryon, = *D. pentagona*, Targ. (Prior name) vide p. 460, Green, C. of C. V., 1922. Noted on cherry stem, Assam vide, Pusa list.
D. calyptroides, Costa and *D. echinocacti*, Bouche are synonyms, though they are put down as separate in the Pusa list, and Bouche's name has the priority.

Fiorinia, Targ.

- F. fenestrata*, Green (Ms.) On *Elæocarpus*. Nilgiris.
F. kandyensis, Green, p. 1016, B. J., xxviii, 1922. On unknown shrub. Kandy.
F. secreta, Green. Leonardi brings this under his new genus *Adisconfiorinia*, vide p. 52 Genera of *Diaspinæ Fioriniæ*, 1906.

Aspidiotus, Bouche.

- A. ambalangoda*, Green, p. 1007, B. J., xxviii, 1922. On unknown plant. Ambalangoda, Ceylon.
A. calophylli, Green, p. 1008. do. On *Calophyllum*, Badulla.
A. calophylli, Green, var. *symplocos*. Green, (Ms.) *Symplocos* leaves Nilgiris.
A. tripinnatus, Green (Ms.) On *Callistemon* leaves, Coonoor.
A. longispinus, Green.* Recently noted on *Bauhinia racemosa*. Tinnevelly, S. India.
A. pinnulifera, Msk. Var *diversicolor*, Green. On Orchid *Cymbidium*, Nilgiris.
A. perniciosus, Comst., p. 304, U. S. Agrl. Report, 1880. On pear, Punjab, vide Pusa list.

Aonidia, Targ.

- A. columnifera*, Green, p. 1008, B. J., xxviii, 1922. On *Turpina* leaves Hakgala.
A. mesochitinosa, Green, p. 1009 do. do. On *Canthium montanum*. Hakgala.
A. mimusopis, Green, p. 1009 do. do. On *Mimusops hexandra*. Trincomalai.

Gymnaspis, Newstead.

- G. diospyros*, Green (Ms.) On *Diospyros embryopteris*. Mundanthorai. S. India.
G. producta, Green (Ms.) do. do. do. S. India.
G. affinis, Green (Ms.) On *Ficus bengalensis*, Sivalapperi, S. India.
G. mangiferae, Green (Ms.) On mango leaves do. do.
Aonidia bullata, Green,* is *Gymnaspis bullata*, Green, vide Green, p. 460, C. of C. V., 1922.

Lepidosaphes, Schimer.

- L. albizziæ*, Green (Ms.) On stem of *Albizia lebbeck*. Coimbatore.
L. leucooplœæ, Green (Ms.) On stem of *Accacia leucophlœa*. Coimbatore.
L. punctatissima, Green (Ms.) On *Albizia lebbeck*. Coimbatore.
L. puniceæ, Green (Ms.) On Pomegranate. Coimbatore.
L. dilatilobis, Green, p. 1010, B. J., xxviii, 1922. On unknown shrub. Sigiriya, Ceylon.

Pseudoparlatoria, Cockerell.

P. pusilla, Green; p. 1010, B. J., xxviii, 1922. On cacao. Peradeniya.

Parlatoria, Targ.

P. camelliae, Comst. Recently recorded from South and N. India on *Melia* and *Aegle*, respectively.

P. cristifera, Green (*Ms.*) On *Citrus* leaves. Maddur, Mysore.

P. limoniae, Green (*Ms.*) On *Limonia alata*. Tinnevely.

P. mesuae, Rth., p. 266, Bull. Ent. Res., v., 1914. On *Mesua ferrea*. Peradeniya.

P. zeylanica, Rth., p. 113, Bom. Journ., xxiv, 1915. On bamboo, Peradeniya.

P. zeylanica, Rth., p. 114 do. do. On *Cinnamomum* Peradeniya.

For this latter species a new name *P. rutherfordi* is proposed by Green, as Rutherford has given same name to two species, *vide* p. 1020, B. J. xxviii, 1922.

P. cinnamoma, Rth., p. 114, B. J., xxiv, 1915. On *Cinnamomum*. Peradeniya.

P. blanchardii, Targ., *vide* Pusa list. On date. Lyallpur.

P. proteus var *crotonis*, Dougl., *vide* Pusa list. On croton. Pusa.

P. cingala var *namunakuli*, Green, p. 1019, B. J., xxviii, 1922. On unknown plant. Badulla.

A NOTE ON THE BIRDS OF KULU

BY

HUGH WHISTLER, F.L.S., F.Z.S.,

Indian Police.

From the administrative point of view Kulu is the Head-quarters Tehsil of the Kulu Sub-Division of the Kangra District of the Punjab. This Sub-Division includes four main areas, namely, Kulu, Saraj, Spiti and Lahul.

From the point of view of Natural Geography we may consider Kulu as the Upper basin of the River Beas.¹

The Beas rises on the Rhotang Range of the Central Himalaya at about 13,000' and its main tributaries are the Parbati, the Sainj, and the Tirthan which all enter it before the river passes from Kulu territory into that of Mandi State.

The whole basin is enclosed by very high ranges with a mean elevation of 18,000' which separate it from the Spiti, Chenab and Ravi Valleys; spurs from these ranges extend down to the Banks of the Beas; a lower range which in its ramifications largely makes up the Saraj Tehsil, separates the Beas basin from the Valley of the Sutlej.

The net result is a mountain area of the wildest character and most beautiful scenery. The lowest altitude is about 3,000' at the Larji Gorge where the Beas enters Mandi State: from there the main valley rises slowly to Manali 6,000', the fertile cultivation and level spaces of the lower parts about Bajaura gradually contracting until the Beas has become a rushing mountain river fringed with beautiful Alder beds and frowned upon by rocky precipices.

Cultivation occupies a very small extent of the area; the greater part of the country is clothed with magnificent forests, while the upper portions of the higher ranges are bare and desolate, wreathed in places with perpetual snow.

The avifauna is purely Himalayan in character with the exception of a small patch of cultivated country about Bajaura 3,600'. Here there are small numbers of species which are typical of the plains of India such as *Corvus splendens*, *Cisticola juncidis* and *Anthus rufulus*, which have somehow colonized this remote area amongst the mountains.

My acquaintance with Kulu is of long standing, having commenced in the year 1910. It has never been my privilege to live in the place but I have altogether been there six times, twice in early winter (November-December) and four times in summer (May-August). This of course is in no way sufficient to allow me to write a proper account of the local avifauna but I have ventured to place my notes on record, so that they may be available as a guide to the casual visitor, or a starting point to any one luckier than myself, who may live in Kulu and prepare a full and detailed account of the birds of a most interesting area. A considerable number of local specimens of birds and eggs are in my collection.

In the main these notes refer to Kulu proper but any observations available regarding Saraj have also been included. I have already published in the pages of the *Ibis* my notes on the avifauna of Lahul and Spiti, the other cantons of the Kulu Sub-Division.

M. Babault who was collecting in India and Ceylon on behalf of the Paris Museum in 1914 visited the valleys of the Beas and Parbatti on his way to Ladakh. His observations have been published in the *Résultats scientifiques du Mission Babault* and the pertinent portions have been included here for the benefit of English readers.

¹ *The Normal Rainfall in the Kulu Sub-division is as follows:—*

	Annual.	Jan.	Feb.	Mar.	Ap.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
Banjar...	41.54	3.06	2.89	3.22	2.34	2.27	3.64	9.03	8.99	3.58	0.81	0.56	1.15
Kulu ...	39.19	4.31	4.18	4.50	3.07	2.17	2.34	5.93	6.25	3.33	0.98	0.62	1.46
Nagar...	49.11	5.17	5.38	6.18	3.70	2.55	2.80	7.82	7.73	4.06	1.03	0.88	1.81

A few notes by other observers have also been incorporated into this account, including a list of species furnished by Mr. C. H. Donald to the local Gazetteer.

***Corvus corax tibetanus* Hodgs.** The Tibetan Raven.

M. Babault includes the Tibetan Raven amongst the list of birds that he met with at Tchari Djony between May 27 and June 2, that is to say about the border between Spiti and Kulu. It is of course fairly common in Spiti and Lahul. According to Stoliczka 'only very few come down in winter to Kulu.'

***Corvus coronoides intermedius* Adams.** The Jungle Crow.

The Jungle Crow is a common and resident species throughout Kulu, occurring alike in village, forest and alpine pasture. I have seen it following caravans to the summit of the Rhotang Pass 13,000 ft. and it may be said to go wherever man goes, occupying in this mountain area the position of the parasite on man that *Corvus splendens* is in the plains. Nests with young were observed in May and June.

***Corvus splendens* Vieill.** The House Crow.

It is curious that there is an isolated colony of this Crow in the Kulu Valley about Bhui and Bajaura: here the species is common for a stretch of about 10 miles, and I have seen a single individual (August 6, 1923) by the Beas a mile above Sultanpur. These birds are resident and have eggs about the middle of June. In the surrounding parts of Kulu they appear to be unknown and I have seen none in Mandi or in the Kangra Valley with the exception of a few birds at Dhelu 4,000 ft. (November 23, 1923).

M. Babault obtained 2 males at Bajaura and attributes them to the race *C. s. zugmayeri*, but these Kulu birds appear to me rather to belong to the typical race. The valley about Bajaura is the habitat of several other familiar birds of the plains, it is interesting to note, such as *Dicrurus macrocercus*, *Anthus rufulus*, and *Cisticola juncidis*.

***Corvus monedula* (L.).** The Jackdaw.

Hume states (N. and E. 2nd ed. I. 12) of the Jackdaw 'I have seen it in the hills in summer as far east as the Valley of the Beas.' It is a pity that he never amplified this record as it has been the basis of a good deal of error regarding the breeding range of this bird, which I have already discussed in the *Journal* (xxix. 161). In any case the occurrence must have been exceptional as in the course of six visits to Kulu, both summer and winter, I have never met with a single Jackdaw.

***Urocissa melanocephala occipitalis* (Blyth).** The Red-billed Blue Magpie.

M. Babault seems to have met with this Magpie fairly commonly in May and June, about the Dulchi Pass, Bajaura, Tchong, and Naggar, and he preserved 4 specimens. This is a slight extension of the known range which was hitherto considered to be bounded on the west by the Sutlej Valley. Although on the look out for the species I have personally failed to meet with it in Kulu.

***Urocissa flavirostris cucullata* Gould.** The Yellow-billed Blue Magpie.

Gould's type of this race was obtained in the Kulu Valley and is now in the British Museum. This Magpie is one of the most common and noticeable birds in Kulu, with its beautiful colour, long graceful tail, and remarkable range of notes. It is found everywhere up to about 7,000 ft. and is particularly common in the Alder groves along the Beas. I have never found the nest but have seen fledged young on June 25.

***Dendrocitta formosæ occidentalis* Ticehurst.** The Himalayan Tree Pie.

I have observed this sombre looking Treepie in Kulu as follows:—A party at Aramghar (5,000 feet) and one at Dhobi on November 30, 1922: a party including young of the year at Aramghar on July 9, 1923; a couple at 5,000 feet in the Parbatti Valley above Jhari on November 10, 1923.

It is common along the Mandi Valley from Mandi to Dhelu.

***Garrulus lanceolatus* Vigors. The Black-throated Jay.**

M. Babault obtained a male at Jhari and a pair on the Dulci ; Pass in May or June. I have observed it as follows :—One on the road at Jaggatsukh on June 18, 1921 ; one by the Beas road at Largi on June 29, 1921 ; one near Katrain on December 6, 1922. It appears to be less common in Kulu than along the outer ranges of the Western Himalayas.

***Nucifraga caryocatactes hemispila* Vigors. The Himalayan Nutcracker.**

Hume says (N. and E. 2nd ed. I. 30) 'my people have taken the nest with young in April below the Jalauri Pass.' M. Babault obtained specimens at Manikaran, Tcho-ti and Pulga in May and June, full feathered young of the year being shot on May 19, at Pulga.

The Nutcracker is a common bird in the Pine Forests of Kulu from 5,000 to 9,000 feet according to the late General Osborn (Journal, B.N.H.S. xv., 712) who says 'in fact it follows the region of the *Pinus excelsa*, the seeds of which furnish it to a great extent with food. The volume of the Journal cited above, contains several notes on a controversy whether the Nutcracker or the Flying squirrel was responsible for damage done to walnuts, which ended in the blame being laid at the rodent's door.

I have observed the Nutcracker in many localities in Kulu. In summer it may always be heard in the Silver Fir Forest about 7,000–9,000 feet on the Bhubhu Pass. I have heard it in June at 9,000 feet in the Nullah below the northern face of the Jalouri Pass. In July at 7,000 feet on the Dulchi Pass, in August above Manali on the Kothi road at 7,000 feet, and in July at 9,500 feet at the entrance to the Hampta Nullah. In November 1923 I saw many pairs from Manikaran to Pulga, 5,000–9,000 feet. In December I have heard it from 6,500 to 8,000 feet behind Katrain and Baragraon.

***Pyrrhonorax pyrrhonorax* (L.). The Common Chough.**

M. Babault observed this species at Tchari-Djony at the end of May, in great numbers at Kothi on June 28, and on the Rhotang on June 29.

During the summer months the Chough may be observed commonly in pairs or in flocks on the alpine pastures from the Beas tunnel at Kothi (8,000 ft.) to the summit of the Rhotang Pass, 13,000 ft. Here I believe they breed here as on May 25, a couple of birds emerged from stick nests in the face of a cliff by Rahla Rest House, which however proved to be quite inaccessible.

They are doubtless common on similar elevations throughout Kulu though I have only the following records :—June 8, 1919, Malasu Nullah a few over the snow ; June 16, 1922, a small party flying high over the Jalori Pass ; September 15, 1923, some at Serahan ; October 31, 1923, 3 on road at summit of Bhubhu Pass. Presumably during the winter they move down to the bottom of the valleys though there is no record on the point.

***Pyrrhonorax graculus* (L.). The Alpine Chough.**

M. Babault obtained a male at Tchari-Djony at the end of May.

According to Stoliczka this chough (which is so familiar in Lahul and Spiti) is in the cold weather specially common in Kulu.

***Parus major kaschmiriensis* Hart. The Kashmir Grey Tit.**

This Tit is a common and resident species along the valleys at altitudes of from 3,000 to 6,000 feet from Manali to Larji along the Beas and in the side valleys at similar elevations. I found a nest with young in the trunk of an alder tree at Raisan on June 19.

***Parus monticola* Vigors. The Green-backed Tit.**

The Green-backed Tit is common in Kulu in a slightly higher zone than the last species, starting in summer at about 5,000 feet. I found a nest with feathered young in a stump at Manali on May 22.

***Lophophanes melanolophus*. (Vig.). The Crested Black-Tit.**

This cherry little Tit is common in summer in the highest forests from about 6,000 to 10,000 feet ; during the winter it descends to a lower zone and then is one of the most prominent members of the hunting parties that are such a feature of Himalayan bird life. I found a nest with naked young on May 25, in a large

hole high in the trunk of an oak tree by Rahla Rest House, and another nest with 5 moderately incubated eggs on June 16, in the trunk of a Yew-tree by Shoja Rest House.

There is little to note on this bird's behaviour; he is always busy in the search for food high in some moss-grown giant of the forest, and the soft gold-crest-like note 'chee chee' will be heard long before the tiny bird is seen in the branches above one: sometimes he is alone, at other times two or three wander about in loose companionship, and in winter flocks of up to 50 in number, collect and join the hunting parties. In spring they rest occasionally from the incessant hunt for food, and from the top of some lofty twig utter the loud clear call note 'want you need you want you need you' or deed you 'deed you deed you'.

Lophophanes rufonuchalis rufonuchalis. (Blyth). The Simla Black-Tit.

M. Babault records obtaining a male from Pine forest in the Valley of Kulu in May without giving more exact data.

I came across a party at 10,000 ft. on the northern slope of the Jalouri Pass on June 17, 1922.

It of course breeds commonly in the Junipir forests of Lahul and is probably more numerous in Kulu than the above records indicate.

Machlolophus xanthogenys xanthogenys (Vigs.). The Yellow-cheeked Tit.

M. Babault obtained a male on the Dulchi Pass on April 29. I have not met with the species in Kulu.

Egithaliscus niveogularis (Moore). The White-throated Tit.

On June 17, 1922, I secured 4 specimens, 2 adults 2 juveniles, from a family party in a patch of willow scrub; the young had clearly been hatched in the neighbourhood. This was at 9,800 ft. on the Kulu side of the Jalouri Pass not far from the summit.

On November 16, 1923, I almost certainly saw one of these tits in some bushes at 4,800' by Banjar but a dog disturbed it before I could verify the identification beyond doubt.

Egithaliscus concinnus iredalei Stuart Baker. The Red-headed Tit.

The Red-headed Tit is fairly common in Kulu and I have met it in summer from 5,500' up to 7,500'. In winter it descends as low as river level though I have also seen it as high as 6,500' at that season. In all probability there is little seasonal altitudinal movement in this species.

Sitta himalayensis Jard. and Selby. The White-tailed Nuthatch.

I procured a male on June 17, 1922, at 6,500' near Jhibbi.

Sitta leucopsis leucopsis Gould. The White-cheeked Nuthatch.

M. Babault procured a pair at Pulga in May and June and remarks that it was more common at Tcho-ti than in the former place. In early November at Pulga I heard a Nuthatch very commonly in the pine forests from 7,000 to 9,000' and believe that it was this species.

On June 17, 1922, I saw one in company with a party of *Parus rufonuchalis* at 10,000' on the Kulu side of the Jalouri Pass.

Garrulax albogularis whistleri Stuart Baker. The White-throated Laughing Thrush.

M. Babault obtained two males and a female in the neighbourhood of Bajaura and the Dulchi Pass at the end of April, and he also found a nest containing 2 young birds with feathers commencing to sprout.

Trochalopteryx erythrocephala erythrocephala Vigors. The Red-headed Laughing Thrush.

Hume says of this Laughing Thrush (N. and Eggs. 2nd edn. vol. i, 55) apparently including Kulu:—'From Kumaon westwards, at any rate as far as the Valley of the Beas, next to *T. lineatum* the most common species of the genus'. I have however only seen a single individual in Kulu and that at Kasol on November 6, 1923.

Trochalopteron variegatum variegatum (Vigors). The Eastern Variegated Laughing Thrush.

This Laughing Thrush appears to be a common bird in Kulu, breeding on the higher ridges from 8,000 to 10,000 ft. and moving down in winter to a lower zone, but I have no very detailed information about it. In Lahul the western form *T. v. simile* occurs and the Rhotang Range is therefore probably one of the boundaries between the two races. It would be interesting to verify which bird is found in the Solang nullah and the forests generally along the southern face of the Rhotang.

Trochalopteron lineatum lineatum * (Vigors). The Nepalese Streaked Laughing Thrush.

The Streaked Laughing Thrush is one of the commonest birds of Kulu from Larji upwards to a height of at least 8,500 ft. in all the valleys. During summer it is scarce along the bottom of the valley from Sultanpur to Larji though some may always be found in the swamp between Bhuin and Bajaura, and in Kulu, as elsewhere, the bird is doubtless subject to a good deal of altitudinal movement according to season. It keeps much more to the ground than the other species of Laughing Thrush, and shuffles along in tangles of grass, bracken and other low herbage, flirting its wings and jerking the long tail, the members of a flock keeping up an incessant conversational squeaking in low querulous notes. It seldom ascends into trees and when disturbed from the undergrowth it merely flies a short distance down hill and pitches into the nearest cover into which it quickly threads its way out of sight.

Grammatoptila striata striata. (Vig.). The Striated Laughing Thrush.

M. Babault met with this large Laughing Thrush at Kothi on June 28. There appears to be no other Kulu record though as it is common at Koteghar in winter Stoliczka may have been right in his conjecture that these birds bred in the higher forests of Kulu.

Pomatorhinus erythrogenys erythrogenys Vigors. The Rusty-cheeked Scimitar-Babbler.

Hume includes the valley of the Beas in his account of the distribution of this species (N. and E. 2nd ed. vol. i, 87) but I have not met with the bird actually in Kulu. It is common in Mandi State and Kangra at elevations from 2,000 to 6,000 ft.

Stachyridopsis pyrrhops (Blyth). Red-billed Babbler.

The Red-billed Babbler is resident and apparently not uncommon in Kulu. I have met with it in June about 5,000 to 5,500' in the main valley from Sultanpur to Naggar, and in November at Sultanpur 4,000' and in the Parbatti Valley about Jari 5,000'.

Siva strigula strigula Hodgs. The Stripe-throated Siva.

I met a party of these Sivas on June 17, 1922, in some thick scrub at 9500 to 10,000' on the Kulu side of the Jalouri Pass: they doubtless breed here and in similar situations on other ranges. In winter they move lower down and I found a flock at 5,000' on the Kraon road on November 27, 1922.

Ixulus flavicollis albicollis Ticehurst and Whistler. The Yellow headed Ixulus.

There is a patch of Chestnut Forest at 5,500' on the road below Kraon where I have met with a party of the Ixulus in the same trees in May, July, August, and November. Otherwise I have only procured it in Kulu on December 5, 1922, from 7,000' in the hill-side jungle behind Katraian.

Pteruthius xanthochloris occidentalis Harington. The Green Shrike-Tit.

Stoliczka says 'Only 3 specimens were shot, in February 1867, in the southern part of Kulu.' There is no other record.

Microscelis psaroides psaroides (Vigors). The Himalayan Black-Bulbul.

In summer this Bulbul breeds throughout Kulu from Larji 3,000' up to 6,000' and doubtless higher in all the valleys and on the hill-sides. I have observed a few in November and early December at about 4,000' to 6,000' but the majority probably leave to winter in the outer ranges.

* *griscientior* of Hartert.

Molpastes leucogenys leucogenys, (Gray). The White-cheeked Bulbul.

A common and resident species in Kulu and Saraj from the bottom of the valleys up to a height of about 7,000 ft., I have found nests in June.

Certhia himalayana Vigors. The Himalayan Tree-Creeper.

A fairly common species throughout the Sub-division, usually seen with the mixed hunting parties except in the breeding season. It is a resident but moves altitudinally according to season.

Tichodroma muraria. (L.). The Wall-Creeper.

A common winter visitor to the sub-division where I have seen it in November and December along the valleys from 3,000' to 5,000'. A few perhaps breed at high altitudes along the Rhotang chain, but of this there is as yet no record.

The Wall-Creeper attracts the attention of the least observant in Kulu by its habit of feeding about the cuttings and retaining walls of the roads, climbing over their surfaces with a jerky mode of progression and a continual downward flick of the outer primaries. It takes to flight with reluctance before the passerby launching out into the air with a hovering undecided flight resembling that of the Hoopoe, a resemblance heightened by the long bill and broad round spotted wings, only to settle again a short distance ahead. It is also very partial to the large boulders that line the sides of the river beds, and it may often be found feeding on the stones close to the edge of the water. I have only once seen it climbing on a tree trunk. It occasionally flies up and endeavours to take an insect on the wing. Nowhere else have I seen this species so common as in Kulu.

Troglodytes troglodytes neglectus Brooks. The Wren.

In Kulu proper I have met with it in small numbers in November and December at heights from 5,000' to 8,000', and it is doubtless a resident in the sub-division breeding in a higher zone about the edge of forest growth.

This is one of the Himalayan birds that is most reminiscent to the exile from England, being, as it is, one of the few birds that is a race of an actual British species. It recalls the home bird in its demeanour and ways. Attention is drawn to it by the harsh and rather scolding metallic note 'chit chit chittr', as it flies from some tangle of herbage at the foot of a stunted bush to the litter of boulders that mark some forgotten landslide; it threads its way amongst the stones, now for a space disappearing into the hollows and crevices below them, then emerging on to some stone. There it waits for a moment, the tail cocked upwards and the head stiffly held back, often so far that the beak points almost directly upwards. It is intensely restless, a mass of nerves, incessantly bobbing and bowing, and turning first to one side and then to the other. Usually it is averse to taking flight but progresses amongst stone and hollow in little jerks and starts, with short flights of merely a foot or two; but once convinced that danger threatens it leaves for distant cover with a flight that is straight and strong and fairly fast. The song is loud and cheerful, as in the English bird, and is freely uttered in the bleakest of surroundings where drifts of snow coil amongst the debris of a rockbound nullah.

Pnoepyga albiventer pallidior Kinnear. The Scaly-breasted Wren.

In November and December I found this quaint little ball of a bird fairly common in the valleys of Kulu and Inner Saraj between 3,600' and 6,500'. It is a most arrant skulker keeping to thick bracken and evergreen undergrowth, preferably in the alder groves, in the dampest and most secluded situations. I have even found it in the water-logged tussocks of the Bhuin marsh. It creeps about close to the ground like a mouse and hardly ever takes to wing, and specimens may be secured only with the greatest difficulty. The call is a loud single note, quite unlike that of *Troglodytes*, but more warbler-like in character.

Tesia castaneocoronata castaneocoronata (Burton). The Chestnut-headed Wren.

The western limit of this beautiful wren is given in the Fauna as Garhwal but in November and December I found that it was not uncommon in the Kulu Valley about Sultanpur, and it was also reported to me from the Parbatti Valley. It was found in similar situations as *Pnoepyga a. pallidior* and has similar habits. The call note is a loud slightly double squeak, rather reminiscent of the note of *Phylloscopus humii*.

***Cinclus cinclus cashmerlensis* Gould.** The Kashmir Dipper.

M. Babault met with this species about Pulga and Garampani between May 17 and June 4. I have not met with it in Kulu though it occurs at the Chandra Lake 14,000' and in Spiti at 13,000' across the border.

***Cinclus pallasii tenuirostris* Pallas.** The Brown Dipper.

The Brown Dipper is one of the most characteristic of the birds of Kulu and Saraj, where it is found throughout the year on the Beas and its various tributary streams: during the summer it is found up to 9,000' and 10,000': but in the winter the birds become distinctly more common in the main valley and it is probable that they then leave the highest zone of their summer habitat. I have not been able to find the eggs but birds in the spotted juvenile plumage are to be seen about in May and it is probably an early breeder.

The Dipper is a most active bird, never still and always busy. The harsh call 'dzchit dzchit' is a familiar sound along every stream, shrill enough to be heard easily above the roar of the waters; it heralds the approach of the small plump brown bird that flies swiftly along a foot or two above the surface of the water, swaying from side to side amongst the boulders and only making a detour over land to avoid some intruder at the water's edge; the wings appear rather small for the stout body and to make up for this they are vibrated very quickly in flight in sustained beats followed by a pause.

Settling on a stone the bird bows and jerks from side to side, or immediately starts feeding, keeping its foot-hold easily on slippery stones and disappearing under water either diving or walking. It swims freely on the broader pools, looking like a miniature moorhen, now and again diving and disappearing for a space.

On one occasion I was watching a Dipper standing on a rock and preening its plumage when the mate arrived. The first bird immediately stretched itself high on its legs, with the tail pointing straight downwards, and the head and neck extended to their full length with the beak pointing to the sky; it stood thus for some time with the wings whirring rapidly, the whole doubtless forming a courting display.

***Larvivora brunnea* (Hodgs.).** The Indian Blue Chat.

M. Babault obtained a male at Pulga in May or June. I believe that I heard the song near Diwal village above Manal's at 7,000' on July 11, 1923.

***Hodgsonius phoenicuroides* (Hodgs.).** Hodgson's Short-wing.

M. Babault obtained 2 females at Tchari-Djony and noted the species about Pulga and Garampani at the end of May or beginning of June.

Mr. Wells kindly presented me with a male which he had shot from a nest containing 3 young (hatched on July 8, 1922) near Chikka 10,000' in the Jagatsookh Nala. The nest was a cup of coarse grass lined with hair, placed about a foot from the ground in a low bush amongst some large boulders. A second nest was also found.

***Saxicola caprata bicolor* Sykes.** The Pied Bush-Chat.

The Pied Bush-Chat is a summer visitor to the lower parts of the Kulu Valley and breeds in abundance up to a height of about 4,000', from Larji to Sultanpur and Manglaur.

***Saxicola torquata indica* Blyth,** The Indian Stone-Chat.

Hume states (N. and E. 2nd ed. ii. 48) that he has found numbers of the nests in the valley of the Beas below Bajaura, but he presumably means at higher elevations on the hillsides; for in my experience the Stone-Chat does not breed in the same area as *S. c. bicolor* but occupies a higher zone from 4,000' upwards, commencing in the valley only about Sultanpur and becoming common about 6,000' as at Manali and Banjar. M. Babault procured specimens at the foot of the Rhotang and in the Parbatti Valley.

During the winter the species is fairly common along the valley from Katrain to Bajaura.

***Oreicola ferrea ferrea* (Gray.).** The Western Dark-grey Bush-Chat.

Hume says (N. and E. 2nd ed. ii. 51), 'I have found many nests in Kulu . . . one near Sultanpur.' This Bush Chat breeds commonly through-

out Kulu and Inner Saraj at elevations from 5,500 to 8,000'. The majority probably leave the Sub-division in winter but in November and December I met with a few along the valley from Sultanpur to Bhuin.

***Oenanthe deserti* (Temm.).** The Desert Wheatear.

According to Stoliczka the Desert Wheatear occasionally winters in Kulu; if so it would be of interest to discover which race is represented by these birds.

***Enicurus maculatus maculatus* Vigors.** The Western Spotted Forktail.

The Forktail is a familiar species on the streams of the Kulu Sub-division being found at all elevation up to about 8,000'. It is a resident bird and probably not subject to any particular altitudinal movements according to season. It attracts the attention of even the least observant by the conspicuous black and white plumage, with the distinguishing white cross of St. Andrew across the back, and the long-forked tail which is swayed upwards and downwards vertically, slowly and with infinite grace, as the bird moves about the stones and banks of the streams that it frequents. The Forktail is not shy and continues to feed even in the near presence of an observer, sidling and turning on the stones, now facing one way, then turning round to the other. The call is loud and distinctive, and in the woodbound streams of the side nullah, is often the only indication of the bird's presence as it flies up or down the stream through the undergrowth disturbed by the approach of an intruder.

Babault found naked young in a nest on June 7.

***Microcichla scouleri scouleri* (Vigors).** The Little Forktail.

The Little Forktail appears to be not uncommon in Kulu and Saraj for I have met with at least 2 pairs at Banjar 4,500' in June, a pair below Manali at 6,000' one at Rahla 8,800' and one at Jhikka 10,000' (Hampta Nala) all in July. In November and December I have found it in the Beas Valley at Sultanpur 4,000', and in the Parbatti Valley at Jari, Kasol, and Manikaran 4,000-6,000'.

***Phoenicurus frontalis* (Vigors).** The Blue-fronted Redstart.

This handsome Redstart breeds at 13,000' amongst the rocky ledges of the summit of the Rhotang Pass, where I have observed it in June. In the same month it also breeds in the scrub on the northern face of the Jalouri Pass at 10,500'. Babault collected a pair in May or June at Ghary and says that he often surprised them in the bushes on the bank of the Parbatti, which seems curiously low for this species in summer. It doubtless breeds along the summit of all the high ranges in the Sub-division.

In November and December I have observed the species on the hillside behind Katraian and at Pulga 7,000'.

***Phoenicurus ochrurus phoenicuroides* (Moore).** The Black Redstart.

On July 27, 1922, I observed a few Black Redstarts below the summit of the Hampta Pass at 13,000' and they probably breed here as they do in Spiti and Lahul, where they are amongst the most characteristic birds of those high valleys. Babault collected a pair at Manikaran and a male on the Rhotang, but he gives no details to show whether the Manikaran birds were breeding or on passage. There are no winter records for Kulu so perhaps the Sub-division, like the Simla and Dharmasala ranges, lies outside the winter range and regular migration routes of this common and wide-spread species.

***Chaimarrhornis leucocephalus* (Vigors).** The White-capped Redstart.

During the breeding season the White-capped Redstart is common and breeds on the Beas from 7,000 ft., above Manali, to its source on the Rhotang Pass 13,000 ft., and doubtless at similar elevations on all the larger streams. I have found fresh eggs at Rahla on July 11, and on the same day another half-built nest, so it appears to be somewhat of a later breeder than most birds. At this season it is distinctly a bird of high elevations and may be found up to 16,000 ft., in Lahul and elsewhere. During the winter months the species moves down to a lower zone and the birds are then extremely common throughout the valleys of Kulu, and even lower down into the foot-hills of the Himalayas.

Like the Plumbeous Redstart, the White-capped Redstart is most strictly a water bird but it differs from the former in its wider interpretation of what the term 'water' includes: for in addition to the rivers, streams and rills, it is found about wet mossy cliffs, precipitous marshy banks and similar situations in which the Plumbeous Redstart does not occur. Like it, however, it is pre-eminently a bird of boulders amongst rushing water, and here the two species are found together, fly-catching with little erratic flights from stone to stone. As with most Redstarts, the tail is an expressive organ. Continuously the bird beats the tail up and down, from well above the line of the back almost to touch the stone on which it is sitting, frequently accompanying the action with a deep bow; and this is done with the tail either closed or partly spread. As it starts to fly or settles the tail is fanned open, a glorious glimpse of chestnut and black. The ordinary call note is a loud plaintive squeak 'teeee' audible some way off.

Rhyacornis fuliginosus (Vigors). The Plumbeous Redstart.

The Plumbeous Redstart is one of the commonest and most characteristic birds of Kulu and Saraj. It is never found away from the rivers and larger streams, and with that reservation, may be looked for anywhere between 4,000 ft. and 10,000 ft., in summer. It does not, however, care for the broader and hotter stretches of the Beas between Sultanpur and Larji. During the winter it leaves the higher zone of its range, and while at that season great numbers pass down to the waters of the foothills, it remains sufficiently common in Kulu up to at least 6,000 ft. It breeds throughout its range in May, June and July. The nests are built on ledges of rocks, in holes under earthen banks, in crevices between boulders and stones, in hollows in the trunks of Alder trees, in ivy on tree trunks and any similar situation, always provided that it is within a few yards of the water's edge.

The nest is a neat cup made of moss mixed with a few roots and leaves, and is lined with fine roots and fibres and occasionally wool and hair. The eggs are pale greenish in colour speckled and blotched fairly heavily with markings of reddish brown. They are usually four in number.

These graceful little birds strike the notice of even the least observant. No patch of stream is without its pair which spend all their time on the boulders in the middle of the rushing water, with occasional excursions to the bank or the bough of some handy tree. They flit from stone to stone and continuously make erratic little fluttering darts into the air after some passing insect; and as they settle the conspicuously coloured tail, chestnut in the cock, black and white in the hen, is slightly fanned and wagged up and down, the movement being repeated at intervals until the next little dart into the air. They are as quarrelsome as restless, and appear to have sharply defined territories, for the male with a provocative little snatch of song is always launching attacks at the intruder from some other pair, dashing at it and chasing it back to its own borders, regardless of sex. The short song is rather sweet and jingling and the hen also shares it; it may be heard occasionally in winter as well as in the breeding season. It is uttered either from some boulder in midstream or in the air as the little bird slowly flies with even movement but rapidly vibrating wings in a short parabola from one rock to another.

Cyanosylvia cyanecula abbotti Richmond. The Bluethroat.

A fine adult male Bluethroat was seen at Manali on May 23, 1921, at 6,000 ft. but I could not obtain it. From the bright colour of the breast I attributed it to this form.

Grandala caelicolor Hodgs. Hodgson's Grandala.

This rare and beautiful bird must breed at high elevations at the head of the Parbatti Valley, as Babault procured a male and female with the organs developed at Tchari-Djony between May 27, and June 2, 1914. It may be noted also that Stewart obtained some specimens from the collection of a Mr. Procknow, a missionary at Koteghar, which the latter said were 'strictly mountaineers and only got several marches in near the snow' (Zoologist) 1886, 439).

Calliope pectoralis pectoralis Gould. The Himalayan Rubythroat.

Babault obtained a male at Ghary on May 13,

Rubythroats breed on the ascent to the Rhotang Pass at about 12,000 ft. amongst boulders on the open hillside, a locality in which I have observed them in June, July and early August.

Tarsiger chrysaeus whistleri Ticehurst. The Golden Bush-Robin.

I saw a male at Jari 5,000 ft. on November 6, 1923.

Ianthia cyanura pallidiora Stuart Baker. The Red-flanked Bush-Robin.

In the middle of June I have found this species common and breeding from Shoja to the summit of the Jalouri Pass in the forest zone between 9,000 ft. and 10,000 ft., and it doubtless breeds on similar ridges throughout the Sub-division. One was seen at 6,500 ft. behind Katraian on December 3, 1922.

Adelura caeruleocephala (Vigors). The Blue-headed Robin.

In Nests and Eggs 2nd ed. vol. ii, 69, Hume records the taking of an egg of this species in the Humpta Nala at about 11,000 ft. He admits to the possibility of error, and the description certainly does not agree with eggs that I have taken in Lahul.

In winter this bird appears in the valleys of Kulu and Saraj and I have seen it in November and December from 4,000 ft. to 7,000 ft. both at Banjar and along the Beas Valley.

Saxicoloides fulicata cambaiensis (Lath.). The Brown-backed Indian Robin.

Stoliczka states that this familiar species is common in Kulu but I have not met with it; its existence there seems unlikely.

Copsychus saularis saularis (L.). The Magpie-Robin.

Observed in fair numbers at Sultanpur 4,000 ft. in May, June, and July, nests with feathered young being found on June 23 and July 29. A pair were seen at Manglaur Rest House, Inner Saraj 3,800 ft. on June 19, 1922. It is a summer visitor only and apparently leaves in August.

Turdus atrogularis Temm. The Black-throated Thrush.

A common winter visitor observed in the valleys in November and December.

Turdus unicolor Tickell. Tickell's Ouzel.

Only observed at about 6,000 ft. towards Manali, where I have seen one or two in May, June and July.

Geocichla wardi (Jerdon). The Pied Ground-Thrush.

I found a small flock of 5 or 6 individuals at 6,000' near Manali on May 24 and 25, 1921. They were feeding in brush-wood in a small wooded nullah by cultivation and flew up into the trees when disturbed.

On July 5, 1923, I met with a pair with newly fledged young in a similar spot by Kraun Rest House 6,000'.

Arceuthornis viscivorus bonapartei (Verr.). The Missel Thrush.

Babault met with the Missel Thrush on the Dulchi Pass between April 29, and May 1. Hume obtained the nest above Juggutsukh. On June 16, 1922, I saw 2 or 3 at 10,000' along the crest of the Jalouri ridge in the forests of Kharshu Oaks, whence they came to feed on the open summit of the ridge.

On December 8, 1922, 2 were seen in the Birch forest above Baragraon near Katraian.

Oreocincla mollissima (Blyth). The Plain-backed Mountain-Thrush.

Babault obtained a male at Tcho-Ti in May or June.

Oreocincla dauma dauma (Lath.). The Small-billed Mountain-Thrush.

I saw what was probably one of these Thrushes on the road below Manali 6,000' on May 22, 1921.

Monticola erythrogastra (Vigors). The Chestnut-bellied Rock-Thrush.

On November 30, 1922, I saw 2 males flying together about the tops of some Alder trees in the Aramghar marsh near Raisan 4,500'. Babault found the species at Ghary and Tchong in the breeding season.

Monticola cinclorhyncha (Vigors). The Blue-headed Rock-Thrush.

A common summer visitor noted in May, June, July and August from 5,000' to 6,000'.

Monticola solitaria pandoo (Sykes). The Blue Rock-Thrush.

On June 23, 1922, I found a nest with 4 partly feathered young in a rocky bank above cultivation in the valley below Sultanpur at 4,000'. A few birds noted about this area in June and July and a male seen at 6,000' above Manali on June 17, 1921, complete my Kulu records for the species. Babault however procured several specimens in May or June at Ghary, Manikaran, Pulga and Tcho-ti.

Myiophoneus temminckii temminckii (Vigors). The Himalayan Whistling-Thrush.

The Whistling Thrush with its deep blue black plumage and yellow bill must often be mistaken in Kulu by tourists and sportsmen for the English blackbird. It can indeed escape the notice of few as it is excessively common throughout the valleys up to 11,000' in summer and to a slightly lower elevation about 8,000' in winter, while its large size, bold demeanour and noisy calls invariably draws attention to it. Primarily it is a bird of the rivers and mountain streams, but its strong flight enables it to wander freely about the precipitous crags and heavy woods that line every valley, and there is scarcely any piece of Kulu scenery that the traveller will stop to regard without sooner or later seeing one of these birds dash across the picture, and without hearing the melodious whistle or the harsh loud song mingling with the roar of the waters and easily rising above it; there is a peculiarly eerie squealing character in the song which truly matches it with its surroundings.

There is something very tight-trussed about the appearance of the Whistling Thrush as he hops and flies from boulder to ledge, from the ground to the broad branches of the alders and chestnuts; his hard shiny feathers are pressed close to the body and as the long tail sways slowly upwards above the long legs he has all the appearance of balancing on tip-toe, like a smart recruit awaiting his sergeants' word of command.

The nest is easily found, for it is a large massive cup of the usual Blackbird type and the bird depends for its protection, not on concealment but on the inaccessibility of its site, placing it on some rocky ledge or in a hollow on the face of a mighty boulder overhanging a rushing stream. Eggs may be found from April to July; they are very fragile and in appearance resemble large and very faded specimens of Blackbird's eggs.

Prunella collaris (Scop.). The Alpine Accentor.

One was seen on the road just above Manikaran at 6,500' on November 10, 1923, but the specimen was not obtained for racial identification.

Prunella himalayanus (Blyth). The Altai Accentor.

On December 8, 1922, I found this Accentor common in parties on the Kharshu Tharch, behind Baragrown, about 9,000'. Some parties of birds seen in similar ground above Pulga on November 9, 1923, were also apparently of this species.

Hemicheledon sibirica gulmergi Stuart Baker. The Kashmir Sooty Fly catcher.

Breeds commonly throughout Kulu in the forests above 5,000' up at least to 8,500' and is a summer visitor only; an occupied but in accessible nest, 40 feet from the ground on a side bough of a silver fir, was seen on June 15 and fledged young were seen on July 1.

Siphia strophciata Hodg. The Orange-gorgetted Flycatcher.

Breeding birds were observed at 9,000-10,000' on the Shoja side of the Jalouri Pass on June 17, 1922.

Cyornis superciliaris (Jerd.). The White-browed Blue Flycatcher.

A common summer visitor to the forests throughout Kulu breeding from 6,000' to 10,000'. I found a nest containing newly-hatched young on May 24, placed in a hollow of the bark of an alder.

Stoparola melanops melanops (Vigors). The Verditer Flycatcher.

A summer visitor observed from the bottom of the valley up to 8,000' but not so common as in the outer Himalayan ranges.

Alseonax latirostris poonensis (Sykes.). The Brown Flycatcher.

Breeds fairly commonly along the Beas Valley from Sultanpur 4,000' to Larji 3,200' and in the Tirthan Valley to Manglaur 3,800'; a summer visitor only. Lays in May and June.

Alseonax rubicaudus (Swains.). The Red-tailed Flycatcher.

M. Babault obtained two males and two females at Pulga at the end of May and judged from their organs that they were breeding.

Culicicapa ceylonensis ceylonensis (Swains.). The Grey-headed Flycatcher.

A common summer visitor to Kulpu breeding in the Zone from 5,000' to 7,000'.

Niltava sundara sundara Hodgs. The Rufous-bellied Niltava.

I procured a female in the undergrowth at 7,000' on the ascent to the Bhubhu Pass on August 9, 1923. According to Hume (N. and E. 2nd ed. ii. 20) the species breeds in the valley of the Beas.

Tersiphone paradisi (L.). The Paradise Flycatcher.

The lovely adult male Paradise Flycatcher with his black-crested head, snowy plumage, and ribbon-like tail feathers is a familiar sight to the traveller along the roads of the valley. It is common from Larji up all the valleys to a height of about 5,000' but of course only as a summer visitor.

Lanius vittatus Valenc. The Bay-backed Shrike.

A common summer visitor to the Beas Valley between Larji and Bajaura breeding early in June.

Lanius nigriceps nigriceps (Frank.). The Black-headed Shrike.

M. Babault obtained a male at Pulga and a female at Bajaura in May or June.

Lanius schach erythronotus (Vigors). The Rufous-backed Shrike.

This fine shrike breeds very commonly in June along the valley of the Beas and Tirthan from Larji 3,200' to Banjar 5,000' and Sultanpur 4,000'; a few pairs extend up to Manali 6,000'. Odd shrikes may be observed in the valleys in winter but they may belong either to this form or to *Lanius tephronotus* which is the breeding shrike of Lahul (as a summer visitor only).

Lanius cristatus subsp. ? The Rufous-tailed Shrike.

Hume's remarks [(N. and E. 2nd ed. i. 327) imply that some form of this shrike appears in the valley of the Beas in the late autumn, on migration from Yarkand, but I have no more definite information on the point].

Pericrocotus brevirostris brevirostris (Vigors). The Short-billed Minivet.

The scarlet and black minivets with their grey and yellow mates attract the attention sooner or later of most travellers in Kulu from their conspicuous colouration and their pleasant calls. They breed throughout the forests of Kulu from about 5,000' to 8,000' and probably higher and in the spring and autumn may be seen anywhere in parties. The majority leave Kulu in winter but I have seen a few about Sultanpur, Kasol and Bhuin, in November.

Pericrocotus speciosus (Lath.). The Scarlet Minivet.

M. Babault obtained a female at Garampani, and a pair at Pulga in May.

Pericrocotus roseus roseus (Vieill.). The Rosy Minivet.

M. Babault obtained a pair at Bajura in May or June.

Dicrurus macrocercus Vieill. The King-Crow or Black Drongo.

The king crow is a summer visitor to the valley of the Beas and the Tirthan at elevations of 3,000-3,500' from below Sultanpur to Larji and Manglaur. Eggs may be found at the end of May and throughout June.

Dicrurus leucophæus longicaudatus Hay. The Ashy Drongo.

The Ashy Drongo is a common summer visitor to Kulu and Saraj from 3,000' up to 8,000' and probably even higher. The majority of eggs are laid in June.

Acrocephalus dumetorum Blyth. Blyths' Reed Warbler.

While marching through Kulu from May 17 to 25, 1921, I met with this species in some numbers from Kraun 6,000' to Sultanpur 4,000' and along the Beas up to Manali 6,000' along the road. It was then on the spring passage, and had for a month previously been similarly common in the Kangra Valley and at Dharm-sala. It was not however observed in Lahul which was obviously out of its line of passage, and by my return from Lahul to Kulu on June 15, it had vanished.

Orthotomus sutorius sutorius (Forst.). The Tailor-bird.

One was heard by the Sultanpur Dak Bungalow 4,000' on June 21, 1921.

Cisticola juncidis cursitans (Frankl.). The Fantail Warbler.

Observed in small numbers about the marshy ground below Bajaura, 3,500' in June 1921 and 1922; it must breed in this locality.

Phylloscopus affinis (Tick.). Tickell's Willow-Wren.

M. Babault obtained 2 males and 2 females at Tchari-Djony and Ghary in May and June.

Phylloscopus pulcher kangræ Ticehurst. The Orange-barred Willow-Wren.

This willow-wren breeds fairly commonly in June about 9,000-10,000' in the scrub and forest about the summit of the Jalauri Pass and doubtless on the other similar ranges. I procured fledged young at Rahla 8,800' on July 11, 1923.

Phylloscopus proregulus simlaensis Ticehurst. Pallas' Willow-wren.

A common species in November and December along the valleys up to at least 5,000'.

Phylloscopus maculipennis (Blyth). The Grey-faced Willow-wren.

A single specimen was obtained on November 15, 1923, at 4,500' by the river at Banjar.

Phylloscopus inornatus humii (Brooks). Hume's Willow-wren.

One was seen carrying food in scrub and oak jungle at 10,000' on the summit of the Jalauri Pass on June 17.

Observed in the valley at Sultanpur and Bajaura in November.

Phylloscopus nitidus viridanus (Blyth). The Greenish Willow-wren.

One or two were observed by the river at Manali 6,000' on May 22, 1921, and a specimen was procured. Willow-wrens apparently of this type were seen at 10,000' on the Jalauri ridge on July 16, 1922.

Phylloscopus occipitalis occipitalis (Blyth). The Large-crowned Willow-wren.

This is the common breeding species throughout Kulu at elevations from 5,000 to 8,000'; probably a summer visitor and passage migrant only.

Phylloscopus trochiloides trochiloides (Sundev.). Blyths' Crowned Willow-wren.

Procured by M. Babault at Ghary and Pulga in May.

Seicercus burkii whistleri Ticehurst. The Black-browed Flycatcher-Warbler.

Breeds fairly commonly in the deep scrub at 9,500-10,000' about the summit of Jalauri Pass in June, and doubtless on all the similar ranges of Kulu. Single specimens were seen in the Bhuin marsh 3,600' on December 10, 1922 and below Kraun at 5,000' on November 1, 1923.

Seicercus xanthoschistos albosuperciliaris (Jerd.). The Grey-headed Flycatcher-Warbler.

A very common species in Kulu breeding throughout the valleys from 3,000' apparently up to 8,000', and probably remaining at much the same elevations throughout the winter.

***Horornis pallidus pallidus* (Brooks).** The Pale Bush-warbler.

This bird appears to breed commonly about 6,000'–8,500' in the nullahs above Manali where I have heard it in song in July and on August 1. In November and December I have procured specimens in the bottom of the valley about Katrain 5,000' and Sultanpur 4,000'.

***Suya crinigera crinigera* Hodgs.** The Brown Hill-warbler.

A common and resident species found from 3,000' up to 5,500' and probably higher in all the valleys.

***Oriolus oriolus kundoo* Sykes.** The Golden Oriole.

A common and general summer visitor to the valleys occurring up to 6,000'; most numerous along the Beas Valley from Sultanpur to below Bajaura. Breeds in the latter half of May and in June.

***Oriolus luteolus luteolus* (L).** The Black-headed Oriole.

According to Stoliczka this oriole is 'occasionally seen in the Kulu Valley between 4,000' and 5,000'.

***Temenuchus pagodarum* (Gmel.).** The Brahminy Mynah.

An abundant summer visitor to the Beas Valley from Sultanpur 4,000' to Larji 3,000', and Banjar 5,000' along the Tirthan Valley. One was seen towards Naggar at 4,500' on August 6, 1923. Breeds in June and July.

***Acridotheres tristis tristis* (L.).** The Common Mynah.

An abundant resident throughout the valleys up to about 7,000'. Breeds in May and June.

***Æthiopsar fuscus fuscus* (Wagl.).** The Jungle Mynah.

Observed in June and July only along the Beas Valley from below Bajaura 3,500' to Manali 6,000' but not very common.

***Uroloncha punctulata* (L.).** The Spotted Munia.

A pair were seen at 6,000' above Kraun at the foot of the Bhubhu Pass on July 2, 1921, and a pair by Banjar Rest House 5,000' on November 16, 1923.

***Perrisospiza icteroides* (Vigors).** The Black-and-Yellow Grosbeak.

A male shot at Naggar on January 1, 1904, was sent to the Bombay N. H. Society by General Osborn who stated that it was a winter visitant only (J. B. N. H. S., xv. 716). This however is incorrect. The species is a fairly common resident in Kulu and Saraj occurring at all elevations (I have seen a pair in an oak wood at 3,000' on the bank of the Beas just above Larji on June 10) but most numerous in heavy pine forest from 7,000' to 10,000'.

***Mycerobas carnipes* (Hodgs.).** The White-winged Grosbeak.

On November 9, 1923, I saw 2 or 3 in Silver-fir forest at 9,000' above Pulga.

***Mycerobas melanoxanthus* (Hodgs.).** Spotted-winged Grosbeak.

M. Babault observed this species at Palga and obtained a pair at Ghary on June 6, with the organs in breeding condition.

***Pyrrhula erythrocephala* (Vigors).** The Red-headed Bullfinch.

M. Babault obtained specimens at Ghary and Pulga in May or June. I found a pair on June 16, 1922, at 10,000' on the northern slope of the Jalauri Pass where they were evidently breeding. In winter I observed a couple near Kasol on November 6, and several about 6,500' behind Katrain in the first week of December. According to Stoliczka numbers appear in Kulu in winter, but this must really refer to a concentration in the valleys of the birds which breed in summer along all the higher ranges.

***Loxia curvirostra himalayana* Blyth.** The Himalayan Cross-Bill.

M. Babault states that he has procured a pair at Pulga (May or June) and also seen it at Manali (June).

***Pyrrhospiza punicea humii* Sharpe.** The Red-breasted Rose Finch.

On July 27, 1922, I shot a male at 13,000' in the Humpta Nala where it was feeding on rocky ground amongst patches of snow.

***Carpodacus rhodochroa* (Vigors).** The Pink-browed Rose Finch.

This Rose Finch breeds at high elevations, probably 9,000' to 11,000', on the ranges in Kulu and Saraj. M. Babault saw fledged young somewhere near Ghary as early as May 11.

In winter it moves down the hillsides and may be found as low as 4,500' along the valleys. It is not uncommon.

***Carpodacus erythrinus roseatus* (Hodgs.).** The Common Rose Finch.

M. Babault met with this finch at Bajaura in May, and at Ghary and Tchong. These birds were probably late migrants as I doubt whether the species breeds to the south of the Rhotang Range. It of course breeds commonly in Lahul and Spiti.

***Carduelis caniceps caniceps* (Vigors).** The Himalayan Gold Finch.

A few gold finches were observed at Bajaura 3,600' on June 22-23, 1921, and of these two were birds in juvenile plumage. In November and December flocks were met with in the Alders along the Beas from Katraian to below Bajaura.

***Callacanthus burtoni* (Gould).** The Red-browed Finch.

Two males with the organs developed were obtained by M. Babault at Tchari-Djoni (Tcho-Ti) at the end of May or beginning of June.

***Acanthis flavirostris brevirostris* (Moore).** Eastern Twite.

According to Stoliczka this Twite visits Kulu in winter.

***Serinus pusillus* (Pall.).** The Gold-fronted Finch.

I observed some flocks about the hillsides at 6,000'-6,500' behind Katraian in the first week of December 1922.

***Hypacanthus spinoides* (Vigors).** The Himalayan Green Finch.

M. Babault obtained a male (with the organs undeveloped) at Kothi 8,000' at the foot of the Rhotang Pass on June 28. In summer it appears to be common anywhere in Kulu up to a height of at least 9,000', but I only once met with it in winter, below Jari on November 5. The movements of this finch in the Western Himalayas are difficult to understand and detailed records regarding it are badly needed.

***Passer domesticus indicus* Lard and Selby.** The House Sparrow.

The House Sparrow breeds abundantly in Kulu and Saraj up to a height of 7,000', laying chiefly in May. In winter it probably deserts the higher part of this range but is certainly common along the valley of the Beas at that season.

***Passer rutilans debilis* Hart.** The Cinnamon Sparrow.

The Cinnamon Sparrow is common in May and June in the Alder groves of the Beas from Katraian to Kothi 5,000'-8,000', and at similar elevations in the other valleys; down to 4,000' it is less common. A certain number were observed about 4,000-5,000' in November, but many probably move to the outer ranges in winter.

***Montifringilla nivalis adamsi* Adams.** Adam's Snow Finch.

A specimen from 'Kulu' is in the Gould collection in the British Museum, and Stoliczka states that this bird visits Kulu in winter. In summer it breeds abundantly in Spiti but not in Lahul.

***Montifringilla nemoricola altaica* (Evers.).** Stoliczka's Mountain Finch.

I have observed this Mountain Finch from May to August about 13,000' on the summit of the Rhotang Pass, feeding round the patches of melting snow on the bare hillsides. In winter it moves down into Kulu (S. F. i. 42) and there is a specimen from 'Kulu' in the Gould collection, but I have little definite information on the point. I observed some on the Kharshu Tharch above birch behind Baragraon on December 8, 1922, and on the Tharches about 9,500' at Pulga on November 8, 1923. It probably is abundant at some seasons in winter as large numbers visit the outer ranges about Simla and Dharmasala.

Montifringilla brandti haematopygia Gould. Brandt's Mountain Finch.

Stoliczka states that this finch visits Kulu in winter.

Emberiza fucata arcuata Sharpe. The Grey-headed Bunting.

Observed at Manali 6,000' on May 23, near Kraun 5,500' on July 1, and above Jaggatsukh at 9,000' at the entrance to the Humphta Nala on July 28. It is not uncommon and in the last locality the males were in song sitting on the herbage in some open patches, which were waist deep in balsam, wild parsley, nettles and the like. The song is only a moderate performance.

Emberiza leucocephala (Gmel.). The Pine Bunting.

A common winter visitor to the valley, observed in November and December at 3,500'–5,500' at Katraian, Bhuin, and Bajaura.

Emberiza stewarti Blyth. The White-capped Bunting.

This Bunting breeds from May to July in some numbers in the cultivation of the Beas Valley just below Sultanpur 4,000', and also apparently along the Kraun Road up to 5,000'. The former is probably the locality from which Hume obtained eggs (N. and E. 2nd ed. vol. ii, 167). The song is of the typical bunting type but short and hard. The alarm note is a sharp 'tit' like that of *Horeites brunneifrons*.

I observed a flock on November 30, and December 9, in some thorn bushes on the road between Sultanpur and Raisan.

Emberiza cia stracheyi Moore. The Eastern Meadow Bunting.

Breeds commonly from about 7,000' to 10,000' and in smaller numbers down to 5,000'. Eggs may be found from May to August. In winter it is common from about 6,500' downwards.

Melophus melanicterus (Gmel.). The Crested Bunting.

Occurs in small numbers in June about Banjar 5,000' and Bajaura 3,600', evidently as a breeding summer visitor.

Chelidon urbica Subsp.? The House Martin.

Stoliczka states that a House Martin which he calls *caschmirensis* is common in the Kulu Valley. On August 5, 1923, I saw a large migrant flight of Martins on the road near Naggar. No specimens were obtained but the birds appeared to belong to the typical form and not *caschmirensis*. Probably both occur on migration and one or other form may breed in the highest nalas.

Ptyonoprogne rupestris (Scop.). The Crag Martin.

In 1922 I saw a few about Katraian on December 1–3, and a large flock near Bandrole on December 9. In 1923 I saw a large flock below Jari on November 5.

Hirundo rustica L. The Swallow.

Hume states (N. and E. 2nd ed. vol. ii, 184) that he found the swallow 'breeding at one or two bungalows between Sultanpur in Kulu and Simla. I have no more definite record for Kulu.'

Hirundo smithii Leach. The Wire-tailed Swallow

M. Babault obtained a male at Sultanpur. One or two pairs breed near Bajaura where I saw fledged young on June 29. A summer visitor only.

Hirunda daurica nepalensis Hodgs. Hodgson's Striated Swallow.

This form of Striated Swallow is a common summer visitor to the valleys from 3,000' up to about 5,000', building its nest under rocks on steep hill sides. Eggs are laid in June. Observed from May to August.

Motacilla alba dukhunensis Sykes. Eastern White Wagtail.

A few were observed at Sultanpur 4,000' in November and December.

Motacilla alba personata Gould. The Masked Wagtail.

I found a nest with 5 young in quill and 2 addled eggs on a Shingle bank of the Beas near Raisan 4,500' on May 21, 1921. A full account of this nest has appeared in the Journal, B. N. H. S., xxix. 282.

A few were observed near Sultanpur 4,000' and Bajaura 3,600' in November 1922 and 1923.

Motacilla alba alboides Hodgs. Hodgson's Wagtail.

M. Babault obtained a female at Bajaura in May or June. I saw several on the Beas at Bajaura 3,600' on November 12, 1923.

Motacilla maderaspatensis Gmel. The Large Pied Wagtail.

A resident in small numbers along the Beas from Larji 3,000' to Katraian 5,000'.

Motacilla cinerea melanope Pall. The Grey Wagtail.

The Grey Wagtail breeds commonly in May and June along the rivers and larger streams of Kulu from 4,000' upwards to at least 10,000'.

The majority leave the Sub-division in winter but in November and December I have observed a few about up to 5,000' in the Beas Valley.

Motacilla citreola citreola Pall. The Yellow-headed Wagtail.

Two were seen by the Beas above Manali 6,000' on May 25, 1921.

Anthus hodgsoni Richmond. The Indian Tree Pipit.

Hume long ago described a single egg of this Pipit taken in Upper Kulu on the snowy range bounding Spiti, about 11,000' (N. and E. 2nd ed., ii. 209).

M. Babault obtained females at Pulga on May 19, and Tchari-Tjony on June 1, the latter was by the snow line with the organs in breeding condition.

It probably breeds about the summits of all the higher ranges of Kulu as on June 16, 1922. I found several nests with eggs in all stages of incubation on the Jalouri ridge at 9,500-10,500'. These have been described elsewhere.

In November parties appear in the valleys about 4,000-5,000'.

Anthus leucophrys jerdoni (Finsch). The Brown Rock Pipit.

Common in June and July along the valley roads from Bajaura to Larji and Banjar.

Anthus richardi rufulus Vieill. The Indian Pipit.

This plains' Pipit breeds in small numbers in June about the marshy ground of the Beas Valley below Bajaura at 3,500'. It is doubtless a summer visitor only.

Anthus roseatus Hodgs. Hodgson's Rosy Pipit.

In May and June this lovely Pipit breeds commonly about 12,000'-13,000' along the alpine pastures of the Rhotang Range, and M. Babault found it on similar ground about forest limit above Pulga; it doubtless breeds along the whole of the highest ranges in northern Kulu. It is one of those birds that are particularly partial to the sodden ground on the edge of melting snow. In winter it is common along the Beas Valley about Sultanpur and Bajaura and there as elsewhere in its winter range is largely a marsh pipit.

Anthus spinoletta blakistoni Swinh. The Altai Water Pipit.

I obtained a specimen from a party in a patch of marshy ground by the Tirthan River below Manglaur 4,000' on November 17, 1923.

Oreocorys sylvanus (Hodgs.). The Upland Pipit.

Common, at least from May to July, from 3,000' to 5,500' in the valley of the Beas and along the Tirthan to Banjar. Hume personally took eggs of this bird in Kulu (N. and E. 2nd ed., ii. 21).

Otocorys alpestris longirostris Moore. The Long-billed Horned Lark.

Kulu is the type locality for this form of Horned Lark and there are 6 specimens from 'Kulu' (without more definite data) in the Gould collection in the British Museum. It is found in summer breeding at 13,000' on the summit of the Rhotang Pass and as M. Babault procured it also at the end of May at Tchari-Djony one may assume that it breeds along the whole of the Rhotang chain.

Otocorys alpestris elwesi Blanf. Elwes' Horned Lark.

A female labelled 'Kulu' and collected by Hay is in the Gould collection in the British Museum.

***Alda arvensis guttata* Brooks.** The Kashmir Skylark.

Flocks of this Lark (which breeds in Lahul but not so far as I know in Kulu unless Hume's eggs belonged to this form) are to be found in November and December in the fields of the Beas Valley from Bajaura 3,600' to Katraian 5,000'. Stoliczka speaks of a Lark which he calls *Alaudula raytal* as a common, winter visitor to Kulu and often caged by the people, but I am not sure what species is indicated.

***Zosterops palpebrosa elwesi* Stuart Baker.** The White-eye.

The White-eye breeds but is not very common in Kulu and Saraj up to a height of about 5,000' and I have observed it from May until August. It is probably a summer visitor only.

***Cyrtostomus asiaticus* (Lath.).** The Purple Sun-bird.

In June 1922 I saw a few of these Sun-birds in the valley of the Beas between Bajaura and Larji about 3,000-3,500'.

***Dicaeum ignipectum* (Hodgs.).** The Fire-breasted Flower Pecker.

Stoliczka states that this beautiful little bird is 'pretty common in Kulu'. I saw a male at some Mistletoe in the garden of Katraian Rest House on December 1 and 2, 1922, and believe that I saw another on November 13, 1923, in the Ban oak forest by the Beas a mile or two above the Outbridge.

***Picus squamatus squamatus* Vigors.** The Scaly-breasted Green Woodpecker.

A common resident in the Kulu and Saraj occurring at least from the bottom of the valleys up to 7,000' in winter and probably higher in summer. At Manikaran I saw one of these Woodpeckers perch on the side face of a great boulder, but near enough to the top to catch the edge with two front claws.

***Picus canus occipitalis* Vigors.** The Black-naped Green Woodpecker.

On June 26, 1922 I saw three of these woodpeckers together on the bank of the Beas below Manali 6,000'.

***Dryobates hyperythrus* (Vigs.).** The Rufous-bellied Pied Woodpecker.

I saw an adult male on May 17, 1921, at 8,000' on the northern face of the Bhubhu Pass.

***Dryobates himalayensis* (Jard. and Selby).** The Himalayan Pied Woodpecker.

A common and resident species, but not observed below 6,500'.

***Dryobates auriceps* (Vigors).** The Brown-fronted Pied Woodpecker.

A common and resident species throughout the valleys from Larji 3,000' up to about 6,000' and possibly higher.

***Liopicus mahrattensis* (Lath.).** The Yellow-fronted Pied Woodpecker.

M. Babault obtained a female at Ghary in May although this is not a bird which might have been expected to occur in Kulu.

***Picumaus innominatus innominatus* Burton.** The Speckled Piculet.

A pair bred probably annually in the rest house compound at Sultanpur 4,000', and I saw a single bird below Jarj at 5,000' on November 5, 1923.

***Megalaema virens marshallorum* Swinhoe.** The Great Himalayan Barbet.

In summer this fine Barbet is common in the great forests of Kulu from about 6,000' upwards; in November and December I have observed it also at 6,000' but it doubtless occupies a lower zone during the winter months.

***Coracias garrula semenowi* Loud. and Tschusi.** The Kashmir Roller.

On May 21, 1921, I saw a Roller by the road side just below Raisan 4,800' on the Beas.

***Merops orientalis* Lath.** The Little Green Bee-eater.

Stoliczka states that this Bee-eater is not uncommon in Southern Kulu from March till the end of October. I have only met with it once, when two were seen in cultivation below Bhuin 4,000' on November 4, 1923.

***Ceryle lugubris guttulata* Steg.** The Himalayan Pied Kingfisher.

This fine Kingfisher is generally distributed and resident throughout Kulu and Saraj from 3,000' at Larji up to about 6,000' on all the rivers and streams. Its large size and handsome black and white plumage and the loud call note 'ping' (similar to one of the calls of the Red-wattled Lapwing) readily attract attention as the bird flies along above the surface of the troubled waters or perches on the trees and boulders that afford handy fishing stations. It occasionally plunges into the water from the wing like the smaller Pied Kingfisher of the plains.

***Alcedo atthis* (Hasselg.)** The Common Kingfisher.

I have observed a few of these Kingfishers along the Beas and Tirthan Rivers up to 6,000' in May and June; they doubtless breed here as M. Babault procured a specimen on May 3, with the organs greatly developed.

***Halcyon smyrnensis smyrnensis* (L.)** The White-breasted Kingfisher.

A resident in small numbers along the valley of the Beas nearly up to Manali 6,000'.

***Upupa epops epops* L.** The Hoopoe.

In summer the Hoopoe breeds commonly along the valleys from Larji up to a height of about 8,000', wherever there is open ground in the vicinity of houses and cultivation. I took c/4 fresh eggs at Sultanpur on May 20 and found several nests with young in June and July, all under the roofs of houses. These birds are summer visitors only, and though an occasional Hoopoe may be seen about the Beas Valley in winter, such birds are probably winter visitors from further north.

***Micropus melba* (L.).** The Alpine Swift.

I have observed this Swift feeding in the Beas Valley about Sultanpur and Katraian in May and June, so it perhaps breeds in some of the ranges.

***Micropus apus pekinensis* (Swinh.).** The Eastern Swift.

On May 21-22, 1921, these Swifts were common along the Beas Valley about Raisan and Katraian. They breed commonly in Lahul.

***Micropus pacificus leuconyx* (Blyth).** Blyth's White-rumped Swift.

On May 21, 1921, I saw these Swifts in company with *M. a. pekinensis* in the Beas Valley about Katraian. Two were seen at Banjar on June 25, 1921. They doubtless breed in the precipices throughout Kulu.

***Micropus affinis* (Gray).** The Common Indian House Swift.

On July 1, 1921, I saw a flight about 5,000' on the road below Kraon.

***Hirundinapus caudacuta nudipes* (Hodgs.).** The White-necked Spinetail.

On June 16, 1922 with great difficulty I secured a specimen of this Spinetail from a flock which were flying about the crest of the Jalouri Ridge 11,000' to the east of the Pass. The flight was exceedingly swift and their wings made a very musical sound in the air different to the usual tearing sound made by the wings of fast flying birds.

***Collocalia fusciphaga brevirostris* (McClell.).** The Himalayan Swiftlet.

Flocks of this swiftlet may be seen feeding about the valleys and lesser ranges anywhere in Kulu and Saraj though I have not observed them over 6,000'. My records refer to May, June, July, August, November and December so the species may well be resident.

***Caprimulgus* sp?** Nightjar.

I have heard Nightjars in the Nagni Nala May 18 and below Shoja on June 15; the call was a rapid chuck chuck chuck with the hollow sound of a stone on ice, but I hesitate to identify the species from this alone.

***Cuculus canorus* L.** The Cuckoo.

The Cuckoo is a common summer visitor to Kulu and Saraj occurring from 3,000' at Larji up to 8,000' and probably higher. A specimen obtained clearly belongs to the race *C. c. telephonus* Heine but the majority appear to belong to the typical form.

Cuculus poliocephalus poliocephalus Lath. The Small Cuckoo.

I have met with this little Cuckoo in the Beas Valley on two occasions, namely a pair in an alder grove above Manali 6,200' on May 25, 1921 and below Sultanpur 4,000' on July 7, 1923. M. Babault however obtained 4 specimens in May and June at Pulga Manikaran and Tchhari-Djony.

Cacomantis merulinus passerinus (Vahl.) The Indian Plaintive Cuckoo.

On June 24, 1922, I heard this Cuckoo calling for a long time near Sultanpur Rest House 4,000', and believe that I also heard it at Banjar 5,000' on June 12.

Clamator jacobinus (Bodd.) The Pied-crested Cuckoo.

M. Babault obtained a male on 28th June with the organs in breeding condition at the upper edge of the forests at 12,500' at Rahla by the Rhotang Pass. In June and July this Cuckoo is not uncommon in the lower Beas Valley round about Bajaura and I have also heard it at Banjar 5,500'.

Eudynamis scolopaceus scolopaceus (L.). The Koel.

On June, 22, 1921, I heard a Koel calling in the swamp between Bajaura and Bhuin at 3,800'.

Psittacula eupatria nepalensis (Hodgs.). The Large Indian Paroquet.

M. Babault records that he observed this species in the valley of Kulu in April and May and also obtained a juvenile on the Dulchi Pass.

Psittacula cyanocephala cyanocephala (L.). The Blossom-headed Paroquet.

According to Stoliczka this bird 'may occasionally be seen in the southern portions of the Kulu Valley'.

Psittacula schisticeps schisticeps (Hodgs.). The Slaty-headed Paroquet.

This is the common Paroquet of Kulu and Saraj where it may be found in summer in every valley and on the ridges up to 8,000' and perhaps higher. It is apparently a resident species.

Strix aluco nivicola Blyth. The Himalayan Wood-owl.

This owl breeds fairly commonly in Kulu from 6,000' probably up to about 10,000' and is doubtless a resident species.

Bubo bubo turcomanus (Eversm.). The Great Eagle Owl.

An Eagle Owl was obtained about 12,000' in Kulu in the summer of 1872 while it was eating a snow partridge on the ground. This bird was recorded by Hume in S. F. I. 315 and named *B. hemachalana* but that is merely a synonym of Eversman's earlier name.

Bubo bubo bengalensis Frankl. The Rock Horned Owl.

Large Owls seen by me in cliffs at Katraian Rest House 5,000' on June 19, 1921, and on the Kraun Road near Sultanpur 4,000' on August 7, 1923 were I believe of this species.

Scops plumipes. Scops Owl.

Under this name M. Babault records a male Scops Owl in breeding condition that he obtained at Pulga on May 18.

Glaucidium cuculoides cuculoides (Vigors). The Large Barred Owlet.

This very diurnal species of owl is a common resident in Kulu occurring from the bottom of the valleys to at least 7,500'. On one occasion when marching along the valley road below Manali I came upon an Owlet and a crow fighting desperately in broad daylight on the road for the possession of a disreputable tattered object, which on my nearer approach proved to be a half-grown nestling owlet.

Pandion haliaetus (L.). The Osprey.

In Mr. Donald's list with the remark 'Not common; probably a migrant; may breed'. I doubt the last clause.

Ægyplus monachus (L.). The Cinereous Vulture.

Included in Mr. Donald's list of birds of Prey in the Gazetteer with the remark 'Probably breeds in Kulu; not very common'. In my opinion it is in the

highest degree unlikely that it has ever bred in Kulu which is too far east of its known breeding range.

Torgos calvus (Scop.). The King Vulture.

In Mr. Donald's list with the remark 'Chiefly found among the Lower hills; nowhere very common'. I saw one on the marsh some 2 miles below Bajaura on June 10, 1922.

Gyps fulvus (Gmel.). The Griffon Vulture.

In Mr. Donald's list with the remark 'Seen near Larji'.

Gyps himalayensis Hume. The Himalayan Vulture.

Fairly common and generally distributed throughout Kulu and Saraj where it breeds in suitable localities.

Gyps tenuirostris Hume. The Himalayan Long-billed Vulture.

In Mr. Donald's list with the remark 'Lower hills, seen near Zakat Khana'.

Pseudogyps bengalensis (Gmel.). The White-backed Vulture.

I saw 3 of these vultures together by the roadside, some 2 miles below Bajaura on November 13, 1923.

Neophron percnopterus (L.) The Egyptian Vulture.

I have observed a few Egyptian Vultures in June at Katraian, Bajaura, Larji and Banjar but they are probably summer visitors only for I saw none in November or December. M. Babault collected a male at Bajaura and a female on the Dulchi Pass and states that they breed in June thereabouts, nesting on cliffs, though in his specimens the organs were not developed. These specimens he identifies as belonging to the race *ginginianus* with the yellow beak, though I should have expected the typical race in Kulu.

Gypaetus barbatus grandis Storr. The Lammergeaier.

Stoliczka has recorded the following note (Jour. A.S.B. xxxvii. 1868) :—'The natives of Kulu, about Plash and the Eastern districts prize the meat very highly, which is not only eaten by the low class, the Kolis, but rather more by the higher class, the Kanaitis. They generally tie a Chukor on a short string, and stick 4 or 5 sharpened spears in the ground crossing each other, so as partially at least to cover the bird, and at the same time to radiate with their points in different directions. The Eagle is watched from some distance and as soon as it throws itself with its usual great force and velocity upon the prey, it is overpowered with large clubs before it can extricate the spears from its body.'

The Lammergeaier is common and resident throughout Kulu and Saraj, and the sight of one or more of these grand birds sweeping round the contours of the hills is a feature of every march. I once saw 4 adults together but it is unusual to see more than a single pair at any time. On one occasion above Manikaran I saw a bird in the dark immature plumage perched on a side bough of a lofty dead tree at the edge of a ravine, but otherwise I have only seen it settle on the ground or on crags and cliffs.

Aquila chrysaetus daphanæ Hodgs. The Golden Eagle.

Mr. Donald's list contains the following statement :—'Resident; lives on crows, foxes, pine martins. Takes Monal and Snow-cock occasionally, but crows are his chief food. Therefore a valuable game preserver.'

The Golden Eagle probably breeds amongst the crags of all the higher ranges in Kulu, but the nest has never been taken in Kulu by any European. I have myself observed the species on the wing by the Bhubhu Pass (9,000') on October 31, at over 10,000' at Pulga on November 9, and as below as 6,000' just below Manikaran on November 10. M. Babault observed it at Tchari-Djoni (end of May) and on the Rhotang Pass on June 29.

Aquila heliaca Savigny. The Imperial Eagle.

In Mr. Donald's list with the remark 'Winter migrant; lives on rats, lizards, carrion.' M. Babault killed a specimen on June 29 on the Rhotang Pass.

Aquila nipalensis nipalensis (Hodg.). The Steppe Eagle.

In Mr. Donald's list with the same remarks as under the last species.

Aquila rapax vindhiana (Frankl.). The Indian Tawny Eagle.

In Mr. Donald's list with the remark 'Not common. Seen in lower valleys. May breed there.'

Aquila clanga Pall. The Large Spotted Eagle.

In Mr. Donald's list with the remark 'Seen near Katraian. Food chiefly frogs and lizards.'

Hieraetus fasciatus (Vieill.). Bonelli's Eagle.

In early November 1923 when in company with Mr. Donald I saw a pair of Bonelli's Eagle frequenting a small cliff two or three miles above Sultanpur on the Kraon road at about 4,500-5,000'. They evidently had an eyrie in this cliff or were intending to nest there. One bird was in the beautiful adult plumage, the other was in a curious phase, pale below uniform brownish above, which at first led us to mistake it for some other species.

Hieraetus pennatus (Gmel.). The Booted Eagle.

'Fairly common probably breeds in the valley. Lives on rats, etc., sometimes takes Chukor. Not very destructive to game.' (Donald).

Ictinaetus malayensis perniger (Hodgs.). The Black Eagle.

Mr. Donald remarks of this species:—'Very rare if known at all in the valley. Mr. Howell's inclusion of this species doubtful.' But in N. and E. 2nd ed., vol. iii. 145, Hume records obtaining a single egg from Kulu with the parent bird taken on January 7.

Spizaetus nipalensis nipalensis (Hodgs.). Hodgson's Hawk-Eagle.

In Mr. Donald's list with the remark 'Resident and probably breeds near Naggar. Very destructive to game.' On December 8, 1922, I saw what was apparently a pair of these Hawk-Eagles chasing a Falcon with booty at a great height across the valley above Katraian.

Circus gallicus (Gmel.). The Short-toed Eagle.

In Mr. Donald's list with the remark 'Seen in Saraj near Sutlej river. Food rats, lizards, snakes, etc.'

Butastur teesa (Frankl.). The White-eyed Buzzard-Eagle.

In Mr. Donald's list with the remark 'Seen near Bajaura. Probably a wanderer but may be resident. Food, rats, mice, worms, lizards.' In 1921 I saw one above Bajaura on June 22, and 2 or 3 about the marshy ground some 2 miles below Bajaura later on June 23 and 29. M. Babault obtained the young in May or June from a nest at Bajaura.

Haliaetus albicilla (L.). The White-tailed Sea Eagle.

Down as 'a migrant' in Mr. Donald's list. He records (J.B.N.H.S., xxvi. 1071) a supposed example that he saw on the Beas in April 1916.

Polioaetus humilis (Jerd.). The Himalayan Fishing Eagle.

In Mr. Donald's list with the remark 'Resident and probably breeds. Destructive to fish.' He states (J.B.N.H.S., xxvi. 1019) that it is fairly common all over the Kulu Valley. I have seen an occasional eagle in June and July along the Beas from Sultanpur to Manali which were probably of this species but no specimens were procured.

Haliastur indus indus (Bodd.). The Brahminy Kite.

'Not common in the lower valleys near rice fields' (Donald).

I saw a pair opposite to Katraian over the rice fields across the river on July 29, 1922. It is doubtless only a rains visitor as in the Kangra Valley.

Milvus migrans govinda Sykes. The Common Pariah Kite.

The Pariah Kite is not uncommon and probably resident along the Beas Valley from Manali to Larji and up the Tirthan Valley to Banjar, up to a height of about 6,000'.

Milvus lineatus (Gray). The Large Kite.

Hume states in Nests and Eggs 2nd ed., vol. iii. p.176, that he has obtained the eggs of the Larger Kite from Kulu. This species is often difficult to

distinguish in the field from the Common Pariah Kite but I possess three specimens collected in Kulu a breeding male from Kraun 5,000' July 1, 1921, a nestling from a nest at Bajaura 3,600' June 8, 1922, and an immature male from Katraian 5,000' December 9, 1922, and believe that I have seen others occasionally along the valley from Naggar to Bajaura in the summer months. A Kite seen at 12,000 feet on the Rhotang Pass on May 26, 1921, was most probably of this species.

Elanus caeruleus caeruleus (Desf.) The Black-winged Kite.

In Mr. Donald's list with the remark 'A migrant. Lives on insects.'

Circus cyaneus (L.). The Hen-Harrier.

'A migrant; fairly common in autumn and spring' (Donald).

M. Babault obtained a female in May at Bajaura with the organs developed but he was I think wrong in drawing the conclusion that this bird was on its breeding ground.

(I saw some species of Harrier in the ring-tail plumage at Pulga 7,000' on November 8, 1923, quartering the fields in front of the Rest House.)

Circus æruginosus (L.). The Marsh-Harrier.

'A migrant; fairly common in autumn and spring' (Donald).

M. Babault obtained 2 males at Bajaura on June 13, with the organs developed.

Buteo buteo japonicus (T. & S.). The Himalayan Buzzard.

In November and December I have met what I believe to be the Upland Buzzard occasionally along the bottom of the Parbatti and Beas Valleys but it is not found there during the summer months, and is then probably breeding in the higher ranges. The whole question of the forms of Buzzard that occur in Kulu and their status requires careful examination.

Buteo ferox (S.G. Gmel.). The Long-legged Buzzard.

Buzzards, probably both of this species and the last, are common in the valleys in winter, and some of them appear to breed on the ranges. Hume states (N. & E. 2nd ed., iii. 125) that typical *ferox* breeds in Kulu and that he possessed an egg with the skin of the parent obtained on January 10.

The race *B. f. hemilasius* Temm. & Sch., also occurs in Kulu as a skin obtained by Hume and made the type of his *Archibuteo leucoptera* (S.F.I. 318) is in the British Museum.

Astur gentilis (Linneus). The Goshawk.

There is very little definite information on record about the Goshawk which probably breeds in the higher forests of Kulu. Certain it is that several are netted annually on the ranges of the Parbatti Valley and Saraj and taken down into India for sale to Falconers.

On December 10, 1922, at Bhuin I met a man with an adult male and two immature females tied to a pole, hooded and with the tails sewn up in cloth according to custom, and he stated that they had been caught in Saraj a week or two earlier.

A bird seen by me in the distance about 10,000 ft. above Pulga, on November 9, 1923 was perhaps an adult Goshawk.

Astur badius dussumieri (Temm.). The Shikra.

'Common: resident and breeds' (Donald).

I shot a fine male in an alder grove (4,000 ft.) just above Sultanpur on May 21, 1921.

Accipiter nisus (L.). The Sparrow-hawk.

'Common; resident and breeds' (Donald).

I have seen an occasional *Accipiter* in Kulu but usually under circumstances that did not allow me to be sure of the exact species: the majority were probably this bird. M. Babault obtained a pair at Tchari-Djony at the end of May and states that he found nests in the forests in this locality.

Accipiter virgatus affinis (J. E. Gray). The Besra Hawk.

'Probably resident and breeds. Nowhere common' (Donald).

Stoliczka on the other hand (J. A. S. B. xxxvii) states that the Besra is far more common than the Sparrow-hawk in the Kulu Valley. I obtained a female at Manali (6,000') on August 4, 1923.

Falco peregrinus L. The Peregrine Falcon.

'A winter migrant' is the remark in Mr. Donald's list.

Falco peregrinus peregrinator Sund. The Shahin.

'Resident and probably breeds. Destructive to small game such as partridges, chukor, etc.' (Donald). I have seen single Falcons above Manali (6,200') on June 17, 1921 and above Katraian (6,000') on December 8, 1922, which were probably of this species. I also have seen an undoubted pair of Shahins at Larji (3,000') on November 18, 1923, when they were bullying an Eagle on the hillside in front of the Rest House.

Hume records (N. and E. 2nd ed., iii. 186) that he possessed an egg taken near Deet in Kulu on February 6, and two eggs taken near Nitta in Kulu on February 3.

Falco severus indicus (Meyer & Wiglls.). The Indian Hobby.

Donald states that it is 'Resident and probably breeds', while Stoliczka says (J. A. S. B. xxxvii) that is not common in Kulu. These statements appear to me to require verification, as I am of opinion that this Hobby has not such a wide range as is usually attributed to it.

Falco subbuteo subbuteo (L.). The Hobby.

I shot a fine adult male Hobby on July 28, 1922, at 9,000' at the mouth of the Humpta Nala, and another adult male on the summit of the Rhotang Pass 13,000' on July 12, 1923. A Falcon seen at 7,000' in the pine forest above Manali on the Kothi Road on August 4, 1923, was almost certainly also of this species. M. Babault obtained a pair in breeding condition at Pulga on May 19.

Falco tinnunculus interstinctus Mc Cl. The Kestrel.

During the summer the Kestrel is exceedingly common throughout Kulu from the bottom of the valley up to heights of 10,000 and 12,000'. A pair seem to breed in every rocky cliff of importance and many pairs also breed in the conglomerate cliffs of earth and boulders which represent the remains of the old bed of the valley. During the winter the species is distinctly less common and at this season many birds doubtless leave Kulu and winter in the plains. These little Falcons are very quarrelsome and the approach of any larger bird of pair to their eyries is the signal for a great display of temper, the intruder being mobbed with an incessant petulant screaming note.

M. Babault found 4 feathered young in a nest at Rahla on June 24.

The typical race may be expected as a winter visitor and passage migrant through Kulu.

Sphenocercus sphenurus (Vigors). The Kokla Green Pigeon.

The Kokla Green Pigeon is a very common summer visitor to Kulu and Saraj occurring in all the valleys up to at least 6,000' and doubtless higher. Stoliczka notes that it is very fond of Mulberry fruit and it is distinctly partial to elms and alder trees.

Columba livia subsp? The Blue Rock-Pigeon.

The Blue Rock Pigeon breeds very commonly in the rocky gorges and precipices throughout Kulu and Saraj up to a height of at least 10,000'. It is a resident species but some of the flocks which feed in cultivation in the valleys during the winter months may be immigrants.

Columba rupestris Pallas. The Hill Blue Rock-Pigeon.

M. Babault met with this species at Tchari-Djony. It is of course common in Spiti.

Columba leuconota (Vigors). The Snow Pigeon.

M. Babault met with the Snow Pigeon at Tchari-Djony. In May and June I have met with a few on the southern slopes of the Rhotang Pass, usually over 10,000' though I have seen it as low as 8,500' below Rahla. There is a specimen from Kulu dated February in the Gould collection.

***Streptopelia orientalis meena* (Sykes).** The Indian Rufous Turtle Dove.

The Turtle Dove is a common summer visitor to Kulu and Saraj occurring at all heights during the passage but probably not breeding below 6,000'.

***Streptopelia chinenses suratensis* (Gmel.).** The Spotted Dove.

The Spotted Dove breeds commonly in the valleys of Kulu and Saraj up to about 6,000' at Juggatsukh. On June 20, I found a nest placed on the rafters of the rest house verandah at Bajaura. It is probably a summer visitor only.

***Streptopelia senegalensis cambayensis* (Gmel.).** The Little Brown Dove.

M. Babault obtained two females at Ghary on May 8, and a male at Bajaura.

***Streptopelia decaocto decaocto* (Friv.).** The Indian Ring Dove.

A common Summer visitor to the valley of the Beas from Larji 3,000' to Manali 6,000', but less numerous above Sultanpur.

***Catreus wallichii* (Hardw.).** The Cheer Pheasant.

Hume records taking a nest with 4 eggs early in May a few miles from Jaggat sukh (N. and E. 2nd ed., vol. iii, 413). My own collection contains 5 eggs which were brought in fresh to me from the neighbourhood of Katraian in May. The eggs of this pheasant however continue to be sufficiently rare in collections.

The Cheer Pheasant is apparently found throughout Kulu and Saraj at moderate elevations above 5,000' but as elsewhere in its range it is sparsely distributed and very local. According to Tyacke (Manual Rev. ed. 1907) the best locality for it in Kulu from the sportsman's point of view is the Dobi Nala. M. Babault collected specimens on the Dulchi Pass. Personally I only met with it in Kulu in the Parbatti Valley at Kasol on open dry grassy slopes amongst scattered cheel trees. It has no seasonal altitudinal movement.

***Pucrasia macrolopha* (Less.).** The Koklas Pheasant.

The Koklas Pheasant is common in Kulu and Saraj in the middle zone of forests from about 7,000 to 9,500'.

***Gennæus hamiltoni* (Griff.).** The White-crested Kalij Pheasant.

This is the common pheasant of Kulu and Saraj and it is very generally distributed at all heights from 4,500' up to 8,000', being especially partial to heavy undergrowth in the neighbourhood of water.

***Lophophorus impejanus* (Lath.).** The Monal.

The numbers of the Monal Pheasant are periodically reduced by netting and snaring in those areas where the slackness or venality of the local subordinate officials encourages the poaching instincts of the Kulu villager; for this pheasant besides affording a welcome dish is considered peculiarly appropriate to form a 'Dali'; while the plumage of the male—and more especially the long crest feathers, has a certain monetary value. When poaching is reduced to the minimum the Monal is common in all the forests of the higher ranges from 8,000 to 12,000'; with the descent of the snowline in winter the birds collect in the lower portion of their range and then excellent sport may be obtained by the sportsman who is sound enough in wind and limb to undergo the necessary climbing. I know no finer sport; unrivalled air and scenery, the magnificence of the quarry, the difficulty of the climbing and shooting and the immense satisfaction as each bird is gathered in to the bag, all combine to afford a day's sport which is seldom equalled and never surpassed in the annals of the shot gun. But the bag is invariably small. Personally I have never killed more than six Monal in one day though other pheasants may also be obtained to swell the bag.

***Tragopan melanocephalus* (Gray).** The Western Tragopan.

The Tragopan is the rarest of the pheasants of Kulu but it may be found in small numbers here and there on the high ranges at the same elevations as the Monal and also slightly lower in winter. It is more of a skulker and is usually found in thick undergrowth, Rhue or light Ringal Bamboo growth. The eggs are very rare in collections but I obtained 4 rather incubated eggs from Saraj in May 1928 through a village headman.

***Coturnix coturnix coturnix* (L.).** The Quail :

Stoliczka states that the Quail may occasionally be obtained in the Kulu Valley between 4,000' and 5,000' ; Kulu is however outside the main migration route of the species.

***Arboricola torqueola millardi* SB.** The Simla Hill Partridge.

I have heard the call of this partridge not infrequently about the summit of the Dulchi Pass and Tyacke states (Manual Rev. ed. 43) that this is the only spot in the Kulu Valley where it is to be found.

***Alectoris graeca chukar* (Gray).** The Chukor.

This splendid game bird may be found throughout Kulu and Saraj wherever the absence of forests provides the necessary open ground, though the stonier and more barren the slopes the greater the likelihood of obtaining sport. At one time good bags were procured, the best that I have heard of being over 40 brace, but of late their numbers have been reduced by poaching. The Chukor is a favourite cage bird in Kulu and numbers are taken thence into the outer hills.

***Francolinus francolinus asiæ* Bonap.** The Black Partridge.

The Black Partridge is found in small numbers in the valley of the Beas from Raisan to Larji, in the Kraun Valley, and up the Tirthan to Banjar at heights up to 6,000'. It is however fairly common only in the open cultivated valley about Bhuin and Bajaura.

***Tetraogallus himalayensis* (Gray).** The Snow Cock.

There are several specimens from Kulu in the Hume Collection but the only definite locality specified is the Jalouri Range (January-February). Stoliczka states that it is found in Northern Kulu. I have no personal knowledge of the bird in Kulu or Saraj but M. Babault met with it at Tchari-Djony. It is probably common enough on some of the highest ranges.

***Lerwa lerwa* (Hodgs.).** The Snow Partridge.

Stoliczka states that the Snow Partridge 'is numerous in the north-western parts of Kulu during the winter, when it descends from the snowy ranges somewhat lower down'. In the Hume collection there are 8 specimens collected in January and February on the Jalouri range, and two from Nunda in February. M. Babault met with it at Tchari-Djony at the end of May. I received a specimen in the flesh which had been obtained somewhere in Saraj and Mr. Shuttleworth, I.C.S. (once Sub-Divisional Officer of Kulu) informed me that he shot a single bird from a covey in February, 1919 on the hump of the Rhotang Pass above Rahla at about 9,500-10,000'. More definite information about this bird is greatly to be desired, as its numbers appear to be diminishing throughout the Western Himalayas.

***Amaurornis fuscus* (L.).** The Ruddy Crake.

M. Babault found it in small parties in the Bajaura marsh on June 13 and secured a male. I obtained a male (with the organs partly developed) on June 30, 1921, at Sultanpur 4,000' in the alder grove between the town and the Tehsil.

***Lobivanellus indicus indicus* (Bodd.).** The Red-wattled Lapwing.

The Red-wattled Lapwing is fairly common in summer along the valley of the Beas from Manali 6,000' to Larji 3,000'. I fancy the majority of these birds are only summer visitors as with the exception of a party of 4 seen at Bajaura on November 12, 1923. I have seen none in winter.

***Charadrius dubius jerdoni* (Legge).** The Little Ringed Plover.

On June 8 and 9, 1922, I found a couple of pairs of Ring Plover which were clearly nesting on the shingle banks of the Beas at Bajaura 3,600'. M. Babault obtained a pair in June in the same locality and also notes that the species was in parties, probably I suppose family parties.

Ibidoryhynchus struthersii Vigors. The Ibis-Bill.

One or two pairs of this beautiful and interesting bird are resident in the valley of the Beas and breed on the shingle beds of the river but I think it desirable to suppress the exact locality.

Tringa hypoleuca (L.). The Common Sandpiper.

The Common Sandpiper may be found here and there on the rivers of Kulu any time in the year, though but rarely in summer. I have no evidence that it ever breeds in the country, and it is hardly likely to do so as in Lahul I did not find it breeding anywhere nearer than Jispar.

Tringa ochropus (L.). The Green Sandpiper.

Not uncommon as a winter visitor and non-breeding individuals are also to be met with in summer, along the valley of the Beas.

Tringa totanus (L.). The Redshank.

On June 8, 1922, I saw a Redshank on the shingle of the R. Beas at Bajaura 3,600'.

Scolopax rusticola (L.). The Woodcock.

The Woodcock must breed fairly commonly in all the larger valleys of Kulu about 8,000' to 10,000'. In winter the birds descend to lower levels and after snow has fallen on the ranges they are found in the 'mahals', the alder groves mixed with swamp and bush undergrowth, along the Beas and presumably the Parbatti and other valleys. Unfortunately there is practically nothing on record about the woodcock in this, one of its most noted habitats in India, and details are most desirable as regards the elevations and nullahs in which the bird has been met in the breeding season, its seasonal altitudinal movements, and the sport that it has afforded in winter.

Capella nemoricola (Hodgs.). The Wood-Snipe.

M. Babault obtained two males with the organs developed at Tcho-ti on the River Tcho at the end of May so it doubtless breeds in the vicinity. Details about the bird are sadly wanting.

Capella solitaria (Hodgs.). The Solitary Snipe.

On November 9, 1923, I flushed a solitary snipe from a small half-frozen stream at 9,000', above Pulga; this was at the edge of a small open 'Tharch' in Silver Fir forest. A few days later on November 17, I shot one from the boulder strewn nala by the Manglaur bridge 4,000'.

Capella stenura (Kuhl.). The Pintail Snipe.

There is a specimen from Kulu collected by Major Hay in the Gould collection in the British Museum.

Capella gallinago (L.). The Fantail Snipe.

Stoliczka says that the snipe has been obtained in Southern Kulu in winter.

Phalacrocorax carbo (L.). The Common Cormorant.

A cormorant, presumably of this species, occurs in Kulu though it apparently does not breed in the country. I have seen a few on the Beas from Bhuin to Larji in June and November, and one in November on the Tirthan below Manglaur.

Anser anser (L.). The Grey Lag Goose.

Occurs in the list of Game birds given in the Gazetteer.

Anser indicus (Lath.). The Bar-headed Goose.

Occurs in the list of Game birds given in the Gazetteer.

Nettopus coromandelianus (Gm.) The Cotton Teal.

Occurs in the Gazetteer list.

Anas pœcilorhyncha pœcilorhyncha Forst. The Spot-bill Duck.

Occurs in the Gazetteer list.

Anas platyrhyncha (L.). The Mallard.

A few were observed in the first half of November on the Beas about Bhuin and Bajaura and on the Tirthan at Manglaur.

A few pairs are said to breed at the little marsh at Dalogi 7,000 feet in the Sainj Valley but the statement has not been verified.

Nettion crecca (L.). The Common Teal.

A very big flock of Teal was observed feeding on water-logged cultivation at Bhuin on November 4.

Querquedula querquedula (L.). The Garganey.

Occurs in the Gazetteer list.

Mareca penelope (L.). The Wigeon.

A small party were observed on the Beas at Bajaura on November 12.

Chaulelasmus streperus (L.). The Gadwall.

A few parties were seen on the Beas from Bajaura to Larji during the first half of November.

Spatula clypeata (L.). The Shoveller.

Two or three were seen at Bajaura and Oot on November 12 and 16.

Marmaronetta angustirostris (Mên.). The Marbled Duck.

Netta rufina (Pall.). The Red-crested Pochard.

Nyroca ferina (L.). The Pochard.

Nyroca rufa (L.). The White-eyed Pochard.

Nyroca fuligula (L.). The Tufted Duck.

The names of these 5 ducks all occur in the list in the Gazetteer.

LADY [ELIZABETH] GWILLIM,

Artist and Ornithologist

BY

CASEY A. WOOD, M.D., M.B.O.U.

[Reprinted from the *Ibis*, July 1925.]

For several years past I have been buying books, manuscripts, paintings, and other items for the Blacker Library of Zoology and for the E.S.W. Library of Ornithology, McGill University, Montreal, Canada. This search for literary material in zoology has finally narrowed itself to rare *desiderata*, the ordinary requirements of these special collections having long been satisfied; consequently, the quest has led me into out-of-the-way localities and into shops and other marts of trade not generally regarded as harbouring works of zoological interest. While following this plan last year I, at a venture, enquired of a small but select London dealer in *objects d'art* whether he had any drawings or paintings of birds or other animals. After a search in his cellar among much half-forgotten stock, he brought out a parcel containing about thirty small (10×14 in.) mounted and coloured drawings of Indian Fishes. Each mat bore an auctioneer's (or dealer's) printed number; a few were signed 'E.G.,' and upon still more were written legends (that Sir Henry Drake-Brockman later translated for me as Urdu) of the native names of the subjects portrayed. With the contents of this package I was shown a portfolio containing paintings in colour of a few Indian flowers, inscribed with both their English and systematic names. Pasted on one of the front pages of this portfolio was a leaf on which was written 'Elizabeth Gwillim, Madras, 1800-1806.' While I was examining these drawings and asking for more, a salesman happened along and said to the proprietor, 'I think that before I went to France in 1914 I saw a collection of bird paintings downstairs.' Shortly afterwards this clerk appeared bearing an immense, dust-laden, but extremely well made portfolio about five feet broad and four high. I noticed that it was brass-bound, provided with a safety lock and had a wide wooden back. It must have weighed thirty pounds. On it were painted barely decipherable initials and a date—'E.C.K. 1800.' The contents amazed and delighted me. I do not claim to be an art expert, but I realized at once that the paintings of Indian birds in the pockets of that giant container were by the hand of no mean draughtsman.

We are all well acquainted with the productions of brush and pencil wielded by 'ladies of quality' during the Georgian and early Victorian periods, and I fully expected to find these amateurish efforts displayed in the bird drawings, despite the rather favourable impressions made on me by the fishes and

flowers. But I was agreeably mistaken; not only were the birds—so far as I then knew them—faithfully depicted as to plumage and posture, but the backgrounds were painted in a fashion worthy of Keulemans or Grönvold. They were in water colour, carefully finished and on fine paper. Some of them were mounted, and all were numbered in the handwriting of the artist. Many bore (on the backs mostly) descriptive notes—measurements, colour-indications, and other data. Generally the systematic title, sometimes only the English or vernacular name, was appended. On some pictures there was no legend; on others only the genus was given. In addition to the completed drawings there was a small collection of unfinished sketches, evidently the usual 'studies' of the artist.

Some of the paintings bore faint, but elaborate, pencilled descriptions of the bird-subject, references to its habitat, life-history, nesting-habits, oology, food, sexual differences, etc., showing that the artist was, for her day and generation, a well-informed zoologist.

In every instance an appropriate background—Indian landscape, trees, shrubs, on which the particular bird was accustomed to roost, flowers, fruit or animal food on which it fed—was provided.

In addition to notations by the hand of the artist herself, the reverse of each portrait showed, as in the case of the drawings of the Indian fishes, a trade number, as if the collection had been prepared for public sale or auction.

After I had acquired this rather remarkable 'find,' I naturally tried to trace the paintings to their original source, but in vain: All the dealers could tell me was that they were purchased by them 'at a sale in the country' many years before, but exactly when and where they could not remember, and were unable to discover, as no record of the sale had been preserved. From time to time the portfolio had been resurrected and an occasional picture sold, mostly for framing but sometimes for decorating fire- and other screens! It was impossible to say who bought these odd lots.

I carefully cleaned the Gwillim pictures of their century of dust and other accumulations, and was much pleased to find that, chiefly owing to the fine quality of the drawing paper, there was very little 'foxing' visible, and that the well-constructed portfolio had further preserved them. The colours (including the iridescence of certain plumage) were barely altered; many a brilliant hackle, mantle, wing, and tail seem as freshly depicted as when they were painted more than a hundred years ago.

In all there were 121 drawings, and as the highest of Lady Gwillim's numbers was 201 (corresponding to 180 of the supposed dealer) it may be assumed that she had at one time and another painted at least the former number of bird-portraits, which before they fell into the hands of the former dealer had probably been reduced by sale or otherwise to 180.

The 121 pictures give one a fair idea of the avifauna—indigenous and migrant—of southern India. Comparing them with Jerdon's list and with the fuller catalogues of Blanford and Oates, there are, of course, many gaps—many genera are not represented at all; but, considering the circumstances, one can easily believe that

if the artist had lived a few years longer she would have made a gallery of Indian bird-pictures of the greatest scientific value, worthy to rank with the major collections of the world. Her sudden and early death, however, prevented the completion of the task, and probably consigned many of her best efforts to oblivion.

The importance of the collection may warrant a few remarks about individual pictures, as well as about the artist herself.

In the first place, almost all the drawings are life-size. If one reflects for a moment that Indian birds include many of the largest of the Old World avifauna, the reason for the immense portfolio I have described is manifest. For example, the painting of *Ardea cinerea* measures, exclusive of the mount, $33\frac{3}{4}$ in. \times 24 in. ; of *Dissura episcopus*, $25\frac{1}{2}$ in. \times 24 in. ; *Herodias alba*, $26\frac{1}{4}$ in. \times 20 in. ; *Ardea manillensis*, $26\frac{1}{4}$ in. \times $19\frac{7}{8}$ in. ; *Botaurus stellaris*, 30 in. \times $24\frac{1}{2}$ in. ; *Pseudotantalus leucocephalus*, 25 in. \times $21\frac{1}{4}$ in. ; *Gallus sonnerati*, $33\frac{1}{2}$ in.

The Accipitres are well represented and proportionately figured; even the largest Vultures are painted life-size.

It has been the proud belief of Americans—myself included—that it was our Audubon who first produced full-length portraits of the largest birds, and certainly the pictures of the male and female Wild Turkey, of Washington's Eagle, etc., and their exact reproduction in the elephant folio bear out that claim. However, so far as originals are concerned, we must now concede the palm to Lady Gwillim, who, so far as I know, is the first artist-ornithologist to paint full-sized and exact pictures of any considerable number of birds whose length exceeds, say, 35 inches.

I asked a number of trained ornithologists to examine and give me their opinion of the scientific and artistic value of the collections, among them Mr. Stuart Baker and Mr. Kirke Swann. The latter carefully revised the systematic names of the Birds of Prey, and expressed his complete satisfaction with the manner in which they had been depicted by the artist. Mr. W. L. Sclater also kindly looked over the portfolio and agreed that its contents would be of distinct value to students and would form a desirable addition to a research library. He pointed out that in the evident faithful delineation of *living* subjects the pictures formed a striking contrast to those common but really valueless drawings supplied by native artists of Indian birds. Moreover, they were likely to be free of those errors (discoloured and faded wattles, legs, ceres, mandibles, etc.) sometimes made by even our best artists who were obliged to make their drawings from bird-skins in which decided colour-changes had occurred after death. Lady Gwillim, having been in a position to command live birds or recently killed specimens, had been able to avoid such mistakes.

Having settled the status of her pictures, it remained to find out something about the artist. I have consulted all the ordinary and most of the extraordinary sources of information that would occur to one interested in the matter, but with meagre results. Many authorities—among them Sir Henry Drake-Brockman (I.C.S.), Sir William Foster of the India Office, Canon Bannister, the well-known genealogist of Hereford, whence Lady Gwillim's husband

came to Madras, the Librarian of the Inner Temple, and several others, who seemed likely to furnish some account of her life and career as the wife of a distinguished Indian judge—very kindly assisted me in this quest. In addition, I have carefully searched all the biographies, both British and Indian, at my command and have diligently explored, while in that city, all the Madras public libraries, including the files of the *Madras Government Gazette* and the records in the Connemara Library at Madras. Through the courtesy of the Garrison Chaplain, the Rev. C. E. De la Bere, I was permitted to examine the Parish registers of St. Mary's Church and to copy the inscription on Lady Gwillim's tomb. I regret to say that the information thus acquired is small in amount and of disappointing character. The *Government Gazette*, the only newspaper published in the Presidency at the time, gives only the usual formal notice of the lady's death, omitting her maiden name, and birth-place. As Sir W. Foster remarks, obituaries of women of title were not fashionable at the beginning of the nineteenth century.

Briefly, all vital data I have been able to secure up to the present time is as follows:—Elizabeth (maiden name unknown) Gwillim was born (about) April 21, 1763, and came to Madras in the year 1800 with her husband, Sir Henry Gwillim (just made K.C.M.G.) of Hereford, one of His Majesty's Puisne Justices of the newly-formed Supreme Court of Judicature. She died Dec. 21, 1807,* at the early age of 44, and was buried in St. Mary's Church, Fort St. George, Madras. Cut in the pavement of this historic church, near the front entrance, is this inscription:—

ELIZABETHA
PIA CONIUX
HENRICI GWILLIM EQ. AUR.
VIXIT ANN. XLIV. MENSES VIII
H. S. E.
MDCCCVII.

If any of my readers can add to the life-dates or personal history of this half-forgotten artist, the information will be gladly welcomed by the present writer.

*In both Mrs. F. Penny's *Fort St. George, Madras* and C. S. Cotton's *Monuments of Madras* this date is incorrectly given as August 7, 1807.

SOME NOTES ON THE SECOND EDITION OF THE FAUNA
OF BRITISH INDIA—BIRDS, VOLS. I AND II

BY

CLAUD B. TICEHURST, M.A., M.R.C.S., M.B.O.U., F.R.G.S.

VOL. I

Excellent as was the Fauna of Oates and Blanford—and few will deny the great help it has been to those of us in India who were ‘ornithologically inclined’, a new edition in light of the extended knowledge of the last 30 years was urgently required. To write such an Avifauna within the limits prescribed covering the whole of India (in the widest political sense) and dealing with every aspect of each species fully, is an Herculean—nay almost impossible—task for any one man. Only those who have studied the subject at all deeply can realize what an immense amount of work in the museum and library such a work entails. It will follow therefore of necessity that those who have specialized at all in any one department or region will find in the new edition of the Fauna statements which are not strictly accurate when applied to their speciality, and here and there they will find errors or slips have unaccountably crept in. Mindful of these facts and of Mr. Baker’s request (vol. xxx, 207–9) for helpful criticisms, I venture to put on paper these few notes which I have jotted down chiefly concerning N.W. Indian birds on reading through the *Fauna*—not that they pretend to be exhaustive—avoiding as far as possible any matters of opinion or subsequent publication. For convenience of reference I have copied the nomenclature used in the Fauna.

p. 21. Raven (*Corvus corax laurencei*). I cannot confirm the statement that the Brown-necked Raven replaces this bird in N.E. Sind. There are very few certain records of *ruficollis* in Sind.

p. 34. Sind House Crow (*Corvus splendens zugmayeri*). The distribution given of this bird is too wide to the west. In Beluchistan it is practically confined to Las Belas and Sibi plains. Zarudny did not meet with it in Persian Beluchistan nor has any one else met with it there or in the Mekran, so far as I am aware. Nor does it enter the hills to any extent. If it occurs at Fao in Iraq at all, it must have been introduced there; there is no recent record. On the other hand it is the form of House Crow over the greater part of the western Punjab plain, but to Afghanistan it must indeed be a straggler or occur in places just on the border. It must be remembered that old records of ‘Afghanistan’ (such as Griffiths’) did not refer to Afghanistan of to-day necessarily but often to Beluchistan (Sibi) or even Upper Sind.

p. 31. Rook (*Corvus frugilegus tschusii*). The Rook does more than occasionally wander to the Punjab plains. It is a regular winter visitor to the plain north of the Salt Range and occurs at Gujranwala, Ludhiana and near Lahore—as recorded by Whistler and Currie. Punjab specimens which I have examined cannot be picked out in any way from the typical race.

p. 29. Himalayan Jungle Crow (*Corvus coronoides intermedius*). It is difficult to see how small birds at Murree can be wanderers from the plains after breeding. No Jungle Crow breeds in the Western Punjab plains and the species is practically absent from Sind. The distribution of *levaillanti* therefore also needs emendation.

p. 38. Kashmir Magpie (*Pica pica bactriana*). It must surely be a straggler to the Simla Hills; at all events no one has met with it there of recent years.

p. 48. Indian Tree Pie (*Dendrocitta rufa*). I have already gone so fully (Ibis. 1922, p. 537) into the races of this bird that I will only say that if any races are to be recognized at all it is surely inconsistent to unite the largest and paler north-western bird (*pallida*) (W 152–170), with the smallest and darker

South Indian bird *rufa* (W 135-152). *Sclateri* and *kinneari* do not appear to me to be very distinct forms.

p. 70. Yellow-billed Cough (*Pyrrhocorax graculus*). The Himalayan bird of which Whistler has obtained a splendid series run much larger than European ones ♂ W 275-298, bill from base 35-38. ♀ W 262-266, bill 34-35. It should, at any rate for the time being, stand as *forsythi*.

pp. 73-77. Grey Tits (*Parus cinereus*, etc.). The Great Tit of Southern Persia *P. major blanfordi* belongs most distinctly to the *major* and not the *cinereus* group. I am unaware of any of the *cinereus* group in N. Arabia.

P. m. intermedius. The measurements given are too small for Baluchi birds, males of which measure W 73-78 T. 63-72. It is a large-winged bird with tail in proportion; some of mine have not fully grown tails so that 63 mm. is probably on the small side.

P. m. planorum. This supposed race wants further examination. I have examined the type, it is an unsexed winter bird with no original label, said to have come from 'South Punjab'! The wing measures 70 mm. It and another apparently from the same source (W 68) I cannot distinguish from *kaschmirensis*. No Grey Tit breeds in the plains of the Punjab, except perhaps on the Delhi side, nor in Sind.

A useful distinction in the various races of Grey Tit is the white wedge on the penultimate tail feather's inner web. This varies in depth in birds from the type localities pretty constantly.

In <i>cinereus</i>	10-20 mm.	up from tip	outer web black.
In <i>mahrattarum</i>	20-28	„ „	„ with a black line.
In <i>kaschmirensis</i>	30-40	„ „	„ no black line.
In <i>intermedius</i>	37-40	„ „	„ „

p. 82. Turkestan Marsh Tit. (*Parus palustris korejewi*). I can find no record of this bird in Beluchistan. I think that it would have been more helpful if, in all cases of very rare stragglers, the records had been given in full.

p. 85. Simla Black-Tit. (*Lophophanes rufonuchalis rufonuchalis*). The English name of this bird only has priority to recommend it, it being given by Jerdon. But it must be very confusing to the inexpert. This Tit does not occur actually in Simla, nor is the type locality Simla. Blyth says 'Tyne range beyond Simla presented by Capt. Hutton.' Stewart (*Zool.* 1886 p. 443) says he gave Hutton this bird and he procured it at Nagteeha two long marches beyond Landour though the date he gives (1861) must have been a slip. In the description it is said that there is a broad band down the abdomen 'black,' and the upper plumage is olive green; both these statements are copied from the Fauna ed.; and are misleading; if there is a broad black band as stated it must be rare as none of our large series have it, and olive green must have been a slip for olive grey.

p. 98. White-Cheeked Tit. (*Ægithaliscus leucogenys*). A reference is made to Whitehead obtaining this bird as low as 1,800 feet at Safed Koh on July 20. This is as recorded by Whitehead, but it is an obvious misprint or slip for 8,000 ft., and as such is nothing noteworthy. The Safed Koh being one of the highest mountains on the Frontier, no part of it of course is anything like so low as 1,800 ft.

p. 100. Penduline Tit. (*Remiz coronatus*). I know of no record in British Beluchistan of this species; Whitehead and Magrath obtained it in Kohat District, N.W.F.P.

p. 133. Velvet-fronted Nuthatch (*Sitta frontalis*). The distribution is given as the whole of India west of Bombay, Gwalior and Kumaon; it should of course be east of this line. The Himalayan bird is smaller than the Ceylon bird in wing and bill, and should stand as *Sitta frontalis corallina* Hodgson (J.A.S.B.V., p. 779. 1836 Nepal), Ceylon birds measure W 75-84. Nepal and Sikkim, W 71.5-77.

p. 199. Afghan Babbler (*Argya caudata huttoni*). The 'Jay River Hills' in Sind is the Gaj Nai, a hill water course in the Khirthar N.W. of the Munchar Lake. I too made the mistake (Ibis, 1922, p. 540) of uniting *celipes* with *Argya*

caudata caudata; it is perfectly distinct and an interesting form. It is rather larger than *caudata* and much darker on the upper parts with larger streakings. Since I wrote (*loc. cit*) I have seen a good series from Rawalpindi, Peshawar (type) Attock and Kashmir foothills (Punch-Gingri road). It is the race inhabiting the lower hills in the N.W. corner of India. Hume's original description is good. The status of this race seems to be on a par with that of *Molpastes leucogenys humii*.

p. 357. Grey Hypocolius (*Hypocolius ampelinus*). The distribution of this bird is still not completely known. It certainly is a summer visitor to Iraq, but Mr. Baker appears to suggest that it is resident in Persia moving south in spring; I know of no support for this.

pp. 383-389. Red-vented Bulbuls. (*Molpastes hæmorrhous*). The measurements given for the various races will need revision, and it may be noted that Bulbuls vary in size according to sex and so it would have been better to have given the measurements for each sex. The wing measurement of *pallidus* is given as 87-97; I measure a large series—♂ 96-101, ♀ 89-94.5. It is not clear from the distribution given that this bird occurs in S. E. Sind and in the plains of the Western Punjab. The measurements of *bengalensis* is given as 103-111; the few I have from Bengal measure 3 ♂♂, 101.5-104.5, 4 ♀♀ 93-99. *Intermedius* is given as '92-105, very few reach 100.' I measure 8 ♂♂. (100.5) 102.5-106, 7 ♀♀ (92), 98-102! In passing it may be noted that the description of *M. h. nigropileus* has been omitted.

p. 390. White-eared Bulbul (*Molpastes leucogenys leucotis*). The type locality cannot be restricted to the Punjab as it has already (J. B. N. H. S., xxviii, p. 271, March 1922) been restricted to Karachi in Sind. The measurements given (W 86-93) surely include the Perso-Iraq form *mesopotamiæ*; not one out of our 17 specimens (Sind and Punjab) exceeds 88. ♂ 81.5-88.

p. 394. Bengal Red-whiskered Bulbul (*Otocompsa emeria emeria*). This must be very rare at Simla; no one appears to have recently come across it though Simla is given in the first edition of the *Fauna* as within its range.

p. 422. Small Olive Bulbul (*Pycnonotus erythrophthalmus*). The distribution is omitted.

p. 423. Black-headed Bulbul (*Microtarsus melanocephalus*). The type locality given as 'Sandwich in mares australis' appears to need an explanatory note.

VOL. II

p. 23. It may be noted that the type of the genus *Saxicola* was designated by Selby in 1825 according to statement (p. 23) but on p. 38 it is accredited to Swainson 1827. The latter seems correct.

p. 24. (*Saxicola caprata*). It seems a pity and confusing to twist round the English names of the Chats. The *caprata* group were well known as Bush-chats and are well named so, conversely the *torquata* group are well known as Stonechats (though Oates called them all Bush-chats) and the English race is actually referred to on p. 27 as the English Stone-chat. Moreover on p. 25 *S. caprata burmanica* is referred to as the Pied Chat while we have another Pied Chat (*E. picata*) on p. 42 and yet another (*E. leucomela*) on p. 45!

p. 41. Hume's Chat (*Enanthe alboniger*). I cannot make out the tail measurements given as 55-71. My Sind birds measure ♂ 71-75, ♀ 66-69 from base of tail.

p. 43. Pied Chat (*Enanthe picata*.) Lest there be any doubt about my statement quoted on p. 43 that this bird is the breeding bird of Eastern Persia, I may say I have examined over 30 specimens, breeding or juveniles, from East Persia, from Serakhs in N. Khorasan to Kohi Taftan in the south. Moreover Blanford also found it breeding in the Kerman and Niriz districts.

p. 47. Barnes' Chat (*Enanthe melanoleuca melanoleuca*). One despairs of the fixity of the Latin name of this bird. *Melanoleuca* had been long used for the eastern form of the Black-eared Wheatear till changed by Hartert to apply to Barnes' bird (Vog. Pal. F., p. 690). It was changed back again in the B. O. U. list Addenda (Ibis, 1921, p. 314) to apply to the Black-eared Wheatear

but now Mr. Baker changes it back again to Barnes' chat. So the nomenclatural pendulum swings! and 'fixity' seems no nearer.

The wing measurements given (85-92) for this bird are too small, as regards Beluchi specimens at all events, as 8 males I have measured vary from 92-96.

p. 53. Tibetan Desert Chat. (*Enanthe deserti oreophila*). Is it certain that this bird winters in Sikkim (whence Stevens does not list it) or anywhere else within Indian territory proper? From my investigations it appears to be a summer visitor to its breeding haunts with a well defined but comparatively narrow winter range which extends from S. Afghanistan, Seistan, Eastern Persia and the Mekran to S. Arabia and Sokotra. It appears to miss the plains of N. W. India altogether.

p. 54. Red-tailed Chat (*Enanthe xanthopyrmyna chrysopygia*.) So far as I am aware this chat does not breed in British Beluchistan. I can find no record of it.

p. 79. Guldenstadt's Redstart (*Phoenicurus erythrogaster grandis*). The distribution given of this bird does not seem quite clear. The term 'N. E. Frontier' is vague and is used in various places with varied meanings. Here perhaps it means the N.E. Frontier of Kashmir?; on p. 124 it apparently means the N.W. Frontier Province. This bird must be a mere straggler to the plains of the Punjab.

p. 84. Western Red-spotted Bluethroat (*Cyanosylvia suecica suecica*). This race may of course occur in the plains of N.W. India, but I think it cannot be common; of a large series in Whistler's and my collection from Punjab and Sind not one is referable to this race.

p. 85. Eastern Redspotted Bluethroat. (*C. s. pallidogularis*). Is it quite certain that this Bluethroat breeds in Ladak? I examined Ludlow's specimens and considered that they were *abbotti*.

pp. 87-88. Turkistan Nightingale (*Luscinia megarhyncha golzii*). On p. 87 this bird is referred to as the Persian Nightingale though named correctly in the heading. The measurement for the tail of the European bird 'about 65' is too small. It runs up to 72 mm. The length of the first primary varies much in European birds and is of itself no guide; sometimes it is equal to, sometimes it is longer than, the primary coverts. *L. m. africana* is referred to as a larger bird than *golzii* where as it is smaller.

p. 105. Blue-headed Robin (*Adelura caeruleocephala*). I can find no record for this bird in Beluchistan.

p. 113. Indian Magpie Robin (*Copsychus saularis saularis*). 'The whole of India' is far too sweeping for the distribution of this bird. There are enormous areas in N.W. India where it is quite unknown or very local and rare.

p. 123. Blackbird (*Turdus merula*). It may be a matter of opinion but it seems to me hardly necessary to consider all these six birds as races of the European Blackbird, and the explanation of their origin is somewhat fanciful. Though *maximus* and *albocinctus* approach on the limits of their distribution, there is no intergradation between the two, which suggests to my mind that they are two distinct species. Again *simillimus* and *bourdilloni* both breed in Palni Hills and so can hardly both be races of *merula*.

p. 143. Salvadori's Thrush (*Turdus obscurus subobscurus*). The wing measurement should surely be 133 not 153 (=5.25 inches).

p. 153. (*Arceuthornis*). The splitting of genera is a question no two people agree on and so, long as this is so uniformity in nomenclature is unattainable. Whether one splits off these three Thrushes from the genus *Turdus* may be purely a matter of opinion; at the same time I think the Fauna is the only modern work in the English language which would put, say, the Black-throated Thrush and the Fieldfare into separate genera, and I fancy few will be disposed to follow this lead. The division between the two genera—sexes alike and sexes not alike—seems too artificial and in some cases a question of degree. Thus on colour the sexes of the Fieldfare are almost precisely alike but in most cases the sexes can be differentiated; on the other hand some males of the

Black-throated Thrush cannot on colour be differentiated from the females ; moreover, in winter habits this bird is very 'Fieldfare-like.' I should be rather puzzled to know which genus to place the Ring Ouzel in !

p. 156. Fieldfare. *Acrocephalus palustris*. The distribution given includes this bird as breeding in the Faroes. If it has ever bred there it must be very exceptional. Müller did not record it as a breeding species, nor Knud Andersen in many years reports on the avifauna of Farøe.

p. 173. Blue Rock-Thrush (*Monticola solitaria*). As I have already pointed out the oldest name for the Transcaspian bird is Blyth's *longirostris*. Mr. Baker makes this race and *pandoo* almost the same size in wing measurement. From a large series of males of both races, I make the measurements to be *longirostris*, ♂ 121-127, *pandoo*, ♂ 117-123. (Ibis. 1922, p. 641). It is of interest to note that Mr. Baker considers the Transcaspian bird to breed on the Beluchi frontier and Kurram Hills whereas Col. Meinertzhagen has identified Beluchi breeding birds as *pandoo*. One would have expected the same race in both areas.

p. 177. Rock-Thrush (*Monticola saxatilis*). Does this bird winter in N. W. India, if so it must be excessively rare? There were no records in the 'First Edition'. Mr. Whistler, never found it wintering in the Punjab and in Sind it is a double passage migrant, as it is in Mekran, almost certainly *en route* to and from Arabia and Africa. Whitehead too only met with it on passage in N. W. F. P.

p. 187. (*Prunellinæ*). Because the generic name *Accentor* cannot be used there is no reason why the English name should be thrown aside ; e.g. we still use the English name 'Ibis' though the Latin generic name *Ibis* has been discarded for *Threskiornis*. Moreover the two groups have for long and widely been known as Alpine Accentors and Hedge Accentors. To call the Alpine birds *Hedge-Sparrows* certainly is *not* technically correct.

p. 197. *Prunella strophiatea jerdoni*. This bird may of course breed in Beluchistan though I know of no record. It breeds as near as S. Waziristan.

pp. 201-3. Eastern Spotted Flycatcher (*Muscicapa striata neumanni*). The statements that this bird is 'to some extent migratory' and is a very rare straggler to India in winter hardly describes its Indian status. It is *inter alia* a summer visitor to N. E. Beluchistan and occurs not uncommonly on autumn passage in the plains of N. W. India. I know of no record of it wintering ; it appears like several other species to winter out of India altogether. The Persian birds which I have examined certainly belong to this form.

pp. 203-6. *Hemichelidon*. Mr. Baker is certainly wrong in discarding Hodgson's names for the Sooty Flycatchers. Hodgson instituted the genus *Hemichelidon* for their reception and described the birds as *Hemichelidon fuliginosa* and *Hemichelidon ferruginea*. If the genus *Hemichelidon* is kept separate from the genus *Muscicapa*, which Mr. Baker does, then Hodgson's names must be used and are not invalidated by *Muscicapa fuliginosa* and *Muscicapa ferruginea* of prior date. If on the other hand *Hemichelidon* is merged into the genus *Muscicapa*, as Dr. Hartert does, these names of Hodgson cannot be used as the combination is preoccupied. This is yet another example, which occurs at least two other times among the Indian Flycatchers alone (viz. *Siphia hyperythra* and *Cyornis melanoleucus*), of the impossibility of uniformity in nomenclature so long as authors are divided on the limitations of genera.

pp. 255-6. Grey-headed Flycatcher. *Culicicapa ceylonensis*. The young in nestling plumage is a slightly duller and paler replica of the adult. No barring and no spotting ! So it apparently is not a Flycatcher ! The birds from N. India are easily separable from Ceylon birds by their paler yellow underparts, paler grey head and throat and paler mantle. It is not quite correct to say that this bird does not occur in the Punjab ; it is common of course in the Himalayan Punjab but it also straggles down into the plains in winter and is recorded from Jhang, Amballa, Lahore and Gujranwala.

I am very doubtful if the name *C. c. meridionalis* can stand for the Malay Peninsula bird. There is no doubt that it is distinct (as pointed out) from the Ceylon bird but three races have already been described from islets off Sumatra, viz. *percnocera* (Pulo Si Maloc Is.), *amphiala* (N. Pageh Is.) *pellonota*

(Samasura Nias Is). It seems questionable from the descriptions whether they are different to the Malay and Sumatra bird and if not one of these names will have priority over *meridionalis*.

p. 264. Indian Paradise Flycatcher. *Terpsiphone paradisi paradisi*. The original reference of this bird should be *Corvus paradisi* (Linn. S. N. ed. x, p. 107) (1758). Linneus' first reference is to Ray. (Av. 195. t 2. f 13) where the bird is described and figured from Fort Saint George, Madras, so that there was no need for Mr. Baker to have restricted it (p. 266) to Madras.

p. 271. North Indian Black-naped Flycatcher *Hypothymis azurea styani*. The distribution given for this bird 'All India north of *sykesi*' is far too sweeping. Although I obtained an unaccountable waif in Sind it does not occur so far as is known in the greater part of N. W. India at all.

p. 277. No. 700 is called the White-browed Fantail warbler instead of Flycatcher. The same muddle is made on p. 281, last para.

p. 283. *Laniidæ*. It is not quite correct to say the shrikes have an autumn moult only, i.e., do not moult in spring. *Lanius minor*, *L. nubicus*, *L. isabellinus* all certainly have a spring body moult and probably *L. vittatus*.

p. 285. Grey shrikes (*Lanius excubitor*). I can see no constant difference in the amount of white on the inner webs of the secondaries and so the key for *aucheri* and *pallidirostris* will fail. The tail length for adults of *lahtora* given as 100-115 is too small. On a good series I measure it as 117-129.

pallidirostris is said to have occurred only once in India; besides this Punjab specimen, I consider it to have been found at least once in Sind; but if Beluchistan is included in India as it seems to be (as the sole occurrence of the woodchat rests on a Quetta specimen) then it should be included as the breeding race in and a not uncommon summer visitor to the hills of N.E. Beluchistan. The nest was described in the Ibis., 1920, p. 150.

pp. 294-7. Rufous-backed Shrikes (*Lanius schach*).

p. 294. The characters relied on for retaining *L. tephronotus* as a species are not constant. That *erythronotus* never lacks the white speculum and has the central tail black and that *tephronotus* always lacks the white speculum and has the central tail brown, no doubt, is true for the majority of the specimens. I have seen *erythronotus* with the white speculum absent (specimens Jhang, Jhelum, Pindi, Quetta, Srinagar) and birds which otherwise are typical *tephronotus* with a white speculum or a trace of it; in some states of plumage *erythronotus* has the central tail brown not infrequently, while in *tephronotus* I have seen the central tail nearly black or the terminal half black. Then again at lower elevations in the N.W. Himalayas one finds typical *erythronotus* but in Lahul 10,000 feet typical *tephronotus* (Whistler) but in the intermediate elevations birds are intermediate between the two and a series can be formed intergrading from one extreme to the other. It is not a question of two species interbreeding as typical birds of the two forms do not occur together. I certainly should consider *tephronotus* to be the high elevation race of *schach*. 'Races of *schach* from Europe'; I have never heard of this bird in Europe.

p. 295. The tail measurements of *erythronotus* are too small; a large series measured as 108-120.

p. 297. The nestling of *tephronotus* is easily distinguishable from that of *erythronotus*; the upper parts are many shades darker grey.

Tephronotus would appear to be very rare in winter in the Punjab plains. The type localities given for these two birds are not correct. Vigors described both birds from the 'Himalayas', good reason for showing that these birds came from the area Simla-Almora has been given in the Ibis (1924, pp. 468-473). Moreover Lucknow certainly is not in the Himalayas nor should I consider Gyantse to be either.!

p. 300. Woodchat Shrike (*Lanius senator niloticus*).

The statement that this bird inhabits the more barren and stony hill sides in Mesopotamia and is not found except as a winter visitor in the open plains is hardly correct; nor have I found any evidence of it breeding in Mesopotamia.

So far as we know it does not winter in Mesopotamia at all but is a through migrant when it usually inhabits much the same sort of country as woodchats do elsewhere (J. B. N. H. S., 1921, p. 235).

p. 301. Brown Shrike (*Lanius cristatus cristatus*). A good distinction between this bird and *isabellinus* and *phoenicuroides* lies in the comparative shortness of the outer tail feather. From the tip of this feather to the tip of the central tail feathers in *cristatus* measures 19–24 mm.; in the other two birds 10–16 mm.

p. 302. Pale Brown Shrike (*Lanius cristatus isabellinus*.) The type locality given as 'Kampada' is hardly recognizable; Gonfoda or Kunfuda seem to be the more usual variants in spelling. A tail measuring 64 mm. surely cannot be fully grown? in 12 specimens measured the length is 80–86 mm.

p. 303. Rufous Shrike (*Lanius cristatus phoenicuroides*).

This bird probably is commoner in India than is recognized at present, but I question whether it will be found to be so in *winter*. In Sind, Punjab, Mekran, as in Iraq, it is a decided passage migrant and it winters as far south as Abyssinia and the Sudan.

p. 329. Small Minivet (*Pericrocotus peregrinus peregrinus*).

I have already (Ibis., 1922, pp. 613–614) gone so fully into the question of placing the type locality at Amballa that for the benefit of those who have not seen this work I will only repeat that when Linnaeus described this bird in 1766 it was very unlikely that he described a Punjab bird at all; and Amballa birds are intermediate between *peregrinus* and *pallidus*; therefore Amballa is a very unfortunate choice.

pp. 356–7. Black Drongo. (*Dicrurus macrocercus*). I am doubtful whether the Himalayan race is sufficiently distinct. No reliance can be placed on average measurements when sex is disregarded or is unknown, as obviously one group might contain more males or more females than the other. The maximal wing length and the minimal tail length in the two supposed races are precisely the same and I cannot find on my own measurements any difference in bill length between the two.

pp. 385–7. Grey-backed Warbler (*Agrobates galactodes familiaris*).

In the generic description, p. 385, it is said that the only representative of the genus in India is a 'somewhat rare winter visitor' but on p. 387 it is said to 'probably breed practically wherever it is found.' The two statements hardly tally. It is very curious and interesting that this bird should breed not uncommonly at Multan; no one else has met with it breeding either in Sind or the Punjab, though Mr. Baker suggests that it may be resident. Doig, Butler, Murray and myself all failed to meet with it in the breeding season in Sind nor has Mr. Whistler discovered its nest in the Punjab. Outside Major Lindsay Smith's records it is only known in the plains as a passage migrant and in two years I only came across it, and then not uncommonly, between September 3 and 24. Mr. Baker does not quite do me justice when he says I have not stated from where to where these migrants were going. I stated my opinion twice! (Ibis., 1920, pp. 531 and 547.) The description of this bird's breeding haunts given as 'in *low stunted* bushes that do not average 20 feet high' makes one wonder what the *really fine* bushes at Multan are like! I saw none there to merit this somewhat 'Americanized' comparison!

pp. 389–90. Indian Great Reed Warbler (*Acrocephalus stentoreus*).

As Mr. Baker says—more breeding specimens are required to elucidate the two forms. I think it is very doubtful if the Sind breeding birds will be found to be *amyæ*; a large series which I collected in Sind are without doubt *brunnes-cens* though none are breeding birds; yet if *amyæ* is resident there one might expect that one or two out of a winter series to belong to this race. Gonda in the U. P. is included in the breeding range of *amyæ* but a March bird thence in Mr. Whistler's collection is as pale as Kashmir breeding birds and no smaller than some; a worn breeding bird is too worn to say much about.

p. 393. Blyth's Reed Warbler (*Acrocephalus dumetorum*).

It is stated that the second primary is equal to the eighth or is a little shorter. In a large series I find the second primary is just shorter than the 5th or than the 6th. The type locality (omitted) should be 'India.'

p. 401. Streaked Grasshopper-Warbler (*Locustella lanceolata*).

The type locality is given as 'Mainz, Irtum.' The latter word does not appear to be a locality at all but a German word (mis-spelt!) meaning an error! The correct type locality should be 'Russia.'

p. 402. Eastern Grasshopper-Warbler (*Locustella naevia straminea*).

This bird may occur more commonly in N. W. India than records indicate but to say that it occurs 'in great numbers' gives rather an erroneous impression. In three winters I saw a few at one spot only, a certain reed-bed on the Munchar Lake and nowhere else in Sind while in the Punjab Mr. Whistler has only met with it at one place.

p. 411. Indian Tailor-Bird (*Orthotomus sutorius sutorius*).

Tail length, 28-112 mm. (summer) is surely a mistake? I have not seen a tail in the breeding season as short as this.

p. 417. (*Luscinola*). Though some ornithologists unite *Tribura* with *Luscinola* I do not think anyone nowadays puts *L. melanopogon* into any other genus than *Luscinola*; indeed it is the type of the genus.

p. 427. Assam Wren-Warbler. The original reference gives the sub-specific name as *assamensis* instead of *austeni*.

pp. 442-5. The Tree-Warblers (*Hippolais*).

The tail length of *H. rama*, given as 47-52, is too small and is much the same as that given for *H. scita*, consequently the comparative shortness of the tail in the latter species is not emphasized though I have found it to be a useful character. I measure *rama*, T. 52-57, *scita* T. 45-52.

As I have already pointed out the evidence for the inclusion of *Hippolais pallida elaeica* in India is very flimsy and rests on Dresser's statement that he had seen a bird from Sind. Some error is possible since no one else has ever obtained it in India. The two birds referred to as being in Col. Meinertzhagen's collection are *not* in his collection but in the Quetta Museum and neither of them is *p. elaeica*.

Hippolais obsoleta is a bird of mystery but if it has a tarsus of 30 mm. it must be a good species!; but an alleged specimen from Sind identified by Seebohm is rather scanty evidence for admission to the Indian list. If this species is a good one the distribution given—Fao, Turkestan and Sind—is a very extraordinary one for a bird described from *Moscow*!

p. 447. Indian Whitethroat (*Sylvia communis icterops*).

Does this bird *winter* in India as stated? Oates only knew of September specimens and Mr. Whistler and myself have only met with it in September and October on passage.

p. 449. Desert Warbler (*Sylvia nana nana*).

The reference given as '*Sylvia affinis* Blanf. and Oates; p. 396' is of course a slip. A 'third hand' clutch of eggs without the bird and without precise data seems somewhat slender evidence for inclusion as a breeder within Indian limits.

p. 457. Indian Lesser Whitethroat (*Sylvia curruca affinis*). I am very doubtful whether this bird breeds at Quetta, Gen. Belham informed me that he may have been mistaken, though he thought he sent breeding birds to Bombay Museum. Enquiries there however have not brought any to light.

p. 452. Small Whitethroat (*Sylvia curruca minula*). This too is stated to breed in Beluchistan; I know of no good record of it. If both these Whitethroats do breed in Beluchistan they cannot both be races very well of *curruca*.

p. 453. Genus *Phylloscopus*. These birds are of course *winter* visitors to the plains, not summer as stated.

The key in the first edition of the *Fauna* failed because it tried to separate birds into one group which have dark heads and coronal bands from those which have not, and into the first group was put *Phylloscopus humii*. The same fault has been repeated in this edition. The majority of specimens of *humii* have the coronal band and the dark part of the crown so ill defined that they are easily put into the wrong division of the key. The key in the second edition I think would work better if for the group with 'no coronal bands'

were substituted 'no wing bars' and for those 'with coronal bands', 'wing bars or a trace of bars present.' But even with this modification a beginner would get mixed up with *sindianus* and *neglectus*.

p. 456. Tytler's Willow Warbler (*Phylloscopus tytleri*). The winter distribution as regards Western India needs modification; this bird is not found in Sind nor can I find any record of it in the Punjab plain. Like some other species it perhaps reaches the Himalayas via the Siwaliks as suggested by Mr. Whistler in other species.

pp. 456-8. (*Phylloscopus collybita*). Mr. Whistler has never been in Ladak and therefore he could not have obtained *Phylloscopus sindianus* breeding there! Specimens which he obtained breeding in Lahul I certainly think are *tristis* as they still retain a trace of the greenish tinge on the wings. When all green tinge has worn off the plumage of breeding birds it is practically impossible to differentiate the two races. In the field the alarm note might be a good guide as Brooks pointed out a very distinct difference.

In winter *sindianus*, so far from 'keeping to semi-desert and poorly wooded tracts,' in Sind is found in the best wooded parts of the province—thick babool jungle in the Indus and canal areas. In anything approaching desert I never saw it as I have already pointed out in the 'Birds of Sind.'

p. 460. Olivaceous Willow-Warbler (*Phylloscopus griseolus*).

By a slip this is called the Olivaceous Tree Warbler. Far from being 'doubtless a common winter visitor to the Punjab' I think it will be found that like *Phylloscopus tytleri* it does not normally winter in these plains at all; at any rate that is in accordance with such evidence as we have as, though Mr. Whistler has found it on migration in the Punjab plain, he has never found it wintering there.

p. 469. Hume's Willow-Warbler (*Phylloscopus humii humii*).

This Willow Warbler does not breed or even occur in Beluchistan so far as is known.

p. 470. Crowned Willow-Warbler (*Phylloscopus humii præmium*).

In view of the other Crowned Willow-Warblers it seems a pity not to have altered this name to the name accepted in England, viz., the Yellow-browed Willow-Warbler.

In the foot-note Mr. Baker says that *Phylloscopus inornatus* of Blyth cannot be used for this bird as Blyth stated subsequently that it does not apply to this species. I do not know where he stated this but how does it alter the facts? Blyth in his 'Catalogue' places *inornatus* as a synonym of *modestus* (= *proregulus*) which, from the description of *inornatus*, could not be correct. He evidently realized this as in a copy of his Catalogue, Blyth altered this in his own handwriting, putting *inornatus* as a synonym of *superciliosus*. Because an author thinks he has made a mistake and says that a bird described by him, he thinks, is in reality another species, the name is not invalidated if the description holds good. For instance Blyth thought erroneously that his *Turdus dissimilis* was only the male of *Turdus unicolor* but that does not invalidate Blyth's name (cf. F. B. I, ed. ii, vol. ii, p. 141). So too Hume renounced his *Pellorneum minus* and said it was only *P. ochraceum*, this was afterwards proved to be wrong and *P. minus* stands.

Mr. Baker cannot separate *Phylloscopus mandellii* from this bird; if that is so then *præmium* is the same bird as *mandellii*, and the latter name has thirty-six years priority over *præmium*! To add to the muddle on p. 47 he calls the bird *P. h. superciliosus*.

p. 490. Kashmir Grey-headed Warbler (*Seicercus xanthoschistus albosuperciliaris*). Mr. Baker may possibly have some source of information not available to me concerning Beluchistan, but like several other species mentioned in the Fauna, I can find no record of this species within the Agency. Most of Beluchistan is quite unsuited to it.

p. 507. White-bellied Bush-Warbler (*Horornis albiventris*). The type locality is given by a slip as Kashmir; it should of course be Hengmai, Manipur Valley, as stated under 'Distribution'

p. 520. Sind Hill-Warbler (*Suya crinigera striatula*). This bird certainly does *not* inhabit the plains. It may wander down in winter to almost plains-level, but it does not leave the hills.

p. 526. Indian Streaked Wren-Warbler (*Prinia gracilis lepida*). The distribution of this bird extends at least to the United Provinces whence I have a breeding bird from Futtehgurh.

p. 533. Aboo Jungle Wren-Warbler (*Prinia sylvatica rufescens*). The name *Prinia rufescens* cannot be used for this bird as it is preoccupied by Blyth's *Prinia rufescens* (= *Franklinia rufescens*) see p. 427. Hume described this Wren-Warbler from Mt. Aboo, Garhwal, Satara District and Etawah (*Ibis*, 1870, p. 111) and he described it again as *Drymoipus insignis* (S.F., i, p. 10) from Mt. Aboo and Central Provinces. Hume was certainly confused (and little wonder!), by the seasonal and age plumages of this Warbler and described twice over the Mt. Aboo bird and mixing up with both, *Prinia sylvatica sylvatica*. It is perhaps questionable whether Hume's names can stand, but as he indicates a greyer tone of plumage in *insignis* and cites Mt. Aboo as a locality, I think perhaps the name *Prinia sylvatica insignis* may stand for this bird which I think is by no means restricted to Mt. Aboo, but to which place I now restrict the type locality.

p. 539. Himalayan Goldcrest (*Regulus regulus himalayensis*).

There is no need to restrict the type locality to Srinagar as Mr. Baker has done, as the correct type locality, even if not well known, can easily be found out. Jerdon described this bird from Blyth's MS. In the *Ibis*, 1870, p. 168, Blyth says he had only seen one specimen which he believes was procured near Simla. In the *Zoologist*, 1886, p. 443 there is the account of obtaining this specimen by Surg.-Gen. Stewart near Kotegurh and this must be the type locality.

LOWESTOFT, July 1925.

REVIEWS

'INDIAN BIRD LIFE' BY DOUGLAS DEWAR. (John Lane, 7s. 6d. net.)

We have again to chronicle the appearance of a new book on Indian Birds by that prolific writer Mr. Douglas Dewar, and have no hesitation in pronouncing this youngest of his literary offsprings as one of the most interesting and valuable.

Its theme is succinctly described by the sub-title 'The Struggle for Existence of Birds in India'. Mr. Dewar has set himself to analyse the various causes that work to prevent the indefinite increase of birds by the rapid mathematical progression of reproduction. He takes us chapter by chapter through the natural causes that work to destroy birds, their eggs and their offspring; and as he goes he cites chapter and verse for his assertions, drawing his examples from his own observations and from the published observations of the various field naturalists that have worked in India.

The purpose of the book is to fill a supposed gap in the writings of Darwin. Darwin enunciated the fact that a struggle for existence must invariably be the corollary of the high rate at which all organisms tend to increase; at the same time he admitted our general ignorance of the nature of the struggle and the exact checks that operate in any particular instance.

Mr. Dewar has set himself to collect some of the evidence that bears on these points in India; and India should surely be one of the best fields for this enquiry with its prolific fauna and its prodigal waste of life.

The chapters before us show how a bird is at the mercy of destructive forces in every stage of life. The adult bird itself suffers from predatory birds and animals, from the greed or the sport of man, from disease and accident and from the forces of Nature acting on the bird direct or through its food-supply. The accidents that befall nest and eggs and young are each in their turn discussed at length.

This tragic recital occupies the main body of the book and is followed by a chapter which endeavours to appreciate the factors contributing to the success or failure in the struggle for existence of certain species like the *Mynah* or the Pink-headed duck.

In the final chapter Mr. Dewar arrives triumphantly at his conclusion that the checks on the multiplication of a species are purely fortuitous in their application to individuals; and that being fortuitous they must therefore have little or no effect on the colour or anatomy or physical characteristics of the species on which they operate, though he admits to a certain effect on features connected with breeding, migration or character.

The book then is in the nature of an attack on the theory of Natural Selection. It is however a reasoned and legitimate attack far removed from the irritating side attacks and pin-pricks on the Darwinian theories that disfigure most of Mr. Dewar's writings. Opinions may differ as to the value of the evidence cited by Mr. Dewar and it is evident that the book has been written hastily without sufficient study or material. The conclusions reached are not necessarily correct. But there can be no doubt that Mr. Dewar has written an interesting book which he might well use as the ground work for a more careful and more important study.

H.W.

'MANUAL OF THE BIRDS OF CEYLON' BY W. E. WAIT, M.A., F. Z. S.,
M.B.O.U., Ceylon Civil Service. (Ceylon Journal of Science, Rs. 10.)

Those of our members who are resident in Ceylon or interested in the Ornithology of that country will have every reason to welcome this Manual of the Birds of Ceylon which Mr. Wait, our member, has prepared on the suggestion of the Director of the Colombo Museum. For some time past there has been need of an up-to-date text book to convey in handy form the advances in method and knowledge since Legge's magnificent but expensive work was published. That work has long since been out of print and is now difficult

to procure, and it was written before Mr. Stuart Baker and other writers introduced the trinomial system to workers in the East; and though information regarding the birds of Ceylon is available in the excellent volumes of the Fauna of India series it is a cumbersome and unsatisfactory arrangement that workers in Ceylon should have to satisfy their needs by delving in a work which applies to so large and diverse an area as that treated of in the Fauna.

Now in Mr. Wait's book they have an excellent and compact hand-book to the birds of Ceylon, with the further merit that its price is such as to bring it within the reach of every student. The usual stumbling block experienced in a work of this nature that the cost of reproduction prohibits a liberal series of illustrations has been partly avoided by securing the services of the staffs of the Museum and the Survey Department to produce a series of plates which illustrate some of the most typical representatives of the various orders and families found in Ceylon. Although the drawings are stiff and conventional they undoubtedly serve their purpose, and it would therefore be unfair to overstress the fact that they share the decline in the standard of illustration which has been rather a marked feature in post-war ornithological publications. An excellent Survey Department map has been included.

For the text of the book we have nothing but admiration. Mr. Wait's introductions to the book, to birds and their classification, and to the various orders and families, are each admirable in their way imparting scientific facts in language which combines scientific accuracy with plain intelligibility to the unscientific student; while his descriptions of each individual species are well framed to supply exactly the information required by the beginner and by the ordinary student. The advanced specialist will not find here or elsewhere in the book all that he requires, but for this minority the book has not been written.

The account of each species commences with the correct names scientific and English, and the local vernacular names when such exist. Those are followed by the first reference to the name now used and by the references to the species in Legge's *Birds of Ceylon* and in the first edition of the *Fauna of British India*, so that where changes of nomenclature have occurred old friends may be recognized under their new names.

Next the bird is described in simple but clear terms, the differences between the sexes being clearly indicated and some idea being also given where necessary of the immature plumage and any marked seasonal changes. The colour of the soft parts and the average measurements in inches are also given.

The paragraph on distribution then describes very briefly the general distribution of the species and so far as is known its status and distribution in Ceylon and where relevant its connection with Southern Indian forms; and we are glad to see that more care has been taken than is usual in a work of this nature to indicate the times of arrival and departure of the migratory species. It is an interesting fact that all migratory birds in Ceylon are winter visitors, as the island lies at the terminus of the routes which run through India from the north, so that there are no passage migrants to the island and no summer visitors; a fact that must greatly simplify all ornithological observation in the island. Apparently also owing to the absence of any well marked procession of the seasons the hill birds do not indulge in any altitudinal movement, but this point Mr. Wait does not seem to have made clear. The Ornithologist in Ceylon has to confine his speculations chiefly to the effects on distribution of the rain belts.

The account of each species then finishes with a short paragraph on the habits and nidification written in clear and homely language. There are the usual keys and a helpful chart of the external parts of a bird and short glossary; and the various lists and indices are fuller and more satisfactory than in the usual run of popular books.

Altogether a satisfactory piece of work on which we heartily congratulate Mr. Wait.

H. W.

'A HANDBOOK OF THE BIRDS OF EASTERN CHINA' BY J. D. D. LA TOUCHE, C.M.Z.S., Etc. (Taylor and Francis.) Part II. December 1925. (7s. 6d. net.)

Students of the Avifauna of the Far East will welcome the appearance of the second part of Mr. J. D. D. La Touche's work on the birds of Eastern

China, the first part of which was reviewed at p. 211, vol. xxxi of the Journal. This new part deals with the families of the *Cinclidæ*, *Turdidæ*, *Muscicapidæ* and *Laniidæ* and is illustrated with four plates, which have been well reproduced from photographs. There is an interesting photo of captive Ruby-throats tethered on perches along a wall, and a picture of the nest of a Paradise Fly-catcher *in situ*. The remaining photographs show typical views of Chinese scenery.

The text of course follows the same lines briefly indicated in our review of the first part, though the accounts of some species are slightly fuller and more interesting than in that part. Many of the species and races dealt with are familiar to ornithologists in India. If Mr. La Touche is right in his identifications, some of our Himalayan species such as *Monticola erythrogastra* and *Chaimarrhornis leucocephala* are found from the Western Himalayas to China without even subspecific variation; though this seems curious in view of the fact that most Himalayan species have developed eastern and western races with their junction about Nepal.

We must however take serious objection to the inclusion of the Indian Grey Shrike, *Lanius ex. lahtora* in Mr. La Touche's list. It is included there as a fairly regular passage migrant and winter visitor purely on the authority of Père David. His record is supported by the statement 'This bird winters in India and, according to David and Oustalet, proceeds in summer to Central Asia'. Mr. La Touche is apparently troubled by this record for he cites Hartert, Dresser and the *Fauna of British India* in their perfectly correct rendering of the distribution of this race of the Grey Shrike. Now it is well known that our Indian bird is absolutely sedentary, and the difficulty in identification of the races of Grey Shrike is equally well known. This record, apparently unsupported by skins, should therefore have been refused by Mr. La Touche whose authority is sufficiently well known to entitle him to suppress older and obviously erroneous records. There are other signs in the work that the author should be more drastic in his treatment of previously recorded observations. It is the part of an experienced authority to sift the materials that have come down to him.

This second instalment of the book however supports our previous opinion that Mr. La Touche is conferring a great benefit on all students of Chinese Ornithology.

H. W.

'THE AQUARIUM BOOK', BY E. G. BOULENGER. (Duckworth, 3 Henrietta Street, London, W.C.).

This book by the Director of the New Aquarium at the London Zoological Gardens should appeal to the scientist and sportsman as well as to the amateur aquarist and man-in-the-street for whom it was primarily written.

In an introductory chapter, for the most part historical and descriptive, the author gives an interesting account of the elaborate mechanism behind the show tanks at the Zoo's Aquarium, the perfect design of which is the result of much patient investigation into the various methods adopted at the other large Aquaria of the world.

The book itself is divided into two parts, one dealing with Salt water, the other with Fresh water organisms. The author does not confine himself to fishes but includes chapters on salt and fresh water Invertebrates, and Aquatic Reptiles, Batrachians and Mammals. Of the vast number of backbone-less creatures which are common to all the waters of the earth, only a comparatively small proportion can be kept alive in an Aquarium. These, however, include many well-known species, and Mr. Boulenger recounts some very interesting observations which he has been able to make on some of the more lowly forms in his tanks.

His remarks upon the fighting habits of the Common Lobster and the description of the moult which takes place from time to time in these and other Crustaceans, show that he has taken full advantage of the unequalled opportunity for the study of the habits and bionomics of his charges. Another astonishing fact about the lobster which he tells us is that 'by stroking it in a certain direction a coma is produced, so that it remains perfectly still, when it can be made to rest tail upwards, standing on its beak and outstretched claws!'

There are many little occurrences of interest which take place in the daily round of the Aquarium; there now flourishes in the filter beds an ever-increasing colony of shrimps and prawns which have escaped down the overflow pipes of the show tanks and are being used as food for some of the other inhabitants of the Aquarium.

It is interesting to learn that the Common Cockle can leap in the water and in fact does so with great energy when its deadliest foe, the Starfish, is introduced into the tank. The leap is produced by a sudden opening and shutting of the valves of the shell.

Practically all fish become tame after a very short time in captivity. This is mainly attributable to regular feeding. There is one six-foot-long Conger at Regent's Park which can be lifted bodily out of the water after he has been fed and many of the smaller fish will come to the surface and nestle in the hand.

The chapter which deals with the Lung fishes and their transport to England in bricks of mud during the period of torpidity is intensely interesting. At present the Zoo is fortunate in having examples of *Ceratodus* from Australia, *Protopterus* from Africa and *Lepidosiren* from South America, all of which have been sent home in the dry state.

Mr. Boulenger has so many other fish stories to tell—all true—and all so equally enthralling that it is impossible to enumerate them here.

Suffice it to say that his book teems with interesting facts from cover to cover. It will appeal especially to those who have already made the pilgrimage to that 'Aladdin's Cave of Treasures' which lies beneath the Mappin Terraces at the London Zoo and will be a source of much enjoyment and instruction to those who have not yet had the opportunity to do so.

The illustrations alone make the volume worthy of a place on the bookshelf. Some are photographs taken in the Aquarium, by no means an easy feat, and there are many beautiful line and wash drawings done from nature by Mr. L. R. Brightwell.

'ACCOUNT OF A PHOTOGRAPHIC EXPEDITION TO THE SOUTHERN GLACIERS OF KANCHENJUNGA IN THE SIKKIM HIMALAYA', BY N. A. TOMBAZI, F.R.G.S. (Maxwell Press, Bombay).

The author has given a delightful account of a tour in an out of the way part of Sikkim in which he succeeded in reaching the pass between the Talung and Zemu valleys which Mr. Douglas Freshfield called the Zemu Gap. This was a notable achievement and has, we believe, never been accomplished before. The author is clearly an enthusiast in mountain scenery and travel, and among his descriptions he has interspread many useful hints on travel which will be of great assistance to those who follow him. Mountaineering in the Eastern Himalayas must always be difficult on account of the peculiar local conditions; except in winter, when extreme cold makes travel at great heights almost impossible, the rains greatly add to the hardships and discomforts of travel, and clouds detract from the enjoyment.

The author suffered severely from bad weather but his enthusiasm enabled him to catch the short but magic period just after dawn when the snows are clear, as his photographs bear witness.

We take liberty to doubt whether the birds seen by him below Jongri were Snow-cock (*Tetraogallus tibetanus*). These birds were seen in juniper forest somewhere below 11,000 feet, whereas the Snow-cock according to the *Fauna of British India—Birds*, rarely, if ever, descends below 15,000 feet. The writer once saw a brood of chicks in dwarf rhododendron at an altitude of about 14,000 feet, but it is doubtful if they descend below this altitude. It is more probable that the birds were Blood Pheasants (*Ithagenes cruentus*); their tameness also points to this probability.

The author actually saw at a distance of 300 yards what the coolies described as a 'Kanchenjunga demon' and what was apparently the foundation for the stories of 'wild men' in these mountains. He writes, 'unquestionably, the figure in outline was exactly like a human being, walking upright and stopping occasionally to uproot or pull at some dwarf rhododendron bushes'. It wore no clothes as far as he could see. The footprints 'were very similar in shape to those of a man, but only six to seven inches long by four inches wide at the broadest part of the foot. The marks of five distinct toes and of the instep

were perfectly clear; but the trace of the heel was indistinct and the little that could be seen of it appeared to narrow down to a point. I counted fifteen such footprints at regular intervals ranging from one and a half to two feet. The prints were undoubtedly of a biped, the order of the spoor having no characteristic whatever of any imaginable quadruped'. It is unfortunate that something more could not have been done to solve this mystery.

The book is beautifully got up and illustrated with a wonderful series of views, the actual contact prints being pasted in the book. In many cases the view-point from which the photos were taken is not given. This and an absence of dates in the narrative are two points which might, we think, be considered should the author contemplate a large edition or a future account of this kind. It seems a pity that a book of such merit should be limited to an edition of only 150 copies. The tables at the end of the book will prove useful but the map is rather indistinct.

F. M.B.

'LIFE OF PLANTS', BY SIR FREDERICK KEEBLE (Clarendon Science Series) xii+256 pages. Oxford, at the Clarendon Press, 1926.

'The growth of Science,' says the author in his preface, 'is like that of a tree. From seed-growing to harvest the time is long and tedious to those who look for fruits. But during that time there go on obscurely and below as well as above ground unending successions of very wonderful developments. Not until they have been accomplished are blossom and fruit produced, and then profusion and beauty are apparent for all to see. Yet to those who tend them the science and the tree are beautiful always, alike in their obscure beginnings and in their brilliant fulfilment.—To show this has been my object: to suggest that science is more than a body of doctrine—an illumination of life.'

Keeble, with the modesty of a true scientist, is afraid that he has not succeeded, for he thinks that the task has proved beyond his powers. I cannot agree with him. His book is certainly a successful attempt to impart understanding of the method and to convey impressions of the beauty of Science. I do not want to imply that Keeble has spoken the final word on the subject. He must realize himself, much better than I do, that we are still far removed from the time when we shall be able to give an adequate picture of the life of plants. And even when science will have succeeded in listing all the manifestations of life, it does not follow that by doing so, we have also established the internal connections between, and causes of, those manifestations.

Linnaeus says somewhere in his *Philosophia botanica*: 'Principium erit mirari omnia, etiam tritissima. Medium est calamo committere visa et utilia. Finis erit naturam accuratius delineare quam alius.' Keeble seems to have followed these words, if not in the composition of his book, at least in preparing its substance. The volume scarcely contains anything that you could not find in many other publications. It deals with the ordinary facts of plant-physiology: the manufacture of food-materials, the absorption of radiant energy by the green leaf, the mobilization of food-materials for construction and power-production, the environment of the land plant, variation, heredity, evolution, etc. And still, this book differs in many ways from others of a similar size and aim. Every page offers a wealth of well-defined information; the reader feels that it has been written by a hand that has felt the pulse of the living plant for many years; he knows that the author could tell him much more than he does and that what he offers you is only a condensed abstract of a vast accumulation of facts. You are not drowned in a sea of ill-digested and badly arranged details. The author is always conscious of what he wants his book to be—an 'illumination of life'. There are botanists in India who write on the same subject and with the same aim, viz., to shed light on the mysteries of plant-life. But they have been induced to dispel the darkness by exaggerating the actual state of our knowledge of certain problems, and by not distinguishing critically between facts and products of imagination. Keeble is not of this type. To him a fact is a fact, a hypothesis is a hypothesis, an attempt to explain a certain process remains an attempt, and no plausible explanation, if it is not supported by solid

data, will induce him to call that attempt a scientific explanation. To give only one example. There is that much-discussed problem of the ascent of sap. Some say they have solved it, and they have found the key to solution in a very simple experiment, and we cannot help wondering why none of the workers before them made use of the same magical wand. Keeble is more modest, and he is modest because he is a true scientist. The problem of the ascent of sap, he says, 'has not even now received an answer which completely satisfies every one who has investigated it.' And after three pages of tentative explanations he is not ashamed to confess that 'the study of transpiration leads from the clear domain of the certain to the dimly uncertain, and beyond to the unknown, the darkness of which only future research may enlighten.'

Keeble's *Life of Plants* is a book that every educated man will read with interest and profit. It is a sober, reliable, and honest account, stimulating at the same time and apt to widen the intellectual horizon without causing mental indigestion.

E. B.

CORRECTION

The note on a 'Panther with Abnormal Feet' which appeared on page 909 of Volume XXX of the Society's Journal was the contribution of His Highness Maharao Khengarji of Kutch, G.C.S.I., G.C.I.E., and not of Maharaj Kumar Shree Vijayarajji as published.

Editors.

EDITORIAL

Our President—His Excellency Sir Leslie Wilson—referred at some length in his speech at the Annual Meeting to the work the Society proposed to do in training teachers in Government Schools to become in their Schools efficient teachers of Nature Study or, as we prefer to call it, Natural History.

He expressed his regret that the knife of retrenchment prevented Government budgeting this year for any financial grant to the Society for the purpose, but gave vent to the hope that some public-spirited citizen of Bombay would help the Society by subscribing funds sufficient for the first year, during which the Education Department would study the work done, and, if it were considered likely to succeed, then there would be strong probabilities of Government budgeting in the future for the expense of maintaining the scheme.

There are many public-spirited men still living in Bombay but so far as the needs of the Natural History Society are concerned the vast majority seem to be suffering from an inability to find their way to the pocket in their trousers where their purse is kept. No doubt, like the small boy when first put into knickers, they have put their hands so often into their pockets that, to protect themselves from perhaps having no pockets in the future, they have had their pockets sewn up. There is one public-spirited gentleman in Bombay who has determined that so far as help in educational matters is concerned his purse pocket should never be closed. We refer to Sir Sassoon David, Bart., who has established a Trust and a Committee of Trustees to deal with that Trust, whereby the interest on a large sum of money shall be available for educational needs of proved utility.

Having a good cause we approached the Trustees and we have now the pleasant duty of expressing our thanks to the Trustees for their right appreciation of the situation and to Sir Sassoon David our gratitude for his princely generosity. We have received from the Trustees a cheque for Rs. 6,000 and have informed the Director of Public Instruction that we are prepared to depute our Assistant Curator, Mr. Salim A. Ali, to be Guide, Philosopher and Friend to all those at the Government Training College, etc., who wish to learn how to make the wonders of Nature legible and intelligible to children of all races in our schools in the Bombay Presidency.

We owe our thanks too not merely to the munificence and foresight of the living but also to the munificence and foresight of those who have passed away but whose name liveth for ever, and surely the name of N. M. Wadia, whose magnificent charity has benefited so many and been on such a Catholic scale, will never be forgotten in Bombay. The Trustees of the Wadia Charities recognizing that besides the lecturer and his salary, apparatus, etc., would be necessary, have given us Rs. 750 and we are very grateful to them. This is not the first time they have helped us. In April 1913 they gave us Rs. 500, in September 1923 Rs. 500, and in 1921 Rs. 1,000 for the Mammal Survey of India, Burma and Ceylon.

We are confident that members of the Society will cordially approve this fresh activity of the Society. We do not exist purely for our own benefit, but to spread the knowledge of the Fauna and Flora of India and so benefit others.

For ourselves, we are more than glad to see again in the Journal an article by Mr. Stuart Baker on the Game Birds of India and to welcome as frontispiece a coloured illustration by Mr. Grönvöld. These illustrations are expensive but we do not think they are luxuries. They are appreciated by members and are therefore of benefit to the Society. They are appreciated by non-members and so are an advertisement to the Society.

We say they are appreciated by members and we know they are, but we wish members would show their appreciation in a practical manner.

There are to be five volumes of the Game Bird series. Two have already been published both in the Journal and in book form. Two have been published in this Journal—'Pheasants' (vol. xxiii to vol. xxvii) and 'Partridges and Quails' (vol. xxvii to vol. xxx) and the fifth and last volume. 'The Waders and other Semi-Sporting Birds' is now being published in the Journal but the publication of this volume in book form and the issue in book form of the preceding volumes iii and iv depends entirely on the appreciation members show of the two volumes now available in book form. We published 1,500 copies of the second edition of volume i and 1,500 copies of the first edition of volume ii. 1,000 copies of No. 1 and 1,100 copies of No. 2 have been bound. We have on the lowest estimate 1,400 members. Yet

in five years we have only sold a total of 512 copies of volume i and 671 copies of volume ii. Members have not supported the Society by subscribing for the volumes as they should have done, yet members can obtain the volumes at special reduced rates and no member who has bought the volumes has considered that Rs. 52 for volume i and Rs. 44 for volume ii is anything but a very cheap price. Even where a member is not interested in small Game Shooting, he must be extraordinarily unfortunate if he has not got a friend who is and who appreciates a good gift. The gift of these volumes does indeed bless both him who gives as well as him who receives. Every book purchased releases for the needs of the Society a much wanted sum of money. The reserve fund of the Society is invested in these volumes and until they are sold that reserve is of no help to us. We wish for a moment that the Society was run on the lines of a trade union and that we could use peaceful picketing against every member, every Officers' Mess and every Club in India which does not possess the volumes. Will members make a special effort and order? Let them remember their sisters and their cousins and their aunts, their nephews or some other youngster who would enjoy life in India better if he had a good knowledge of the Game Birds of this country and Burma, and, remembering, fill up the order form which he will find loose in this Journal.

We said farewell on the 1st June to the Curator, Mr. S. H. Prater, who left Bombay on deputation to America. Mr. Prater intended to spend a few days in Central Europe visiting Museums and, after a brief stay in England, will reach New York about 15th July. He will return to Bombay at the end of December 1926 and we look forward with interest to the reforms and innovations which will result—his return will however not be an unmixed blessing to the Honorary Secretary. Will he have learnt in the land of the Almighty Dollar to make bricks without straw or will it be a duet of 'They do it like this in America and we must do the same.' 'It can't be done, Prater, there is no money.' Will there ever be an occasional chorus 'Well the Maharaja of ————— became a Vice-Patron this morning, you can mount your group of Elephants'. It's the occasional chorus we want and with His Excellency the Viceroy and His Royal Highness the Prince of Wales as Patrons surely some of India's Princes who are members of the Society might become Vice-Patrons and give us Rs. 5,000 to commemorate their name in the Prince of Wales' Museum.

Since writing the above we have received, through the kindness of H. H. the Maharaja of Rewah, the skin of a white tiger. Unfortunately it was shot at the hottest time of the year and in a distant jungle and its condition causes great anxiety to our Taxidermists. Even should they be unable to preserve the skin so as to mount the trophy in our Museum—and what a splendid trophy would it not be!—yet the possession of the skin of a white tiger in our research collections is something to be very grateful for, and we are sure all members of our Society will join in our expression of thanks to His Highness.

Major Colvin writes us in connection with the shooting of the trophy as follows. 'It was shot at the hottest time of the year in a distant jungle on the borders of Rewah and the Korea State. Two other white tigresses were seen in beats by His Highness in the same vicinity, but were spared. In all three white tigers have been shot by His Highness, two others (tigresses) have been seen recently and one (a tiger) died in captivity here in 1925 during the hot weather.'

He adds, 'If you mount the head, the eyes should be very pale, almost "wall-eyed" colour. There is no pink about them at all.'

For further particulars about white tigers in Rewah State and for a photograph of one in captivity (evidently the one which died in 1925) we would refer members to an article on the subject written by the Editors and published on page 932 of volume xxvii of this Journal.

As regards the occurrence of a Black Tiger we would refer members to Mr. T. A. Hauxwell's note on the 'Possible occurrence of a Black Tiger' which appeared on page 788, vol. xxii, Major Stewart Capper's note 'Black Tigers', page 343, vol. xxiii, and to General R. G. Burton's note on page 216 of this volume.

We have much pleasure in announcing that Miss Comber has presented to the Society the valuable collection of Hymenoptera made by her father, the late Edward Comber, when in India. We also congratulate Col. R. W. Burton on his recovery from the severe injuries he received from the fall from his Machan. We look forward to his return to India and shikar in November 1927.

OBITUARY

MR. S. BASIL—EDWARDES

We regret to announce the death on the 2nd December, 1925, of Mr. S. Basil Edwardes, a rising young Naturalist.

Mr. Basil-Edwardes was a member of the British Ornithologists' Union and joined this Society on the 8th February, 1918, ever since which time he has been a fairly regular contributor to our Journal.

He was a keen observer of Nature and we have a number of extremely well written notes from his pen in our Journal.

Mr. Basil-Edwardes *forte*, however, was Ornithology, for which branch of Natural History he displayed great aptitude and still greater promise.

Some of his contributions are :

On the Homing Flight of the Common House Crow (*Corvus splendens*),
vol. xxviii.

The Nidification of the Himalayan Tree-Creeper (*Certhia himalayana*),
vol. xxix.

'Notes on Birds uncommon in or unrecorded from the Simla Hills'.
vol. xxxi.

The first part of his latest paper 'A Contribution to the Ornithology of Delhi,' appears elsewhere in this number.

We, who have recently had the pleasure of handling a collection of birds made by Mr. Basil-Edwardes, and others who were acquainted with his work, realize to the full the loss that Indian Ornithology has sustained through his untimely death.

H. H. SIR UDAJI RAO MAHARAJA OF DHAR

The Society has sustained another loss in the death on 30th July after a prolonged illness of His Highness Sir Udaji Rao Puar Sahib Bahadur, Maharaja of Dhar, K.C.S.I., K.C.V.O., K.B.E.

His Highness was an enthusiastic member of the Society which he joined in 1904. He was an ardent Sportsman and took a very intelligent interest in Natural History. He paid a great deal of attention to the problem of bird migration and lately the Society had the privilege of supplying him with a large number of identification rings such as are in use at a number of Bird observation stations in Europe and America. Should any member ever happen to shoot a ringed duck or other Game Bird bearing the inscription 'Inform Maharaja Dhar' he may at once put it down to the naturalistic zeal of this enlightened ruler.

The following articles written by H. H. Sir Udaji Rao Puar have appeared in our Journal :—

'Comments on the colour change of the Black Buck *Antelope cervicapra*',
vol. xxix.

'Records with Snipe', vol. xxx.

To Her Highness the Maharani of Dhar, who is one of our esteemed Vice-Patrons, the Society tenders its respectful condolences.

MISCELLANEOUS NOTES

No. I.—MEASUREMENTS OF GAME ANIMALS

In the course of many visits to museums, and after inspecting hundreds of mounted heads of game animals, it has been evident that there are certain faults in mounting animals which seem to be common to most taxidermists.

1. **BODY.** The sternum is not carried far enough back, so that the animal appears to have a very thin and elongated waist. The buttocks are usually made globular, no indication of the two very distinct muscles being made.

2. **LEGS.** The hind legs are often badly 'cow-hocked' while both fore and hind legs are not far enough apart.

3. **NECK.** This is usually much too slender. Animals with large horns have thick necks to carry them.

4. **MUZZLE.** The nostrils are not carried back at a slant but are square with the mouth. This is due to the fact that in skinning out, the cartilage of the nose is divided to remove the small muscles between the nostrils, and the taxidermist does not join it up again. The upper lip should project beyond the lower lip, and the sides of the upper lips slightly overlap the lower lips. Many taxidermists seem to forget that a lot of small muscles and cartilage are skinned out, and they place the skin of the lips direct on to the bones, thus shortening down the muzzle and making it too small for the rest of the face, giving it a sort of pug-shaped appearance.

It will be seen from 1, 2 and 3 that there is a general tendency to make an animal too thin; with, I suppose, the object of giving it a graceful, traditional 'gazelle-like' appearance. Ibex and markhor are heavy, bulky animals, while even black-buck and chinkara have sturdy necks and bodies.

I append a list of measurements of specimens of big game of sixteen species. The girths of shoulder and thigh were taken at the junction with the body. Heights at shoulder were taken by straightening the fore legs with the heels together and taking the measurement from the top of the shoulder to the bottom of the heels. The waist measurement was only taken when normal: I have deliberately omitted several measurements taken when the animal was distended with grass at the end of a morning feed. I may say that such increase of girth may amount to 5 inches in the case of quite a small animal like an oorial.

From this list, measurements of the following animals seem worthy of special note:—

1. *Kashmir Stag and Sambhar.*—Although the height at the shoulder of the sambhar is slightly greater, it will be noticed what a very much bulkier animal is the Kashmir Stag. I have shot several specimens of both and had expected to find that the Kashmir stag (undoubtedly I thought the bigger beast) would be a couple of inches more at the shoulder. The length of tail in the sambhar is noticeable.

2. *The Female Serow* (No. 12) is quite abnormal. As is seen she was bigger than the big male which carried the record horns for Kashmir.

3. *The Astor Markhor* (No. 14) and the *Straight-horned Markhor* (No. 16). The measurements of the Haramosh specimen are measurements of a male perhaps slightly above the average, those of the *Straight-horned* are of an adult male of normal size, but not quite as big as one of the three I have shot in previous years.

The measurements indicate a very different type of beast to the Haramosh specimen, and I think the *Straight-horned Markhor* of this area should rank as a separate species. The horns are very distinct from those of the Sufed Koh and Baluchistan, and I have never seen any variation from type, while the geographical distribution is limited and distinct.

The Sind Ibex (No. 20).—The measurements are much smaller than those usually given in books. The buck is about the same size as an oorial ram, though thicker legged and more cobby in build. The height of three other bucks averaged the same.

[illegible]

Cannon bone	DEPTHS							Locality where shot	Remarks
	Thigh	Hock	Throat	Base of neck	Chest	Waist	Weight in pounds		
...	Khru, Kashmir	Horns 47".
...	Kishtwar
5½	22	10	7½	10	17½	15	...	Khru, Kashmir	Probably above the average.
6	31	12	...	14½	21½	Jaipur State
...	Kishtwar	26½" at quarters.
3	15½	6½	6½	7½	11½	10	...	Patiala State
...	Hissar
2½	11	5¼	5½	6½	9	8¼	...	Patiala State
...	43	Ladakh
..	41	„
...	Kashmir Valley
...	„	An abnormally large female.
...	„
5¾	22¼	9	7½	9½	18½	17	...	Haramosh
4¼	17	6¾	„
4	17	7¼	6½	9	14¼	13¾	...	Kohat District
...	196	Baltistan
...	193	„
...	188	„
4½	19	6¾	5	8	14¼	13½	...	Khirthar Range, Sind.
...	131	Ladakh	An exceptionally big ram.
...	118	„
...	125	„
...	243	„
3¾	19	7	Kala Chitta Range, Punjab.
3¾	18¼	6¾	6	10½	12½	10½	...	„ „
3	14	5¾	4	6¼	19½	9½	...	„ „
...	Western Siam	Exceptionally fine horns.
...	Upper Burma

Skin Measurement of Big Game

The following table of measurements of big game skins may be of interest in view of much recent controversy :—

Species	Length straight.	Length on curves.	(Skin)	
			Pegged out.	Dressed.
	ft. in.	ft. in.	ft. in.	ft. in.
Tiger	9 3	9 9½	10 10½	10 3
Tiger ♀	8 3	8 8½	10 0	9 3
Panther	7 1¾	7 5½	8 11½	8 8½
Himalayan Black Bear	4 10		6 1	5 6
"	5 4½		6 10½	6 6½
" ♀	4 10		6 1	5 8
"	5 3		6 8	6 4½
" ♀	5 2		6 7½	6 2½
Brown Bear	4 10		5 10¾	5 5
"	5 3		6 9½	6 4
"	5 5		6 11	6 4½
" ♀	5 1		6 6	6 0

It will be noticed how constant the difference is between the various skins. I have always aimed at securing a good breadth to a skin, so it is possible the lengths of the skins might have been slightly increased, but this would render even less probable the stories of gigantic tigers of 11½ and 12 feet 'as they lay'. I have yet to see a 12 feet dressed tiger skin.

C. H. STOCKLEY,

KOHAT,

March 2, 1926.

Major,

2/1, Punjab Regt.

No.II.—COMPARATIVE SHAPE AND MEASUREMENT OF THE FORE FEET IN TUSKERS AND TUSKLESS ELEPHANTS

There is a statement in Mr. E. L. Walker's *Elephant Hunting and Shooting in Ceylon*, p. 73, to the effect that a Ceylon Tusker elephant has a slightly oval forefoot, similar to, though not so pronounced as, the hind foot.

I have personally investigated this point; and of four captive tuskers' feet which I have examined I find that this is actually so to some extent.

Tuskers being rare in Ceylon I have only been able so far to examine four.

I find, however, that there is a difference of approximately an inch in the back-to-front as against the breadth measurement in Ceylon non-tuskers' feet also, and that in the case of the *tuskers* examined the difference is 2 to 2½ inches.

I should be very much obliged if you can inform me whether any difference has been noted in the shape of the forefoot as regards tuskers and tuskless elephants in India as the point is an interesting one.

EHELIYAGODA, CEYLON,

June 24, 1925.

I. L. CAMERON.

[Commenting on the above Mr. A. J. W. Milroy, I.F.S., Deputy Conservator of Forests, Gauhati, writes :—

'I have never taken any measurements to prove that tuskers' forefeet are 'rather more elongated than those of Mukhanas, because I have always 'accepted it as a fact.

'Experienced natives will generally state without hesitation whether a track 'in the forest was made by a tusker or a Mukhana, but of course one only 'occasionally actually comes on the animal itself so they may often be wrong. 'However, they have had so many generations of experience in the tracking 'of elephants turned loose to graze and so many opportunities of seeing the 'tracks of elephants that have raided their crops, that the probability is that 'they are not mistaken.

'Personally, given a distinct track, I have always been accustomed uncon- 'sciously to ascribe it to a tusker or a Mukhana, as the case might be, but 'I cannot remember to have proved definitely that I was right, though it has 'never occurred to me till the present moment that there could be any mistake


'in most cases. I will forward your letter to Mr. N. L. Bor, who will be 'conducting Kheddahs on a big scale next season, and will ask him to take 'measurements.'

Mr. Milroy has forwarded us the following letter from Mr. N. L. Bor, Deputy Conservator of Forests, Kohima, Naga Hills, Assam, in reference to the subject:—


'I made some measurements last cold weather and append results. It will 'be seen that the front-and-back measurement of the feet of tusker and tuskless 'is greater than the breadth in every case.

'I do not think a study of the figures would justify one in laying down a 'definite rule as to whether the difference in the two measurements is *greater* 'in the case of *tuskers*.

'I think it is a misnomer to apply the term oval to a forefoot though the 'term can be applied correctly to hind feet. The greatest breadth of a forefoot 'occurs well behind the centre of the foot to back axis.'



HIND FOOT



FORE FOOT

Mr. Bor has appended a list of measurements of the forefeet of twenty-nine tuskers and thirty-five tuskless males (*Mukhanas*) the average difference between the back-to-front (A-B) and breadth measurements (C-D) in the tuskers is 1.56", in the *Mukhanas* 1.45". The forefoot in the largest tusker measured 1' 7 $\frac{1}{4}$ ", back-to-front with a breadth of 1' 5 $\frac{1}{2}$ " the same measurements in the largest *Mukhana* were 1' 6" by 1' 2". Eds.]

No. III.—A WONDERFUL SIGHT

Late in the afternoon of a day in January last I witnessed a wonderful sight from my observation machan over the Dod Sampagi tank in the jungle here. At 5 p.m. twenty-eight Elephants came into the tank and for nearly two hours enjoyed themselves thoroughly in the water. The biggest of the herd—an old cow—went through the most extraordinary antics, at times she would sit up like a dog and from this position she would frequently go through a motion of half standing on her head as if she was considering turning a somersault. Most of them were at one time or another completely immersed in the water except for their trunks. Two 'half-grown tuskers sparred with each other for a while. There were two young ones in the herd, one not more than three feet high and I should say about two or three weeks old; the other was about four feet high—both ran in and out between their mother's legs and the smaller one seemed to spend its time stumbling into holes made by its elders' ponderous feet and making frantic efforts to scramble out of them again. One of the herd, a small cow, had a dislocated hind leg, its hip-joint being out. This caused it to walk like a ship pitching and tossing. The poor brute's leg was as a result of the dislocation much longer than the other three and to drag it out of the mud the elephant had to lean right forward. The leg seemed to be very swollen. An elephant Khedda was held on these hills last December and I think it is likely that this elephant was one of those caught, and that it pulled its leg out of joint in its efforts to release itself from its leg-ropes, and was consequently released. This is of course only a surmise. On reaching

one end of the tank the herd got our wind and bunched together and then hurried away into the surrounding jungle on hearing a low whistle from me.

There is nothing more fascinating than observing Game in this tank on a moon-light night. My observation machan is large enough to allow four or five people to lie on it at full length. I recently watched a fine stag Sambhur and two does romp and wallow in the water for about three hours.

HONNAMETTI ESTATE, ATTIKAN P.O.,

via MYSORE,

April 28, 1926.

R. C. MORRIS.

NO. IV.—TIGERS AND ELEPHANTS

When out shooting the other day, I came across the tracks of a small herd of elephants in the bed of a stream. By the tracks it was clear that there was a young calf with the herd, and when I saw the tracks of a tiger following those of the elephants, I suggested to my Burmese hunter, a very experienced old man, who has lived all his life in the jungles, and shot quite a lot himself, that the tiger was following the elephants in the hope of having the calf for dinner. He seemed amused at my crediting the tiger with so much audacity, and told me that the tiger was merely following in order to eat the excrement of the calf. He explained that the diet of the calf was milk, and that tigers looked upon the excrement as a *bonne bouche*. The idea is a nasty one, but from the point of view of natural history, it is an interesting thing in connection with the habits of tigers. I have never read of this anywhere, and it would be interesting to know whether any other members of the Society have ever heard of it.

The tracks of the elephants were a day old, and there were droppings of the adult members along the tracks, but strangely enough, there were none of the calf's. In the hope of finding some I followed the tracks for some miles, coming on places where the herd had stopped to feed for a considerable time, but no droppings were found. Personally, though no naturalist, I am inclined to believe the hunter's explanation. At any rate it is reasonable enough.

I wonder if you could get the views of some of those of the members who are qualified to give opinions on this matter?

THAYETMYO, BURMA,

February 18, 1926.

R. K. ANDERSON.

[Commenting on the above, Mr. S. F. Hopwood, Conservator of Forests, Chindwin Circle, writes:—

'I was very interested in the letter from the Superintendent of Excise, Thayetmyo. Elephants are frequently killed by tigers in Burma though it is very rare that a full-grown animal is so killed. This year several tame half-grown elephants have been killed by tigers in the Mu Division and I hear that a large firm of timber contractors lost a full-grown female from the same cause. There is nothing unusual in tigers killing elephants. What I consider to be much more unusual is the suggestion that a tiger could follow the tracks in the way a tracker tracks game. I do not believe that a tiger is capable of running any animal by scent like a dog or by following tracks like a tracker. Mr. Anderson does not say whether the tiger tracks followed the elephant tracks for some miles. I find it very difficult to believe that they do unless the elephants were changing their feeding ground and had marched for some miles along the bed of a large stream or along a road as they sometimes do. It is well known that tigers habitually use the beds of streams and roads and in my opinion the fact that tiger tracks followed those of the elephants was merely a coincidence.'

In vol. vii, p. 119 of the Journal, we published a note by Mr. A. G. Corbett, Deputy Conservator of Forests, Burma, in which three casualties among elephants due to attacks from tigers are recorded—the first was that of a two-year old elephant calf which was attacked and killed. The mother had come to

the rescue but was unable to do anything and only got badly mauled about the hind quarters and was apparently driven off. The second case mentioned is that of a contractor's elephant which escaped after a severe mauling, and the third case that of a very big tusker which was similarly attacked and succumbed five days later. Most authorities agree that tigers do not find their prey by scent but depend rather on sight and hearing. Eds.]

No. V.—THE 'SAMBHUR' CALL OF TIGERS

May I add still another letter to the many which have appeared on this subject, not as an authority but as one who has many times heard this call during well over twenty years of almost continuous touring and residence in the jungly and out-of-the way parts of S. India and Burma.

My own view has long been that held and stated by the writer in the last Journal (Dunbar Brander), viz., that the cry is (if not actually a sex-call), a call to a mate, and that it is never a 'hunting' cry, or one of alarm pure and simple.

In the letter quoted from (Cardwell), the statement is made that 'no one who has heard it a few times could possibly mistake it with the . . . note . . . of sambhur—but my experience is that on hearing one solitary cry when one is not on the *qui vive* it is generally impossible to say off-hand *which* it is, but that once alert and listening for it the next and subsequent calls can seldom be mistaken. One's mental 'phonetic picture' of sounds depends largely upon the personal equation but I don't like the 'fook' or 'koi' spelling for any calls I have heard; the sound to me is best expressed as something between 'moop' and 'meep' (the initial consonant is probably really entirely absent), and if one tries to pronounce 'moop' with the lips pursed as for whistling, one gets, to my ear, a very fair imitation of the sound as I hear it, although of course it is quite possible that different animals, in different districts, may have a modified pronunciation, as it were.

In Lower Burma I have listened almost nightly for several consecutive nights, during the months of February to April, on different occasions, to the call at quite close quarters, and the impression given was almost as if it were of 'conversational' character, the call being quietly uttered perhaps seven or eight times while the animal was passing over say three or four hundred yards of ground, and sometimes much less, close behind the bungalow in bamboo forest.

Personally I don't think I have ever heard a call replied to, and certainly not from a different direction, though possibly sometimes they might have been more or less alternate calls by two animals in the same radial direction from one and not very far apart. On this point I can offer no evidence but the calls have always *sounded* to be by the same animal repeating its cry. Examination has never shown evidence of two tigers though here again at that season and in such ground the tracks of even one are often difficult to trace.

These nights of calling never coincided with a kill in the vicinity although over twenty kills of dudh-wallahs cattle within a couple of years took place less than a mile away.

The domestic cat has a habit sometimes of wandering about giving vent to a plaintive call or cry which in some ways is distinctly similar to the tiger cry on a small scale, and I have wondered whether perhaps the call in both cases is not so much a definite mate call as a sub-conscious, questing, complaint from an indefinite sense of loneliness; and I am inclined to think, on very slight grounds perhaps, that it is generally made by the female.

By a coincidence I was listening to and thinking over this matter of the calling of tiger, in Lower Tennasserim, two days ago and on the following day received my mail containing the December number of the Journal with the Note under discussion.

On one point I am perfectly satisfied myself, and that is that the call is not ever made as a decoy call in *imitation* of that of the sambhur, as with their acute sense of hearing it is incredible that a note or cry which is capable of recognition by such an imperfect instrument as the human ear should ever *deceive* a sambhur; although the curiosity of the deer tribe might possibly tempt an inquisitive animal to investigate the curious sound, just as one can call up barking deer with a blade of grass or bamboo held between thumbs and blown through. But as I have said, I have never known of a call being heard and followed by the find of a sambhur (or other) kill, which is negative evidence

certainly, but is pertinent; nor have I heard of anyone else having noticed kill and call as being in *post* or *proper hoc* relation.

MERGUI, BURMA,

February 5, 1926.

W. R. COLERIDGE BEADON.

NO. VI.—WILD DOGS IN MYSORE

I have never heard of more than one species of wild dog in India, the rufous-coloured animal, but a few days ago I came across at any rate, a variant from the usual type and should like to ascertain what authorities have to say about it.

I was on bison tracks with an experienced Curumba tracker in ordinary teak forest near the Mysore frontier, when I noticed a movement in the long grass about seventy yards off across a small ravine. I was using field glasses and, pointing out the spot to the Curumba, we saw what I at first thought was a large jackal, but it stood higher and had an independent, fierce look that no jackal ever possessed. Colour was yellowish, almost golden in the sunlight (about 7.30 a.m.) on head and shoulders, greyish-yellow body, full brush deepening into black towards the end. The Curumba whispered *Chennai*, wild dog. I was doubtful as the colour was so different to the ordinary wild dog. We crossed the ravine, the animal running up the hill and disappearing, and heard a series of chattering barks, which the Curumba said were made by these dogs. He was quite right and I saw several more in front, not more than thirty yards off, leaping up in the grass to see us better and now and again giving their extraordinary bark; I don't know what else to call it. I regret I did not shoot one or more but the bison tracks turned off to the right and I followed them.

Subsequently I gathered the following information:—

These dogs have been known to the Curumbas in the Mudumalai Forest for years and are much dreaded by them, as they are said to attack men. The Curumba does not mind the ordinary 'red dog', which is very common eight or ten miles from Mudumalai, nor does he fear tigers, etc., the ways and habits of which he understands. Asked if he knew of a case of a man being attacked, the tracker said that anyone attacked did not come back to tell about it. These dogs are reputed not to allow the ordinary wild dog in the same forest. I asked the Forest Ranger, a very intelligent man, if he had heard of more than one kind of wild dog and he said that he had been told of these by the Curumbas, but had never seen one himself and was interested to hear that I had done so. As to the colour, I have never before seen a wild dog with a black end to his brush in South India, but have read of this in North India; the lighter colour may be due to these dogs being purely forest animals and growing longer coats? Would a cross with a Mysore wolf be a possibility? I think the account of their fierceness may be accepted. The Curumbas are a jungle tribe, living amongst and thoroughly accustomed to all wild animals; good trackers and shikaries, and the man with me was a veteran whose coolness I can vouch for. As far as I know, the ordinary red wild dog does not bark; the only noises I have heard him make are a sort of whistling, apparently to communicate with each other, and occasionally whining.

WELLINGTON, S. INDIA,

March 15, 1926.

G. E. WINDLE,

Captain.

[In colouring the Wild Dog varies from uniform red to rufous-grey or even light brownish-grey. As regards the colour of the terminal portion of the tail—of a series of thirty-four skins obtained in India and Burma all except four have black tips to the tail including specimens from South India and Canara. Of the four white tipped examples three were obtained in the C.P. and one in the U.P. Dunbar Brander in his *Wild Animals in Central India* mentions that the white tip is much more common in the C.P. than the black. Of ten specimens in our collection obtained in the Berars and the C.P. seven have black tips and three white. It is interesting to note that, like the Curumbas of Mysore, the Nagas of the Mishmi Hills, Assam, appear to believe in two varieties of wild dogs, one of which they credit with fiercer and more rapacious qualities than the other. There are, notwithstanding, only two recognized

species of Wild Dogs, the Indian *C. dukhenensis* and the Malay *C. rutilans*. The latter has never been recorded from Indian limits. It would be of very great value if sportsmen obtained skins of the two species alleged to co-exist in Mysore and South India. Eds.]

No. VII.—SOLITARIES

Writers on Indian Big Game Shooting have in the past generally led one to believe that solitary elephants and bison have become 'solitary' through being turned out of a herd. From careful observation during the last five or six years I am quite convinced that this is not the case. They turn 'solitaries' of their own free will: they arrive at an age when, for many reasons, the herd is a nuisance to them, and they get morose and desire to be alone. If it was a fact that younger bulls in the herd turned them out, this surely would apply to the rutting season more than any other period. Yet it is not denied that all solitary bulls do rejoin the herds during this season and they leave the herds as soon as this period is over. In December 1924, I watched a tussle between a magnificent old Bull Bison, undoubtedly a solitary out of the rutting season, and a younger herd bull. The former won hands down! It is inconceivable that, veritable giants of the forest as solitary elephants and Bison are, with their enormous size and strength, they should allow themselves to be turned out of a herd by smaller and less powerful bulls.

December 23, 1925.

R. C. MORRIS.

[Commenting on the above note Mr. Dunbar Brander writes:—

"Writers on Indian Big Game Shooting as a rule are intent on telling the public what they did and how they did it: and although it is generally thought necessary to throw in a certain amount of information about the animal this information is frequently misleading and incorrect.

"To contradict all the misleading information supplied by writers on Big Game Shooting will prove an endless task. I am in no position to speak with any authority on Elephants. With regard to Bison I consider that the great mass of bull bison are solitary by their own choice and I am in general agreement with what Mr. Morris writes.

"The remarks made in 'Wild Animals in Central India' on this subject are as follows:—What Mr. Morris has written is a confirmation thereof.

"At this season (the breeding season) bulls which are solitary during the rest 'of the year join the herd.' 'Young herd bulls will be found with the cows at all times, but during the breeding season the master bulls, who lead semi-solitary 'lives drive these youngsters away. These master bulls however are of no great 'age. The really old beasts lead solitary lives, and this animal seems to lose 'all sexual instincts at a comparatively early age.'

"It will be seen from the above that to a great extent the bulls are solitary by preference. The exceptions to this are the young bulls driven out by the master bulls during the breeding season. These young bulls do not like leading solitary lives, they are seldom met with and when excluded from the herd can often be seen grazing in the vicinity—*near* but not *in* the herd. Then again it must sometimes happen that two master bulls contend for the same herd in which case one must be driven out and unless he can find another herd he has to remain solitary only during this period involuntarily. With regard to the really old beasts, I have often known these make no attempt to join with the herd during the breeding season: the master bulls which join the herd are of no great age. I was camped for some time on a stream in Chanda during the breeding season: there were four old solitary bulls on the banks of the stream which never attempted to join the herds. The same thing has been observed in other places.

"In connection with this subject the habits of the Domestic Bull are of some interest. Stock breeders are chary of purchasing a bull unless fairly young as they cannot otherwise be certain of the bull breeding. It is not uncommon for the purchaser of a bull of say only five years of age, insisting on a guarantee that the bull will perform his duties. It is exceedingly rare for a bull of over ten years of age to be used for breeding purposes: there are reasons, it is true, besides the loss of breeding instinct which account for this, but the loss of the desire to breed on the part of the bull is a big factor in the case. No animal that I know of is comparable with the genus *Bos* in this

respect: in fact the species of this genus with which I am familiar seem to lose the desire to breed long before they can be considered sterile in other respects.

"From the above and from what I wrote in 'Wild Animals in Central India' it will be seen that my observations and Mr. Morris' agree and the only possible difference is that I distinguish between a 'master' bull and an 'old' bull. Mr. Morris refers to a 'magnificent old bull': this beast is probably what I would define as a 'master' bull." Eds.]

NO. VIII.—A SOLITARY COW GAUR

(With a photo.)

While after Bison on September 27, 1923, in the forest round Eindayaza in the Tavoy District (Lower Burma) we picked up the spoor of a solitary animal, a patriarchal old bull as it seemed to us at the time. We followed the trail for a couple of hours meandering in and out of the thicket, in loops and semi-circles in the manner typical of the gaur. On getting into dense bamboo cover at about midday, the dogs flushed our quarry from almost under our feet, but all that we could see of him was glimpses of black as he rushed madly past. A hasty shot point blank told, and fifty yards further with the help of the dogs we came upon the animal, who tried to rise at our approach but was spared



the effort by a second bullet in the neck behind the ear. A closer view of the animal, sad and surprising to relate, disclosed the fact that it was a venerable old cow. The horns, a photograph of which I append, are distinctly of the bull's formation, though more slender and with a somewhat narrow sweep. They are in perfect condition with fourteen annulations on each, all heavily corrugated.

The animal measured 5' 9" at the shoulder, between pegs, and appeared to be in sound healthy condition.

I do not know if it is quite usual to find solitary cows in this manner, but the Karens with me who were keen hunters and trackers of life-long experience, assured me that they had never met with a case of this sort before. It would be interesting to know if any of our members have come across a similar instance, and how this perversion of nature in the old lady may have been brought about.

BOMBAY,

SALIM A. ALI.

January 29, 1926,

No. IX.—A CROSS BETWEEN AN IBEX AND A TAME GOAT

(With a photo.)

The photograph of this curious hybrid which was caught when young in the Kishtwar Hills when amongst a flock of Ibex, is very interesting. The horns are exactly like those of the Sind Wild Goat but are thicker and not so much curved backwards at the tips.

The colouring is much like that of the Baltistan Ibex in its summer coat. The legs of this hybrid are nearly white and the sides are whitish and dark slate.

The photograph was taken in the autumn whilst the Chenar leaves were still green on the trees, as depicted on the print enclosed.

Now in December the hair, which is very coarse, is much longer.



The animal is very sturdy and has the habits of an Ibex, it feeds on leaves, and takes dry hay and has now learnt to eat sugar beet, cabbages and occasionally takes turnips. When I first got it as a present from one of the Kishtwar Wazeers it was very wild and used to raise itself on its hind legs and attack any one who came its way.

SRINAGAR, KASHMIR,

January 1926,

A. E. WARD,

Col. (Retd.)

No. X.—THE NILGIRI TAHR (*HEMITRAGUS HYLOCRIUS*)

(With a plate.)

This interesting wild goat has a very circumscribed habitat and its range is limited to the cliffs and precipices of the Western Ghats, from the Nilgiris southwards. On the Nilgiris themselves it had become almost extinct, though in recent years strict preservation has considerably increased its numbers. The Nilgiri heads, however, run very small compared to those on the more southerly ranges. On the Nelliampathy Hills, to which range my remarks apply exclusively, herds of 60 to 90 still occur and heads up to 16" are to be got. The record buck tapes 17½" and doe 14".

The Nilgiri Tahr spends its life on the open cliffs, and the feats of agility it is capable of across the face of the most terrifying precipices must be seen to be believed. A herd will gallop across or down a cliff which does not appear to offer foothold to a lizard. Where grassy downs top the cliffs, the herds will often lie upon them but seldom wander far across them and jungle is disliked intensely. It is amusing to watch a herd crossing a patch of forest, a feat which seems to be considered a most perilous adventure judging by the careful, wary progress through it. With a knowledge of these habits it is curious that an identical species should be found on both sides of the fifteen miles of flat level plain known as the Palghat Gap. What could have induced the first emigration from the southern ranges across such a great area of what then must have been forest?

'Ibex', as these animals are generally misnamed in the south, are restless animals. Though, generally speaking, they feed in the early morning and late afternoon and lie up during the heat of the day, yet, even at noon, one individual after another will rise from its siesta, walk a few paces, browse a while and sink down again to rest, while the kids are eternally playing, butting each other, frisking about and generally refusing 'to lie still and go to sleep'. While a herd is supposed to be resting, one or more does will be noticed as having posted themselves as sentries and extraordinarily effective watchers they are against any danger coming from below or from their own level, but an enemy approaching from well above them has them at his mercy. I have seen a panther, and on another occasion a tiger, watching a herd graze up towards his ambush. Once they got within reach: one spring and the herd was scattered leaving one of their number behind. Similarly, provided the sportsman is well above his quarry, he can fire several shots at his leisure (should his first miss) before the ibex decides where to run to. Once he does start however, it is at incredible speed considering the ground.

Nilgiri Tahr may occasionally have two young at a birth but one is the rule. Kids are dropped most commonly in the early hot weather. During the heavy rains of the South-West Monsoon the tahr generally move down the cliffs to a lower elevation, ascending to the higher cliffs again with fine weather.

Old bucks desert the herds in the hot weather as a rule, and live a bachelor existence. In fact they are found more often solitary than accompanying the herds. In colour the does are light grey and the bucks, as they age, assume a very deep brown, almost black, coat with the well-known saddle on the back almost white: from a distance indeed this distinctive mark looks perfectly white. The bucks, which stand about 3½ feet high at the shoulder, are far heavier and stockier animals than the does, and except in the case of very young males, their build alone is sufficient to differentiate the sexes. The eye is bright golden-yellow.

Nilgiri Tahr shooting is absurdly simple provided that the sportsman approaches the animals from *above* and that he has a good head for heights. The ground they live in provides the only thrills in this branch of 'Shikar' but these thrills are very real and far from free from danger. A man who has *not* got a really good head for heights and who is likely to become unnerved at having to clamber down an almost perpendicular rock-face by his fingers and toes, with the world opening out like an aeroplane photograph 3,000 feet sheer below him had better try for other game. This is no exaggeration as any one who knows the Nelliampathy Ibex country will certify.

Panthers levy the heaviest toll on the herds but tigers not unfrequently visit the cliffs on a dinner of ibex intent, which sounds unlikely but is nevertheless true.



A GOOD HEAD IN THE SOCIETY'S COLLECTION, R. = $16\frac{1}{2}$ " ; L. = $16\frac{1}{4}$ "



YOUNG FEMALE NILGIRI TAHR

Wild dogs do not stand much of a chance of a successful hunt on this ground and I doubt if any tahr fall to them on these hills.

The Poacher accounts for a good many head on the lower and more accessible cliffs, the meat, locally supposed to possess medicinal value, being the attraction.

The flesh of the doe is delicious but the intolerable smell of the buck would turn any but the most hardened stomach from any gastronomical experiments in that direction.

One uncomfortable habit (from the shikari's point of view) frequently indulged in by saddle-backs, unless shot dead, is to hurl themselves into space in their death struggle, and the trophy when retrieved after its 3,000 feet fall is thereby apt to be ruined.

Young tahr make delightful and most amusing pets, but as they grow up their horns, in play or anger, are capable of inflicting a serious wound. A friend of mine has a tame doe which recently neatly skewered his cook through the thigh, causing a nasty sore which took a long course of hospital treatment to cure.

NELLIAMPATHY,

A. P. KINLOCH, F.Z.S., M.B.O.U.

February 17, 1926.

NO. XI.—THE 'CLICKING' OF THE MUNTJAC (*M. VAGINALIS*)

In volume xxx, No. 3, page 694 of the Journal, Mr. Ralph C. Morris refers to the rattling or clicking noise heard by himself and others, from Barking Deer. I have had several opportunities of hearing a kind of rapping sound from this animal, before being able to find out how it was produced.

One day I was sitting up in a place in the Ghat Forests near Lonavla, where I knew Barking Deer had the habit of moving. It was during the hot weather, and I must have been dozing, when I was roused by a rapping just behind me. I was sitting on a large stone, with my back to the trunk of the tree, large enough to hide me completely, as long as I kept quiet. As I soon recognized the sound as coming from 'Bhekar' I waited, in the expectation that the animal would slowly stalk away through the undergrowth and give me the chance of a shot. However, he was putting my patience to a trial and went on rapping, at intervals, evidently without moving. The sound was such as might have been produced by knocking together two dry sticks, once or several times. After a while I could not master my curiosity any longer. Bending slowly to the left and turning my head backward I could get a good view of the Deer. His head was slightly turned away from me, so he did not notice me. And now I could clearly see that the rapping was produced by the Deer striking the ground (or, more likely, a root or a stone) sharply with the hoof of his right foreleg. Evidently, he had not advanced since he started his performance, nor did he budge from the place during the short time during which I could observe him. I was in a very uncomfortable position, and when trying to improve it I must have frightened the animal, which bounded off with a sharp bark, which he repeated when he came to a stop in the dense undergrowth, perhaps thirty yards away. As he evidently had not seen me I resumed my former position and kept still. After about a quarter of an hour the animal approached again, this time a little from the left, and I could get an easy shot at him.

What prompts the Barking Deer to rap? It may be curiosity or alarm. Barking Deer usually move through the Forests in pairs, at some distance, and the experience I have given an account of above, might justify the conclusion that the animal scented danger and wanted to warn his mate.

As Mr. Morris calls the noise a clicking or rattling it is possible that he refers to a sound which is different from the one I have tried to describe. I may, however, say that the rapping sound will vary according to the ground, or object on the ground the animal's hoof strikes upon.

DELHI,

MAX FREI,

February 15, 1926.

No. XII.—MEASUREMENTS AND WEIGHT OF A MALAY TAPIR

A recent copy of a report on the post mortem examination of a Malay Tapir which died in the Zoo at Karachi gives the measurements and weight of the animal as follows:—

Length	84 inches.
Height at shoulder	44 "
Girth	68 "
Weight	1,200 lbs.

The post mortem examination revealed that the animal died of Gastro-enteritis and a considerable quantity of sand and small pebbles were found in the small intestines.

BOMBAY NATURAL HISTORY SOCIETY,

S. H. PRATER, c.m.z.s.

February 2, 1926.

No. XIII.—THE OCCURRENCE OF THE PINK-FOOTED GOOSE (*ANAS BRACHYRYNCHUS*) AND THE MALLARD (*ANAS BOSCAS*) IN JAIPUR, RAJPUTANA

The occurrence of the following birds in Rajputana may be of interest, particularly as I see Stuart Baker mentions the Mallard as being scarce in Rajputana and the Pink-footed Goose has occurred only sparingly anywhere:—

On a jhil in Jaipur territory, the following were shot on December 27:—

One Pink-footed Goose (*A. brachyrynchus*). (The only one seen, a single bird came to one of the butts.)

Three Mallard (*A. boscas*). (Large number seen, at least fifty on the jhil.)

The Bar-headed Goose was there in thousands.

I also noticed the following birds:—

Common Flamingo (*Ph. ruber antiquorum*). One flock of six. (Two adult in pink plumage, four immature white plumage).

White Pelican (*P. onocrotalus*). Flock of ten.

Dalmatian Pelican (*P. crispus*). Two only.

AJMER, RAJPUTANA,

W. M. LOGAN HOME,

December 31, 1925.

Major.

No. XIV.—ON THE OCCURRENCE OF PALLAS' SANDGROUSE (*SYRRHAPTES PARADOXURUS*) WITHIN INDIAN LIMITS

(With a plate.)

An example of a female Pallas' Sandgrouse (*Syrrhaptes paradoxurus*) was forwarded to the Society by H. H. Shree Sadul Singhji Bahadur, Maharaj Kumar of Bikaner. It was shot at Gujner, Bikaner State, on December 31, 1924 and was identified by Mr. Stuart Baker, who very kindly arranged with the British Museum to provide the photo illustrating the note. Dresser in his *Palaearctic Birds* states that Pallas' Sandgrouse inhabits the steppes of Southern Russia and Asia, east to North China, north to Lake Balk; he writes, 'Large flocks have visited Europe at uncertain intervals and it has been obtained in almost every country, while it has bred in Great Britain and Denmark'. The present is, as far as we are aware, the first authentic record of its occurrence in India. Like the Tibetan Sandgrouse (*Syrrhaptes tibetanus*), Pallas' Sandgrouse has completely feathered toes and tarsi; while the hallux or hind toe is entirely wanting—it can be distinguished from its Tibetan congener by the possession of a large black patch on the abdomen. In habits it is said to resemble other Sandgrouse. The food consists of seeds. The call note which is uttered on the wing is described by Dresser as a loud *truck turuck truck turuck*. The nest is a mere depression in the soil sometimes lined with a few grass bents, and the eggs, three in number, are deposited late in May or early in June; in colour they are stone-buff, often with a greenish tint, marked with purplish-brown shell-blotches and dark-brown surface spots. They measure about 1.69×1.16 (Dresser).

BOMBAY NATURAL HISTORY SOCIETY,

S. H. PRATER, c.m.z.s.

April 15, 1926.



PALLAS' THREE-TOED SANDGROUSE (*Syrhaptes paradoxurus*).

No. XV.—THE OCCURRENCE OF THE GIANT HERON (*ARDEA GOLIATH*) IN THE KHULNA DISTRICT, BENGAL

Blanford writing on the distribution of this Heron states that its occurrence in India is remarkable and somewhat mysterious. He mentions a few records of its having been obtained in the Calcutta Bazaar (1845-1846) by Blyth, of two shot in 1878 and another in 1879 and a third seen there in 1880. It is interesting to record that a Giant Heron (*Ardea goliath*) was presented to the Society in December 1925 by Mr. L. R. Fawcus. It was shot by him in the Khulna Sunderbuns, Bengal, where according to Mr. Fawcus it is not 'uncommon'. The following are the measurements:—

Bill from angle of forehead feathers, 205 mm.; wing, 580 mm.; tarsus, 235 mm.

BOMBAY NATURAL HISTORY SOCIETY.

S. H. PRATER, C.M.Z.S.

No. XVI.—THE ROCK HORNED OWL IN KASHMIR

(With a photo.)

This fine owl, which is nearly as large as the Eagle Owl of Europe, is not uncommon in the valley of Kashmir frequenting steep rocky hillsides between 5,000 and 6,000 feet altitude.

They spend the hours of daylight sitting on some rock on a more or less precipitous rocky hillside protected from the direct rays of the sun. About half an hour after sunset they begin to get on the move, and to call. Their call is a loud 'Hoo'—slightly drawn out, but still a single note and not disyllabic as stated in the F. B. I. Birds.

This call is also heard frequently in the very early dawn.



The food of this owl judging from the large number of 'casts' I have examined in Kashmir, consists almost exclusively of rats, and perhaps five per cent of the

musk rat or rather musk shrew. Breeding commences early in March, holes or sheltered cavities among rocks on steep or precipitous hillsides being used as nesting sites.

Three is the full complement of eggs laid which are broad ovals, pure white of course, measuring about 60×46 millimetres.

The chief enemy of this owl during incubation is the jungle crow. I was fortunate enough to catch one red-handed—so to speak. He had just perforated one egg out of three when disturbed by me. In another case a single egg, the first laid was similarly destroyed. The birds use the same nesting sites year after year. The enclosed photograph of three young owls in their nest on the Takht-i-Suliman (Srinagar, Kashmir) was taken on May 6, when the young were about three weeks old.

The remains of two large rats may be made out to the right and left in the foreground.

SRINAGAR, KASHMIR,
February 1926.

B. B. OSMATSON.

NO. XVII.—MATING HABITS OF THE COMMON KITE (*MILVUS MIGRANS GOVINDA*)

The following are some notes from my diary with regard to the breeding of our Common Pariah Kite (*Milvus migrans govinda*) which might prove interesting. The observations mainly extend over the breeding season of 1925-1926 which still continues.

In the preceding season (1924-25) the first date on which a pair were seen in copulation was October 11, and the last February 2.

In the current season I have been enabled to note happenings more carefully as, close to my window at Chowpati, there are a number of cocoanut palms which are extensively patronised by the kite fraternity of the neighbourhood for nesting purposes. There is one pair in particular which has appropriated a particular palm nearest my window, and whose operations I am going to record. I do not find any reason to doubt the identity of this pair because, as far as I have been able to observe, pairs take up definite quarters, for instance one pair is always to be seen near a particular spot on the ridge of Wilson College roof, another has established itself on a palm farther away from my window, and so on.

Copulation was first noticed on September 13. On December 29, the following note appears in my diary: 'Copulation among the kites continues generally. The act is performed many times each day. Have not been able to determine so far a full day's number, but by 10 in the morning, before leaving for office, have frequently observed it from three to six times. The birds usually return to the same neighbourhood for the act time after time, and very often the female does not leave her perch in the interval at all. The male bird alights on her back from the air, and balancing himself by flapping his wings, flies off directly the act is completed. The female whistles in a particular manner when wishing to attract her mate, and the cock bird keeps up a series of distinctive short, sharp, crescendo whistles or 'screams' which end up abruptly when the act is over.'

On January 21, I noticed this pair busying themselves with nest construction for the first time, but copulation appeared to continue simultaneously with this. On January 31, I thought a marked decline in the number of times per day was noticeable, and this I ascribed to the increase in the actual nesting activities; in this however, I was deceived, for on February 6, my diary has the following: 'Copulation still continues as before. Usually the act takes place three or four times of a morning' and another note on the seventh says, 'The pair were seen to copulate seven times to-day; five times before noon and twice after 5.30 p.m.' This was a Sunday and I was thus enabled to keep a full day's record. On February 13, copulation was observed at 6.45 p.m. just as dusk was setting in. It is February 18 to-day, and copulation still continues though the frequency is somewhat diminished, I think, and the birds at last seem more preoccupied with nest building.

It is astonishing to find this extraordinary protraction of the mating period. Moreover, from September 13 (when the pair were first seen copulating) until

January 21, or say the middle of January, the birds appeared to give no thought to the building of a nest, and it would seem that the idea of bringing up a family had never occurred to them all this time.

The astounding sexual capacity of the birds and the period over which the mating operations continue, surpasses anything known to me in this line with undomesticated animals, and the capacity of the lady kite in particular tends rather to put one in mind of a notorious Queen of Aragon who had certain very original notions on the subject.

SALIM A. ALI.

BOMBAY NATURAL HISTORY SOCIETY,
6, APOLLO STREET,
February 18, 1926.

[Commenting on the above Mr. C. H. Donald writes:—

"The note, I think, is of considerable interest in showing how very often these birds can copulate without apparently any result. Of course, it is not uncommon to see these birds going through all the actions at odd times but whether they actually do the deed, or not, is very difficult to see.

"I am afraid I can shed no light on the subject as I have never taken any special note of it, but I cannot see how they can go on copulating half through the year without producing any result, unless there is something wrong with one of them. I think this note would be of greater value if the writer could continue his observations a little longer and let you know the result, i.e., if they succeed in bringing up a family in the end."

Mr. H. Whistler considered the incident as abnormal. He writes, 'I have not come across such a case, and it would not normally occur, as the organs in a male bird are usually in a quiescent state and merely enlarge temporarily at the beginning of the breeding season. Presumably pairing does not take place ever until the organs have enlarged and stimulate the male to action.'

The question arises however, as to whether the breeding season had not already commenced when these observations were made. In Bombay the normal breeding season commences early in October.

Asked to continue his note in the light of further observations, Mr. Salim Ali writes on May 16th as follows:—

"I give below notes from my diary in regard to the copulation of the pair of kites subsequent to the 18th of February:

Feb. 21 "Copulation has become markedly less frequent."

Feb. 24 "Pairing has practically ceased, though it does still take place occasionally."

Feb. 28 "Witnessed pairing. This has now become very infrequent.

"After the above I have no further record, and the mating would therefore appear to have ceased from this date. The pair now took up the work of collecting material for the nest in right earnest. Building commenced in the same palm and a good deal of 'screaming' (quite distinct from the 'screaming' in copulation) accompanied the operations. Crows evidently proved a great nuisance and it was a common sight to see one of these rascals being assailed and chased away from the vicinity by the infuriated owners.

"Unfortunately, before much progress had been made with the nest, the whole thing was pulled down by a toddy-drawer who climbed the tree regularly, and further attempts of the birds met with the same fate. After three or four misadventures in this manner, the pair left the locality and I did not notice them making further endeavours. They were still casually to be seen in the neighbourhood, but for nesting they probably found a safer haven elsewhere. In fact, I am not certain if they eventually did succeed in bringing up a family at all. One particular I omitted to mention in my previous note was that during the act, and while the male was on the female's back and emitting the short sharp "screams" to which I have referred, the female very often turned her head back to look at him, bringing her bill quite close to his. I was reminded of this by a note I was reading the other day on the "Courtship and Sexual Habits of Birds" (*Ibis*, Series xii. vol. i (1925), p. 889) where writing of the Oyster Catcher (*Hæmatopus ostrelegus*) J. S. Huxley observes as follows:

"During the time the cock is on the hen's back, she may turn her head right

round to look at him, and on one occasion I thought their bills touched." ' This remarkable similarity in the behaviour of birds of such widely divergent species is interesting indeed.

"Referring to the period which may elapse between the actual union and the laying of the eggs, Huxley observes the following on p. 893 which I quote *in extenso*, to show that the facts recorded by me, are not altogether abnormal, although it must be admitted that the period during which copulation was carried on by the kites was a very considerable one.

"In Warblers buntings, etc., " "The hen appears to pass through two distinct phases—one in which she seeks the territory and company of the male, the second in which she will copulate. This separation of phases does not occur in the Oyster Catcher, although it is clear that unions achieved in March can have no physiological effect on the fertilization of eggs not laid till after April 25, i.e., in Oyster Catchers the physiological capacity of the eggs to be fertilized is not synchronous with the physiological readiness for insemination, while in Buntings the two are closely linked—an interesting difference. I have evidence that the Crested Grebe resembles the Oyster Catcher in this respect. . . . It had best be styled the 'pre-incubation period' ". Eds.]

NO. XVIII.—BRAHMINY KITE (*HALIASTUR INDUS INDUS*) SWIMMING

Three or four Brahminy Kites, all young birds, were flying over a small dock situated opposite my house picking up refuse floating in the water, when suddenly one of them in making a swoop flopped down on the water. It seemed in no way disconcerted but rode the water easily, swimming like a duck. After the space of a minute or so, it rose rather awkwardly and rejoined its companions.

Mr. H. B. Tilden, with whom I discussed the incident, has made a similar observation and has sent me the following note:—

'On a voyage from Karachi to Bombay per s.s. *Chakla*, while anchored off the port of Porbandar on the 21st instant, I observed a Brahminy Kite (which had been hovering round the ship in company with several sea gulls) take to the water gracefully and swim in quite a choppy sea for at least five minutes. He then rose into the air with the ease and grace of a sea gull.

It occurs to me that this is an unusual proceeding especially as I saw a Common Pariah Kite (*Milvus govinda*) drowned in Madras harbour. In stopping for food thrown overboard the s.s. *Modasa*, the kite struck a wavelet and apparently could not take to the air again. She swam or rather beat the surface of the water with her wings for quite half an hour and progressed about 50 yards to the ship's side.

We passed a stout rope, with knotted loops, overboard but the unfortunate kite was too exhausted to take hold and just as a lascar was about to climb down to the rescue the kite sank.'

BOMBAY NATURAL HISTORY SOCIETY,
6. APOLLO STREET,
May 1, 1926.

S. H. PRATER, C.M.Z.S.

NO. XIX.—AN ALBINO BUSTARD (*EUPODITIS EDWARDSI*)

On January 30, 1926, at a spot about eighteen miles, N.W. of Kutch Mandvi, on an open and barren tract of country, I saw a flock of nine bustard. One of these was a pure white—apparently as white as an egret—and was clearly visible from a long distance, long before the others were distinct. I tried for a long time to get within range but owing to the openness of the ground and the number of the flock, I could not do so. I should be glad to know if such albinos have been noticed before. I myself had never heard of nor seen one till this occasion.

BHUJ CUTCH.

VIJAYARAJJI.

[In January this year we received from the Maharaja of Dhar an Albino Coot (*Fulica atra atra*). Albinism was not complete and there was a considerable mixture of black feathers among the white which gave the bird a speckled

appearance. The feet and bill were fleshy white and the iris pale blue. In complete albinos the pupil and iris are red owing to the blood vessels shining through the otherwise strongly pigmented parts. When albinism is partial it may mean a general deficiency in the pigment available or a disturbance in the normal blood supply of a certain area of growing feathers. A lesion of the pulp of a growing feather not infrequently prevents the deposition of the normally present black pigment in the feathers and other parts, but the pulp may recover after one or more moults and normal colouration be assumed. Eds.]


NO. XX.—THE TRANSPORT OF BIRDS' EGGS

(With a text figure.)

Carrying eggs about India, where one, so to speak, moves with one's house on one's back, is not an easy matter, so possibly the method I employed may be of use to others, who may be experiencing trouble; anyhow I give it for what it is worth.

In 1895, I was quartered at Baroda, where I had the privilege of making the acquaintance of Professor Harold Littledale. Though probably unknown to the present generation, he was then one of the foremost sportsmen in the land, in every way. Like myself, he collected eggs and initiated me into the method I now write of and which I have found most successful. Should this article, by any chance, come to his notice, I hope he will forgive me for writing about his inventions.

To commence with, get an ordinary kerosine tin: have one end removed and converted into a lid, on a hinge, and a clasp to close it. Then, have trays constructed of tin with lids (either all hinges or removable) to fit into the kerosine tin easily, so that they can be drawn out without force. These trays should be made in three *depths*. For big eggs, three trays to a tin, for medium size four trays, for small eggs five trays. A small loop or handle should be made to each set of trays to facilitate withdrawal. Each kerosine tin would, also, have a handle to carry it by.



Kerosine Tin containing
3 trays suitable for big eggs

Any tinsmith, in any bazaar, can make these things without difficulty. Having got your tin with trays ready, cotton wool or wadding should be placed at the bottom of each tray. In these the eggs, for transportation, should be wrapped up, separately, in wool and packed, while, on the top of all, another layer of wadding should be placed. Eggs, thus packed, will travel, with the utmost safety.

Eggs can also be shewn off to great advantage, when not travelling, in these trays, by removing them out of their wrappings and placing them in clutches, separated by a little wool, paper, etc.

Another advantage of this system is that one can put one's hand on any clutch in a moment, as follows. Kerosine tins should bear Roman numbers, thus I, II, III, etc. Trays ordinary, thus 1, 2, 3, 4, etc. Eggs should be packed in them by variety and a record kept. For instance, the eggs of *Corvus splendens* are packed in tray 7, box III. A record should be placed against it thus:—III. 7. All that is necessary, should you wish to look up *Corvus splendens* are packed in tray 7, box III. A record should be placed aid is to place a slip at the top, in each tray, showing, by clutches, the eggs it contains, thus *Corvus splendens* 4. 4. 5. 3. *Corvus corax* 4. 3. 3. 5., etc. This also gives a census of all eggs.

To move your tins, wooden boxes to hold 4, 6, 8 or any number to suit your convenience, can be made.

For use in the districts, shooting or other expeditions, the ordinary kerosine wooden case holding two tins is recommended. A lid on hinges, with a fastener is easily made. This is a load that can be easily carried by a coolie or placed on a mule, pony, yak, etc.

I have found the above method very satisfactory. It is simplicity itself. Kerosine tins can be obtained almost anywhere in India, as also the tinsmith and sutár to do the necessary.

JAMAICA, B.W. 1.,

R. M. BETHAM, M.B.O.U.,

NO. XXI.—EARTH SNAKE (*SILYBURA* SP.) AND CHICKEN.

Last month while proceeding, accompanied by my mother and sister, to call on a neighbour, we came across a young fowl, lying helpless in the middle of the road, with an earth snake (the common *Silybura*) wound tightly, in several loops, round both its legs, which were so tightly tied together that it took me several seconds to disentangle the knot and release the bird. Did the chicken take the snake for a particularly juicy worm? But how did the snake get so firm a grip round both legs? And why would it not let go? Had I not had witnesses with me at the time, I should hesitate to record a fact to me so incredible.

NELLIAMPATHY HILLS,

A. P. KINLOCH, F.Z.S., M.B.O.U.

January 1, 1926.

[The power to coil itself tightly round an object has so impressed the natives that in some parts of the Western Ghats, as at Mahableshwar where the Earth Snake (*Silybura macrolepis*) is common, the belief is held that were one of these snakes to coil round one's finger amputation would be the only means of ridding oneself of it! Eds.]

NO. XXII.—AN INTERESTING ENCOUNTER WITH A COBRA

While motoring into Mysore the other night, my search light picked up a snake lying in the middle of the road, which I took to be a Russel's Viper, and I promptly steered the car, so that both the front and the back wheels went over it. I was travelling at some speed but pulling up as soon as possible, I switched over the searchlight to the rear, and backed the car to see what had happened and saw the snake lying still. Imagining the heavy car had killed it, I steered close by when, to my surprise, the snake reared up with expanded hood and struck the wheels in a most vicious manner. I now realized, I was tackling a cobra and an unusually a large one at that! The snake started to move across the road, and I once again went over it, slowly this time, and distinctly felt the front wheel go over, and I am sure, the back followed suit. It did not, however, seem to harm the snake much as, when I looked back it was still moving, so I promptly reversed the car and as I tried to get the back wheel over it, it reared up once more with distended hood and struck repeatedly at the wheel which just missed it; but my front off side wheel caught it squarely as I gave the necessary swing and I felt the wriggling of the snake on the steering wheel, as it appears to have got entangled in the spokes and drawn back a yard or two.

At this critical moment the head lights went out owing to a faulty switch contact, which I had to push into place and just in time to see the cobra get off the road and crawl into the bushes alongside it where I could not follow in the car nor, did I think it advisable to investigate matters further on foot.

One would have fancied that a heavy car of this type, a Six Cylinder Buick, would have crippled, if not killed, the snake; but, beyond appearing a bit sluggish in its movements, it did not seem to have been hurt seriously.

MYSORE,
March 17, 1926.

WILLIAM H. THEOBALD.

No. XXIII.—THE OCCURRENCE OF THE BUTTERFLY *APPIAS*
INDRA VAR *ARISTOXENUS* IN THE NILGIRIS

As it may be of interest to some members of the Society I beg to report the capture in February 1925 in the Nilgiris at about 4,500' of a *O Appias indra* corresponding to the form '*aristoxemus*' figured on page 50 of the plates of 'The Rhopalocera of the Indo-Australian Faunal Region' Seitz, which flies in the mountains of Formosa.

I have also caught a few females of '*Statilia*' which have a kind of earthy brown under surface very similar to the form '*thrasia*' figured on the same page and which also flies in Formosa.

COONNOOR, S.I.,
March 27, 1926.

J. FLORENCE.

[Commenting on the above Col. W. H. Evans writes:—

'*Appias indra shiva*, Swin. is the correct name for the South Indian race of *narendra*, M: the name *Statilia* Fruh, designates the more heavily marked wet season form. Wherever it occurs *indra*, as well as the other members of the genus *Appias*, is extremely variable more particularly in respect of the colour below and the extent of the black margin above. The variations mentioned by Mr. Florence are normal.' Eds.]

No. XXIV.—ON *PSARA PHŒOPTERALIS* AS A PEST ON GRASSES
IN SOUTH INDIA

(With a plate.)

INTRODUCTION

It is highly probable that many of the minor pests the Entomologist is confronted with are kept in their present status by the combined action of various little known factors which serve to bring about what is vaguely termed as the balance of life. At any time any of these minor pests may turn out to be formidable enemies of crops, if by chance the equilibrium is disturbed by the disappearance of one or more of these factors. *Psara phæopteralis* presented a remarkable instance of such a change of status in the year 1924. In November of that year it was noted by the junior writer to be one of the pests concerned in the destruction of fodder grasses in Ernad Taluk, Malabar. It was subsequently found that *Psara phæopteralis* was present also at Coimbatore at that time though not in such large numbers as to cause alarm. Under these circumstances it was felt expedient to study its life-history and binomics and in the present paper a short account of the observations made is presented.

LIFE-HISTORY

The Imago.—This is a Pyralid moth a little over 10 mm. in length and with a wing expanse of 25 to 30 mm. It is dark fuscous in colour with a prominent spot on the forewings. It is very much like *Psara licarsisatus*.

The Egg.—It is creamy white in colour and almost oval in shape. It is moreover somewhat convex in shape being deepest at the centre and thinner towards the margins. The egg is about 0.25 mm. long and 0.2 mm. broad. Eggs 300 to 500 in number are generally laid on the food plants but in captivity the female moth readily laid eggs on the sides of glass jars. Eggs are

generally found in clusters varying from 5 to 25. It takes four to five days for the eggs to hatch. The egg acquires a reddish tint about the middle after a day, making thereby the beginning of development of the embryo. After four days the young larva could be seen lying curled up inside the egg shell. Soon it tries to move about and after travelling round inside the shell many times over it breaks its passage out with the help of the mandibles.

The Just Hatched Larva.—This has its head and prothorax black and the rest of the body pinkish red. The cephalic and the prothoracic shields are markedly well developed. The body bears dorsally and laterally tufts of short fine hairs which arise from warts. The just hatched larva is 1.25 mm. long. After escaping from the egg shell it wanders about and gets attached to a grassblade. It then makes a webby shelter at the base of the leaf blade and feeds by scraping the tissues from the leaf. At this stage it has a curious habit of curling itself like a millipede when disturbed. At the end of three days it undergoes its first moult.

Second Instar.—The larva after the first moult changes from pinkish red to light pink colour. Now it is about 2.5-3 mm. long. The second instar lasts three days.

Third Instar.—The colour of the larva changes from light pink to light green. The larva is 6-8 mm. long. The third instar lasts three days.

Fourth Instar.—The colour changes to green and the warts become thickened. The larva is 14 mm. long. The fourth instar lasts two days.

Fifth Instar.—The larva is dark green in colour. It is 20-22 mm. long. When ready to pupate it is about 25 mm. long and somewhat stout. The hairs are very prominent.

Pupa.—It is dark brown in colour and is 10 mm. long. Pupation takes place on the surface of the soil at the base of the food plants, the cocoon being made of silk and excreta. Pupal period is seven days.

FOODPLANTS

The pest was noted on *Panicum javanicum*, *Panicum miliaceum*, *Andropogon annulatus*, and *Cynodon dactylon*. It is probable it may feed on any common grass. In one instance it was reared from wheat.

DISTRIBUTION

Malabar, Coimbatore, Bangalore and Bellary.

NATURAL ENEMIES

A single specimen of an Ichneumon parasite, *Syzeuctus annulipes* (see Fig. 7) was one reared from a larva.

METHODS OF CONTROL

As the parasite noted is not found in large numbers control by this means is out of the question. The only method of dealing with the pest would appear to be the collection and destruction of adults. Since large numbers of moths were found attracted to electric light at the Agricultural College and Research Institute the possibility of controlling them by means of light traps was tested and the results proved to be fairly satisfactory. The use of hand net was also found efficient in minimising their numbers.

Reference to Plate—

1. *Panicum javanicum* plant showing the various stages of the pest.
2. Egg x 43.
3. Just hatched larva x 31.
4. Mature larva x 3.
5. Pupa x 5.
6. Adult x 3.
7. Larval parasite x 2.

M. C. CHERIAN, B.A., B.S.C., D.I.C.

AND C. J. GEORGE, M.A.



For explanation see end of article.

PROCEEDINGS

Proceedings of the meeting held on June 24, 1926

A meeting of the members of the Bombay Natural History Society and their friends took place on Thursday, at 6.15 p.m., in the Board Room at the Prince of Wales' Museum, Rev. E. Blatter, S.J., Vice-President, presiding.

NEW MEMBERS

The following fifty-seven new members were elected since the last meeting :— Mr. G. Cunningham, C.I.E., O.B.E., I.C.S., Kabul ; Mr. A. Boller, Bombay ; Major W. Brooke-Purdon, D.S.O., M.C., O.B.E., R.A.M.C., Wellington, Nilgiris ; Mr. J. E. Clutterbuck, R.E., Bhurkunda ; The President, Mess Committee, 4th Bn. 10th Baluch Regt. (D.C.O.), Quetta ; The Director, Institute of Plant Industry, Indore, C.I. ; Miss R. Navalkar, Poona ; The Mess President, 4/4th Bombay Grenadiers, Razani, N.W.F.P. ; Mr. Fred Hagenbach, Bombay ; Mr. H. A. Hyde, M.C., Indore, C.I. ; The Curator, Sarawak Museum, Kuching, Sarawak ; Mr. K. H. McIntosh, Mussoorie ; The President, Mess Committee, 4/12th F.F. Regt. (Sikhs), Ambala Cantt. ; Mr. G. H. Farquharson, Misa, Assam ; Mr. J. A. G. Campbell, Bokakhat, Assam ; Mr. K. R. Eates, Karachi ; Mr. J. D. Finlay, Clarkabad, Lahore Dist. ; H. H. the Thakor Saheb Lagdhirsinhji of Morvi, Morvi ; H. H. The Thakor Saheb Bahadursinhji of Palitana, Palitana ; Lt.-Col. Gordon Casserly, Europe ; Mr. F. G. Burgess, Maymyo, Burma ; Mr. T. G. Cule, Lumding, Assam ; Mr. I. C. Dare, Bombay ; The Mess President, Officers' Mess, 3/4th Bombay Grenadiers, Lucknow ; Mr. D. G. Cameron, Mahendragiri, Nagercoil, S. I. ; Mr. D. G. Sevastopulo, F.E.S., Lyallpore ; Mr. Ali Asghar H. Fyze, Bombay ; The Mess Secretary, 3/17th Dogra Regt., Alipore ; Col. St. G. L. Steele, C.B., Abbottabad, N.W.F.P. ; Mr. A. Macleod, I.C.S., Peshawar ; The Mess President, 3/10th Baluch Regt., Lahore Cantt. ; The Mess President, Officers' Mess, Royal Engineers, Roorkee, U.P. ; Mr. C. H. Thompson, I.F.S., Rangoon ; Mr. W. G. Crawford, I.F.S., Rangoon ; H. E. Lord Irwin, P.C., G.M.S.I., G.M.I.E., Viceroy of India, Simla ; Capt. Malik Anun Singh, M.B., I.M.S., Taiping, F.M.S. ; Capt. L. D. W. Hearsey, M.C.V.D., *via* Mailani, Kheri, U.P. ; Mr. A. F. Gibbs, Umaria, B.N.Ry. ; Mr. G. Atkinson, Dehra Dun ; Mr. V. A. N. Sausman, Dango, *via* Billimora ; Mr. J. E. Bisset, Meppadi, P.O.S.I. ; Mr. E. H. Peacock, B.F.S., Mawlaik, Burma ; Mr. F. D. S. Richardson, England ; Mr. F. Gersperger, Bombay ; Mr. F. I. Morgan, Kadur ; Mr. K. Johnson, Nazira, Assam ; Mr. L. G. Knight, Kalthurity, S.I. ; Mr. C. W. Clode, Coorg ; The Agricultural Engineer, Burma, Mandalay ; the Mess Secretary, 1/4th Bombay Grenadiers, Bombay ; the President, Mess Committee, 3/15th Punjab Regt., Allahabad, U.P. ; H. H. Durjansal Singh, Ruler of Khilchipur ; Mr. Duncan T. Merson, Nilgiris ; Capt. V. R. Wilton, I.A.S.C., Dalhousie, Punjab ; Mr. R. L. Gamlen, Hyderabad, Deccan ; Capt. A. R. Spurgin, Bombay ; Mr. C. C. Fowler, M.I.E.E., Bombay.

The Honorary Secretary announced that not only had H. E. Lord Irwin become a member of the Society in his private capacity but he had also, as Viceroy of India, been pleased to become a Patron of the Society and the Society therefore now had the honour of having as their Patrons both His Majesty the King-Emperor's representative in India and His Royal Highness the Prince of Wales, the Heir Apparent to the Throne.

This might be considered another tribute to the work of the Society. Work which was appreciated by those who bore rule in India, for not only was His Excellency the Governor of Bombay a member and President of the Society, but the Governors of Bengal, Madras, Burma, and the United Provinces were also members.

GENEROUS DONATIONS

A hearty vote of thanks was passed to the Trustees of the Sir Sassoon David Trust Fund for a donation of Rs. 6,000 and to the Trustees of the Wadia Charities for a donation of Rs. 750. These donations would enable the Society, in conjunction with the Director of Public Instruction, to carry out the scheme for teaching nature study in schools in this Presidency referred to by H. E. Sir Leslie Wilson at the annual meeting of the Society. Mr. Salim Ali, who had recently been appointed Assistant Curator to the Society, would be appointed the guide, philosopher and friend of those teachers in schools who took the nature study course about to be inaugurated at the Museum.

THE LATE MR. E. COMBER

The Honorary Secretary referred to the great loss sustained by the Society through the recent death of his old friend and colleague Mr. E. Comber. Mr. Comber became a member of the Society in 1894, joining the Committee in 1901. He was keenly interested in Birds and also in Hymenoptera. He contributed several papers to the Society's Journal. By the last mail it was learnt that Mr. Comber's daughter had presented the Society with her father's valuable collection of Hymenoptera.

NEW EXHIBITS AT THE MUSEUM

Sir Reginald Spence drew the attention of the members and those present to the beautiful exhibits which had been added to the various galleries since the last meeting.

The first of these was the magnificent group illustrating the nesting of the Great Indian Hornbill. The material for this case was obtained through the energies of Mr. M. S. Tuggerse, Divisional Forest Officer, Honavar. The tree in which the nest was situated was 150 ft. high. The nest itself was 75 feet from the ground—photographs in the case show the position of the nest quite clearly. The tree was felled and the section containing the nest had to be transported a considerable distance to the coast and then shipped to Bombay. The female and young (which was a little over two weeks old) were despatched to Bombay alive. The two occupy the nest, while 'Old Bill' who lived in the Society's room for twenty-six years has taken on the responsibilities of a father—here he is seen feeding the family through the small opening which is left after the entrance hole has been sealed up. These birds inhabit the evergreen forests of the lower Himalayas, Kanara, and Burma.

Besides this, there was a fine specimen of a Panther which was secured for the Society by Col. R. W. Burton, I.A. (Retd.) and also a Scaly Ant-eater or Pangolin which was obtained by Mr. F. Ludlow from Baluchistan. Several other smaller exhibits were also on view in different galleries all of which had been set up in the Society's laboratory.

Rev. E. Blatter, S.J., then read a paper on the 'Luminescence of Animals and Plants,' a full account of which will appear in the next number of the journal.

**WARD TAXIDERMY FAMOUS FOR
OVER 100 YEARS.**

Revised Edition, Illustrated, Rs. 16.
LYDEKKER'S

**'Game Animals of India, Burma,
Malaya and Tibet.'**

By J. C. DOLLMAN.



**ROWLAND WARD'S
RECORDS OF BIG
GAME**

EIGHTH EDITION
Profusely Illustrated.

Price, Rs. 43-12.

Rowland Ward's Sportsman's Handbook.
Eleventh Edition.

Rs. 3-8.

Increased facilities, and the most modern methods in the ROWLAND WARD STUDIOS, have enabled us to make a considerable reduction in our charges for modelling heads.

New Price List on Application,

ROWLAND WARD, LTD.,

Naturalists,

By Appointment to H. M. The King,
167, Piccadilly, LONDON, W.

**FRANCIS EDWARDS,
Bookseller,**

83a, High Street, Marylebone, London, W. 1.

Catalogues of Books on the Indian Empire, Ornithology and Natural History are issued at Intervals and the latest will be sent post free on application.

**'OUT OF PRINT' BOOKS SEARCHED FOR,
FREE OF CHARGE.**

FOR SALE

Indian Pigeons and Doves by Stuart Baker, in good condition

Rs. 25

B. N. H. S. Journal, Vol. XV-XXIII, Bound in cloth boards and Vol. XXIV-XXX, unbound, complete and in good condition

Rs. 150

Apply

**H. H. F. M. TYLER, Esq., I.C.S.,
Anantapur, M. & S. M. Railway.**

A GUIDE TO SHIKAR ON THE NILGIRIS

BY

'BIG BORE'

Crown 8vo. 270 pages with a map

The latest publication on Shikar dealing almost exclusively with the Nilgiri Mountains. The information given is reliable, up to date and invaluable to Sportsmen :: :: :: ::

The most comprehensive and interesting book on the subject yet published

CONTENTS

BIG GAME AND RECORDS
TABLE OF GAME SHOT during the
past fourteen years
GAME HAUNTS AND SEASONS: Chief
centres and routes
POPULAR SHOOTING CENTRES
SHIKAR HINTS AND NOTES
BEATING FOR TIGER
STANDARD METHOD OF MEASURING
TROPHIES
CAMP ARRANGEMENTS
TRAVELLERS' BUNGALOWS
ELEPHANTS
NILGIRI GAME RULES
FISHING

SHOOTING CENTRES OUTSIDE THE
NILGIRIS
CROCODILE SHOOTING ON THE WEST
COAST
REPTILES
THE LEPIDOPTERA OF THE NILGIRIS
THE COIMBATORE DISTRICT
THE WYNAAD, MALABAR DISTRICT
THE PULNI HILLS
RULES AND REGULATIONS FOR HUNT-
ING AND SHOOTING IN THE
TRAVANCORE STATE
THE SPORTSMAN'S SHIKAR MAP OF
THE NILGIRIS

Price Rs. 4 post free

THE DIOCESAN PRESS

Post Box 455

::

Vepery, Madras.

